

ESTIMATION OF CARCASS FAT AND PROTEIN IN NORTHERN PINTAILS (ANAS ACUTA) DURING SPRING MIGRATION

Authors: Dombrowski, Pascale, Bourgeois, Jean-Claude, Couture, Richard, and Linard, Christian

Source: Journal of Wildlife Diseases, 39(3) : 620-626

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-39.3.620>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

ESTIMATION OF CARCASS FAT AND PROTEIN IN NORTHERN PINTAILS (*ANAS ACUTA*) DURING SPRING MIGRATION

Pascale Dombrowski,¹ Jean-Claude Bourgeois,^{2,3} Richard Couture,¹ and Christian Linard^{4,5}

¹ Département de Chimie-Biologie, Université du Québec à Trois-Rivières, C.P. 500, Trois Rivières, Québec G9A 5H7, Canada

² Société de la faune et des parcs du Québec, Direction de l'aménagement de la faune, 5575, Saint-Joseph, Trois-Rivières-Ouest, Québec G8Z 4L7, Canada

³ Département de Chimie-Biologie, Université du Québec à Trois-Rivières, C.P. 500, Trois Rivières, Québec G9A 5H7, Canada

⁴ Département de Chiropratique, Université du Québec à Trois-Rivières, C.P. 500, Trois Rivières, Québec G9A 5H7, Canada

⁵ Corresponding author (email: christian.linard@uqtr.ca)

ABSTRACT: Foraging in stopover areas influences nutritional condition of birds during spring migration. Our purpose was to determine if body mass, percent carcass water, and serum biochemistry would predict energy reserves (carcass fat and protein) in northern pintails (*Anas acuta*) at a spring staging area, Lake St. Pierre in Québec, Canada (46°11'N, 73°08'W). Northern pintails were collected during spring 1997 (14 April–9 May). In this staging area, body mass and percent body water successfully estimated carcass protein and fat in male northern pintails, but only carcass protein in females. None of the seven blood parameters we used accurately estimated nutritional reserves in staging northern pintails. These findings suggest that investigators must use direct estimates of carcass reserves to examine nutrient reserve requirements for egg production, migration, or body maintenance during spring migration.

Key words: *Anas acuta*, carcass composition, northern pintail, nutritional status, serum chemistry, spring migration.

INTRODUCTION

Nutritional and reproductive status of waterfowl are supported by carcass fat and protein reserves (Whittow, 1986; Raveling and Heitmeyer, 1986; Harder and Kirkpatrick, 1994). This physiologic condition is partly a function of foraging in appropriate habitats (Gauthier et al., 1984; Whyte and Bolen, 1984); therefore spring staging areas may be important to allow birds to build energy reserves during and after migration to breeding areas (Olsen, 1994; Austin and Miller, 1995). Changes in body mass during spring or fall migration are common in many North American bird species (Dunn, 2001). Furthermore, the stopover allows waterfowl to build protein reserves that support egg production (Krapu, 1981; Barzen and Serie, 1990; Alisauskas and Ankney, 1992; Krapu and Reinecke, 1992; Mann and Sedinger, 1993; Esler and Grand, 1994). The use of carcass protein during migration (Marsh, 1984) would necessitate its replenishment during the stopover. Some migrants may gain carcass protein equivalent to a third of their body-

mass gain during a stopover (Karasov and Pinshow, 1998). Proteins may be stored as muscle or as other functional tissue (Klaassen et al., 1997). The date that a bird arrives at each staging area, and how long it stays, may affect carcass nutrient reserves. Furthermore, the previous winter location and the ultimate nesting region could also influence nutrient reserve status (LaGrange and Dinsmore, 1988).

Lipids are the dominant fuel source during migratory flight in many bird species, and carcass fat has generally been determined by petroleum ether lipid extraction (Dobush et al., 1985; Johnson et al., 1985; Ringelman and Szymczak, 1985; Miller, 1989; Gauthier et al., 1992; Harder and Kirkpatrick, 1994; Dabbert et al., 1997). However, lipid extraction is both expensive and time consuming (Johnson et al., 1985). Because of this, easier and less expensive methods have been developed to estimate body nutrient reserves. These methods involve equations or indices based on: 1) morphometric measurements including body mass (Johnson et al., 1985;

Ringelman and Szymczak, 1985; Miller, 1989; Smith et al., 1992; Dabbert et al., 1997); 2) amounts of fat and protein in different tissues (Chappell and Titman, 1983; Whyte and Bolen, 1984; Johnson et al., 1985; Hohman and Taylor, 1986; Miller, 1989; Smith et al., 1992; Dabbert et al., 1997); or 3) water content of the carcass (Child and Marshall, 1970; Campbell and Leatherland, 1980; Johnson et al., 1985; Miller, 1989; Conway et al., 1994; Karasov and Pinshow, 1998). For example, body mass was found to be a good predictor of nutrient reserves in northern pintails from Texas (Smith et al., 1992) and California (Miller, 1989).

Blood biochemical analyses reflecting lipid or protein metabolism seem to offer possibilities for estimating carcass reserves. For example, Ksiazkiewicz et al. (1993) found significant correlations ($P < 0.05$) between cholesterol levels and abdominal fat deposition ($r = 0.48$) and between intestinal fat and cholesterol ($r = 0.39$) in female mallards (*Anas platyrhynchos*). Dabbert et al. (1997) found that blood triglyceride levels predicted ($r = 0.57$; $P = 0.007$) total body fat in mallard drakes. Measurements of plasma total protein, albumin, uric acid, urea, free fatty acids, cholesterol, β -hydroxybutyrate, and thyroid hormones may also give reliable information about nutritional status of birds (Totzke et al., 1999; Gannes, 2001; Hollmén et al., 2001; Alonso-Alvarez et al., 2002).

In this paper, we examined how well body mass, percent body water, and different biochemical compounds predicted carcass fat and protein during spring migration. We expected some blood biochemicals related to lipid metabolism (free fatty acids, triglycerides, cholesterol, and β -hydroxybutyrate) or protein metabolism (total protein, uric acid, albumin) would allow estimation of carcass fat and protein, respectively.

MATERIALS AND METHODS

Study area

We conducted the field study from 14 April to 9 May 1997 at a migratory stopover on the

north shore of Lake St. Pierre (St. Lawrence River) in Québec, Canada (46°11'N, 73°08'W). This stopover (246 ha) consists of 80 agricultural fields (plowed fields, stubble fields [corn, barley, buckwheat, soybean], prairies, and abandoned fields) that are flooded during springtime. It is crossed by five drainage ditches and forms a deep basin. The staging area is managed in order to maintain water levels during spring waterfowl migration. It is part of a Ramsar wetlands site as well as a UNESCO Biosphere Reserve and is heavily used by northern pintails, which represent 80% of the 10,000 dabbling ducks present in spring. The spring staging period, which lasts nearly 5 wk, coincides with a period of high food availability for ducks (seeds and invertebrates). In addition, some female pintails nest in the Lake St. Pierre floodplain as reported by Bélanger and Couture (1989).

Specimen gathering

We collected a total of 36 male and 31 female feeding northern pintails using a .22 rifle. We sexed and weighed (hereafter termed "body mass") each bird, obtained blood samples (5–10 ml) by cardiac puncture (Donham, 1979; Hannon, 1979) with a syringe (22G×1½" needle), and put the blood samples into additive-free glass tubes (Vacutainer®, Becton Dickinson, Rutherford, New Jersey 07070, USA). We retrieved only 30 males and 27 females rapidly enough to obtain an adequate volume of blood. We stored all blood samples at 4 C in the dark until reaching the laboratory. We centrifuged the tubes within 3 hr after their arrival at the laboratory and obtained the sera before freezing them at -20 C pending analysis. We excluded esophagi, gizzards, and ovarian follicles from carcass analysis but stored them for other studies. We distinguished yearlings from adults by first observing differences in the cloacal bursae (bursae of Fabricius), but because this criterion may be not valid during spring for waterfowl (Dane, 1968), we also used wing-feather characteristics that differentiate yearling from adult northern pintails during breeding season (discriminant functions DF and DF2 for males and females, respectively) as proposed by Duncan (1985). Our sample was therefore composed of 47 adults (25 males, 22 females) and 10 yearlings (5 males, 5 females). We stored carcasses at -30 C in plastic bags.

Measurement of body reserves

In the laboratory, we sheared all feathers from the thawed carcasses using electric clippers and scissors. We weighed carcasses again,

TABLE 1. Variables measured on northern pintails sampled at Lake St. Pierre, Québec, in spring 1997.

	Male		Female	
	Adults (<i>n</i> = 25) ^a	Yearlings (<i>n</i> = 5) ^a	Adults (<i>n</i> = 22) ^b	Yearlings (<i>n</i> = 5) ^b
<i>Body components</i>				
Body mass (g)	1115 ± 110 ^c	1110 ± 69 ^c	984 ± 85	912 ± 74
Carcass fat (g)	169.0 ± 61.5	174.8 ± 85.0	148.6 ± 46.4	175.7 ± 55.2
Carcass protein (g)	264.5 ± 80.8	269.7 ± 72.5	247.0 ± 75.6	174.4 ± 20.9
Percent body water	52.74 ± 5.15	51.37 ± 3.34	51.25 ± 4.16	53.83 ± 3.78
<i>Blood metabolites</i>				
Free fatty acids (mmol/l)	1.679 ± 1.120	1.750 ± 1.052	1.465 ± 0.803	2.123 ± 1.428
Albumin (g/l)	17 ± 3	18 ± 3	18 ± 3	15 ± 2
Uric acid (μmol/l)	1039 ± 427	1134 ± 347	1005 ± 393	691 ± 82
Cholesterol (mmol/l)	7.00 ± 1.01	7.57 ± 1.15	6.90 ± 1.26	4.98 ± 0.75
Total protein (g/l)	38.0 ± 4.8	39.1 ± 4.6	43.3 ± 6.2	38.8 ± 0.8
Triglycerides (mmol/l)	4.11 ± 1.18	4.31 ± 0.78	5.78 ± 1.50	5.77 ± 2.18
β-hydroxybutyrate (g/l)	0.074 ± 0.038	0.072 ± 0.028	0.069 ± 0.030	0.040

^a For β-hydroxybutyrate, *n* = 28 males (24 adults, 4 yearlings).

^b For β-hydroxybutyrate, *n* = 22 females (21 adults, 1 yearling).

^c Mean ± SD.

then ground (Hobart meat grinder, Hobart Corporation®, Troy, Ohio, USA) and homogenized them.

We dried a 100 g sample of the homogenate from each northern pintail to constant mass in a forced-air oven (60 °C) to calculate percent body water ("percent water"). We ground each lyophilized sample in a high-speed grinder and weighed 1 g duplicates using an analytical balance. We extracted lipids (total body fat ["carcass fat"]) following Gauthier et al. (1992) using petroleum ether. Finally, we calculated dry lean mass ("carcass protein"): Carcass protein = fat-free dry weight – mineral ash (obtained after incinerating the lyophilized sample at 550 °C for 12 hr) (Miller, 1989; Dabbert et al., 1997).

Biochemical analyses

We examined free fatty acids, triglycerides, cholesterol, and β-hydroxybutyrate, which are blood metabolites related to lipid metabolism and total protein, uric acid, and albumin, which are related to protein metabolism (Amand, 1986; Hochleithner, 1994). These serum factors could potentially be used to predict nutritional status (Ksiazkiewicz et al., 1993; Dabbert et al., 1997; Domingo-Roura et al., 2001). Because we collected variable amounts of blood, not all metabolites could be determined in all individuals (we analyzed samples from only 28 males and 22 females for β-hydroxybutyrate).

We used a Hitachi 704 automatic chemical analyzer (Boehringer Mannheim GmbH, Mannheim, Germany) to perform serum bio-

chemical analyses with the procedures and reagents supplied by the company: albumin (Doumas et al., 1971), total protein (procedure proposed by Boehringer), free fatty acids (Shimizu et al., 1980), cholesterol (Siedel et al., 1983; Kattermann et al., 1984), triglycerides (Siedel et al., 1993), β-hydroxybutyrate (Bergmeyer and Bernt, 1965), and uric acid (procedure proposed by Boehringer). We did not conduct duplicate analyses because serum samples were too small (1–2 ml). All analyses were submitted to internal and external quality control processes (Boehringer Mannheim Canada [now Roche Diagnostics Canada], Laboratoire de santé publique du Québec [LSPQ]). Our laboratory was accredited by the LSPQ.

Statistical analyses

We analyzed all data by sex because of the differences in metabolism and the timing of physiologic events between male and female birds. However, for each sex, we grouped the yearlings and adults together due to the small sample size and because there was no significant difference (Mann-Whitney *U*-test, *P* > 0.05) between yearlings and adults for the different variables used in the analyses, except for cholesterol in females (*P* = 0.003). Table 1 presents the values (for the different variables measured on the male and female birds) separately for yearlings and adults.

We first tested all of the variables for normality and judged all as normal (albumin values were arc-tangent transformed and free fatty ac-

TABLE 2. Correlation values (r) of body mass, percent body water, and serum chemistry variables with carcass fat and carcass protein (g) for 30 male and 27 female northern pintails sampled at Lake St. Pierre, Québec, in spring 1997. Significant values are identified with asterisks (*: $P < 0.05$, **: $P < 0.01$, ***: $P < 0.001$).

	Males		Females	
	Carcass fat	Carcass protein	Carcass fat	Carcass protein
Body mass (g)	0.41	0.70***	0.24	0.58**
Percent water	-0.53*	-0.60**	-0.08	-0.71***
Free fatty acids (mmol/l)	0.14		-0.35	
Cholesterol (mmol/l)	0.16		0.09	
Triglycerides (mmol/l)	-0.29		-0.05	
β -hydroxybutyrate (g/l) ^a	0.35		-0.35	
Albumin (g/l)		0.29		0.42
Uric acid (μ mol/l)		0.44		0.35
Total protein (g/l)		0.04		0.26

^a Analyses of 28 males and 22 females.

ids values were natural-log transformed). We tested all null hypotheses at the $P < 0.05$ significance level. We used the Pearson correlation coefficient to identify the variables (body mass, percent water, or blood metabolite) that best explained carcass fat and carcass protein. We adjusted P values using the Bonferroni method. We used SYSTAT[®] software for all analyses (Geissler, 1988; White and Clark, 1994).

RESULTS

Male northern pintails collected at Lake St. Pierre during spring 1997 weighed (mean \pm SD) $1,114 \pm 103$ g and females weighed 971 ± 86 g (Table 1). Body mass and percent water correlated significantly with carcass protein (for both males and females), but there were no significant relationships between indices and energy reserves except for carcass fat and percent water in males (Table 2). We found no statistically significant or consistent relationships between serum metabolites and carcass fat or carcass protein in either males or females (Table 2).

DISCUSSION

At Lake St. Pierre, carcass protein was consistently predicted by body mass and percent water. Percent water reflected, at least partially, variations in carcass fat in male northern pintails. For females, no relationship was found between carcass fat and percent water; maybe there was not a sufficient range of carcass fat in the sam-

pled birds. Lastly, we must not forget that excluding the digestive and reproductive organs might have biased the analysis. In other studies, body mass was a fair predictor of carcass fat (Miller, 1989 [$r^2 = 0.63$]; Smith et al., 1992 [$r^2 = 0.69$ – 0.74]) and carcass protein (Miller, 1989 [$r^2 = 0.71$]). However, carcass fat was estimated more precisely by percent water (Miller, 1989 [$r^2 = 0.92$ – 0.95]; Conway et al., 1994 [$r^2 = 0.86$]).

None of the seven blood parameters used in our study accurately estimated nutritional reserves in staging northern pintails. Biochemical parameters only partially reflect nutritional status, because factors other than condition influence physiology (Domingo-Roura et al., 2001). For example, age (Puerta et al., 1990), daily rhythms, starvation, stress, and method of capture can alter blood biochemistry profiles (Harder and Kirkpatrick, 1994). Stress can cause an increase in cholesterol (Franzmann and Thorne, 1970) and protein levels (Laidley and Leatherland, 1988; Marco and Lavin, 1999). Triglyceride levels decline with time between the last triglyceride-rich meal and sampling (Domingo-Roura et al., 2001). Levels of blood proteins are affected by metabolic interactions, such as tissue repair, that create demands on protein reserves and, subsequently, decrease albumin in the blood

(Domingo-Roura et al., 2001). Moreover, protein and lipid catabolism during long-distance flight (Jenni and Jenni-Eiermann, 1998) can cause an increase in uric acid, free-fatty acids, and β -hydroxybutyrate (Gannes, 2001; Gannes et al., 2001). Blood metabolite composition is strongly affected by diet (Gannes, 2001) and varies according to age, sex, and reproductive condition (Fairbrother et al., 1990). Adults may have lower plasma uric acid and triglyceride levels than young birds (Puerta et al., 1990). Ksiazkiewicz et al. (1993) found significant correlations between cholesterol levels and fat deposition in females, but they did not find any correlation between blood cholesterol levels and fat deposition in mallard drakes.

Estimating carcass fat and protein may be useful and less costly than complete carcass analyses; however, our results suggest that investigators must use direct estimates of carcass reserves to examine nutrient reserves during spring migration or reproduction. Indeed, the destination and origin of the pintails we collected was unknown, nor did we know how long any individual pintail had been in the area. Pintail spring migration through any given area is very complicated and might affect technical results. Additional research with larger samples from additional spring staging areas, together with an understanding of the dynamics of migration of northern pintails through the areas, is needed.

ACKNOWLEDGMENTS

We were financed by La Fondation Héritage Faune (Fédération québécoise de la faune), La Fondation Universitaire du Centre du Québec, and l'Université du Québec à Trois-Rivières (UQTR). We thank the staff of La Société de la faune et des parcs du Québec for technical support and G. Gauthier (Université Laval) for kindly allowing us to use his equipment for carcass analysis. We appreciate the helpful comments of H. Glemet (UQTR) and G. Gauthier who reviewed a previous version of the manuscript, A. Aubin (UQTR) who helped in study design, and G. Daigle (Consulting Service in Statistics, Université Laval) for his advice on the statistical analysis. We thank the following

people for reviewing the manuscript: L. Devine, B. Duval, M. R. Miller, and an anonymous Journal of Wildlife Diseases reviewer.

LITERATURE CITED

- ALISAUSKAS, R. T., AND C. D. ANKNEY. 1992. The cost of egg laying and its relationship to nutrient reserves in waterfowl. *In* Ecology and management of breeding waterfowl, B. D. J. Batt, A. D. Afton, M. G. Anderson, C. D. Ankney, D. H. Johnson, J. A. Kadlec and G. L. Krapu (eds.). University of Minnesota Press, Minneapolis, Minnesota, pp. 30–61.
- ALONSO-ALVAREZ, C., M. FERRER, AND A. VELANDO. 2002. The plasmatic index of body condition in yellow-legged gulls *Larus cachinnans*: A food-controlled experiment. *Ibis* 144: 147–149.
- AMAND, W. B. 1986. Avian clinical hematology and blood chemistry. *In* Zoo and wild animal medicine, 2nd Edition, M. E. Fowler (ed.). W. B. Saunders, Philadelphia, Pennsylvania, pp. 264–276.
- AUSTIN, J. E., AND M. R. MILLER. 1995. Northern pintail (*Anas acuta*). *In* The birds of North America, No. 163, A. Poole and F. Gill (eds.). The Academy of Natural Sciences, Philadelphia, and The American Ornithologists' Union, Washington, D.C., 32 pp.
- BARZEN, J. A., AND J. R. SERIE. 1990. Nutrient reserve dynamics of breeding canvasbacks. *Auk* 107: 75–85.
- BÉLANGER, L., AND R. COUTURE. 1989. Pintail, *Anas acuta*, use of man-made ponds in Quebec. *Canadian Field-Naturalist* 103: 547–550.
- BERGMAYER, H. U., AND E. BERNT. 1965. Enzymatische Bestimmung von Keton-Körpern im Blut. *Enzymologia Biologica et Clinica* 5: 65–75.
- CAMPBELL, R. R., AND J. F. LEATHERLAND. 1980. Estimating body protein and fat from water content in lesser snow geese. *Journal of Wildlife Management* 44: 438–446.
- CHAPPELL, W. A., AND R. D. TITMAN. 1983. Estimating reserve lipids in greater scaup (*Aythya marila*) and lesser scaup (*A. affinis*). *Canadian Journal of Zoology* 61: 35–38.
- CHILD, G. I., AND S. G. MARSHALL. 1970. A method of estimating carcass fat and fat-free weights in migrant birds from water content of specimens. *Condor* 72: 116–119.
- CONWAY, C. J., W. R. EDDLEMAN, AND K. L. SIMPSON. 1994. Evaluation of lipid indices of the wood thrush. *Condor* 96: 783–790.
- DABBERT, C. B., T. E. MARTIN, AND K. C. POWELL. 1997. Use of body measurements and serum metabolites to estimate the nutritional status of mallards wintering in the Mississippi Alluvial Valley, USA. *Journal of Wildlife Diseases* 33: 57–63.
- DANE, C. W. 1968. Age determination of blue-winged

- teal. *Journal of Wildlife Management* 32: 267–274.
- DOBUSH, G. R., C. D. ANKNEY, AND D. G. KREMENTZ. 1985. The effect of apparatus, extraction time, and solvent type on lipid extractions of snow geese. *Canadian Journal of Zoology* 63: 1917–1920.
- DOMINGO-ROURA, X., C. NEWMAN, F. CALAFELL, AND D. W. MACDONALD. 2001. Blood biochemistry reflects seasonal nutritional and reproductive constraints in the Eurasian badger (*Meles meles*). *Physiological and Biochemical Zoology* 74: 450–460.
- DONHAM, R. S. 1979. Annual cycle of plasma luteinizing hormone and sex hormones in male and female mallards (*Anas platyrhynchos*). *Biology of Reproduction* 21: 1273–1295.
- DOUMAS, B. T., W. WATSON, AND H. G. BIGGS. 1971. Albumin standards and measurement of serum albumin with bromocresol green. *Clinical Chemistry Acta* 31: 87–96.
- DUNCAN, D. C. 1985. Differentiating yearling from adult northern pintails by wing-feather characteristics. *Journal of Wildlife Management* 49: 576–579.
- DUNN, E. H. 2001. Mass change during migration stopover: A comparison of species groups and sites. *Journal of Field Ornithology* 72: 419–432.
- ESLER, D., AND J. B. GRAND. 1994. The role of nutrient reserves for clutch formation by northern pintails in Alaska. *Condor* 96: 422–432.
- FAIRBROTHER, A., M. A. CRAIG, K. WALKER, AND D. O'LOUGHLIN. 1990. Changes in mallard (*Anas platyrhynchos*) serum chemistry due to age, sex, and reproductive condition. *Journal of Wildlife Diseases* 26: 66–67.
- FRANZMANN, A. W., AND E. T. THORNE. 1970. Physiological values in wild bighorn sheep (*Ovis canadensis canadensis*) at capture, after handling, and after captivity. *Journal of the American Medical Association* 157: 647–650.
- GANNES, L. Z. 2001. Comparative fuel use of migrating passerines: Effects of fat stores, migration distance, and diet. *Auk* 118: 665–677.
- , K. A. HATCH, AND B. PINSHOW. 2001. How does time since feeding affect the fuels pigeons use during flight? *Physiological and Biochemical Zoology* 74: 1–10.
- GAUTHIER, G., J. BÉDARD, J. HUOT, AND Y. BÉDARD. 1984. Spring accumulation of fat by greater snow geese in two staging habitats. *Condor* 86: 192–199.
- , J.-F. GIROUX, AND J. BÉDARD. 1992. Dynamics of fat and protein reserves during winter and spring migration in greater snow geese. *Canadian Journal of Zoology* 70: 2077–2087.
- GEISSLER, P. H. 1988. Criteria for evaluating microcomputer statistical packages. U.S. Fish and Wildlife Services Resources Information Bulletin 88-15, 2 pp.
- HANNON, S. J. 1979. Plasma calcium as an indicator of reproductive condition in female blue grouse. *Journal of Wildlife Management* 57: 463–465.
- HARDER, J. D., AND R. L. KIRKPATRICK. 1994. Physiological methods in wildlife research. In *Research and management techniques for wildlife and habitats*, T. H. Bookhout (ed.). The Wildlife Society, Bethesda, Maryland, pp. 275–306.
- HOCHLEITHNER, M. 1994. Biochemistries. In *Avian medicine: Principle and application*, B. W. Ritchie, G. J. Harrison and L. R. Harrison (eds.). Wingers Publishing, Lake Worth, Florida, pp. 223–245.
- HOHMAN, W. L., AND T. S. TAYLOR. 1986. Indices of fat and protein for ring-necked ducks. *Journal of Wildlife Management* 50: 209–211.
- HOLLMÉN, T., J. C. FRANSON, M. HARIO, S. SANKARI, M. KILPI, AND K. LINDSTRÖM. 2001. Use of serum biochemistry to evaluate nutritional status and health of incubating common eiders (*Somateria mollissima*) in Finland. *Physiological and Biochemical Zoology* 74: 333–342.
- JENNI, L., AND S. JENNI-EIERMANN. 1998. Fuel supply and metabolic constraints in migrating birds. *Journal of Avian Biology* 29: 521–528.
- JOHNSON, D. H., G. L. KRAPU, K. J. REINECKE, AND D. G. JORDE. 1985. An evaluation of condition indices for birds. *Journal of Wildlife Management* 49: 569–575.
- KARASOV, V. H., AND B. PINSHOW. 1998. Changes in lean mass and in organs of nutrient assimilation in a long-distance passerine migrant at a springtime stopover site. *Physiological Zoology* 71: 435–448.
- KATTERMANN, R., D. JAWOREK, G. MOLLER, G. ASSMANN, I. BJORKHEM, L. SVENSSON, K. BORNER, G. BOERMA, B. LEIJNSE, AND J. P. DESAGER. 1984. Multicentre study of a new enzymatic method of cholesterol determination. *Journal of Clinical Chemistry and Clinical Biochemistry* 22: 245–251.
- KLAASSEN, M., A. LINDSTRÖM, AND R. ZIJLSTRA. 1997. Composition of fuel stores and digestive limitations to fuel deposition rate in the long-distance migratory thrush nightingale, *Luscinia luscinia*. *Physiological Zoology* 70: 125–133.
- KRAPU, G. L. 1981. The role of nutrient reserves in mallard reproduction. *Auk* 98: 29–38.
- , AND K. J. REINECKE. 1992. Foraging ecology and nutrition. In *Ecology and management of breeding waterfowl*, B. D. J. Batt, A. D. Afton, M. G. Anderson, C. D. Ankney, D. H. Johnson, J. A. Kadlec and G. L. Krapu (eds.). University of Minnesota Press, Minneapolis, Minnesota, pp. 1–29.
- KSIASZKIEWICZ, J., H. KONTECKA, AND L. NOGOWSKI. 1993. A note on blood cholesterol as an indicator of body fatness in ducks. *Journal of Animal and Feed Science* 1: 289–294.
- LAGRANGE, T. G., AND J. J. DINSMORE. 1988. Nutri-

- ent reserve dynamics of female mallards during spring migration through central Iowa. In *Waterfowl in winter*, M. W. Weller (ed.). University of Minnesota Press, Minneapolis, Minnesota, pp. 287–297.
- LAIDLEY, C. W., AND J. F. LEATHERLAND. 1988. Cohort sampling, anaesthesia and stocking-density effects on plasma cortisol, thyroid hormone, metabolite and ion levels in rainbow trout, *Salmo gairdneri* Richardson. *Journal of Fish Biology* 33: 73–88.
- MANN, F. E., AND J. S. SEDINGER. 1993. Nutrient-reserve dynamics and control of clutch size in northern pintails breeding in Alaska. *Auk* 110: 264–278.
- MARCO, I., AND S. LAVIN. 1999. Effect of the method of capture on the haematology and blood chemistry of red deer (*Cervus elaphus*). *Research in Veterinary Science* 66: 81–84.
- MARSH, R. L. 1984. Adaptations of the gray catbird *Dumetella carolinensis* to long-distance migration: Flight muscle hypertrophy associated with elevated body mass. *Physiological Zoology* 57: 105–117.
- MILLER, M. R. 1989. Estimating carcass fat and protein in northern pintails during the nonbreeding season. *Journal of Wildlife Management* 53: 123–129.
- OLSEN, J. H. 1994. Anseriformes. In *Avian medicine: Principles and application*, B. W. Ritchie, G. J. Harrison and L. R. Harrison (eds.). Wingers Publishing, Lake Worth, Florida, pp. 1237–1275.
- PUERTA, M. L., J. C. ALONSO, V. HUECAS, J. A. ALONSO, M. ABELANDA, AND R. MUNOZ-PULIDO. 1990. Hematology and blood chemistry of wintering common cranes. *Condor* 92: 210–214.
- RAVELING, D. G., AND M. E. HEITMEYER. 1989. Relationships of population size and recruitment of pintails to habitat conditions and harvest. *Journal of Wildlife Management* 53: 1088–1103.
- RINGELMAN, J. K., AND M. R. SZYMCAK. 1985. A physiological condition index for wintering mallards. *Journal of Wildlife Management* 49: 564–568.
- SHIMIZU, S., Y. TANI, H. YAMADA, M. TABATA, AND T. MURACHI. 1980. Enzymatic determination of serum-free fatty acids: A colorimetric method. *Analytical Biochemistry* 107: 193–198.
- SIEDEL, J., E. O. HAGELE, J. ZIEGENHORN, AND A. W. WAHLEFELD. 1983. Reagent for the enzymatic determination of serum total cholesterol with improved lipolytic efficiency. *Clinical Chemistry* 29: 1075–1080.
- , R. SCHMUCK, J. STAEPELS, AND M.-H. TOWN. 1993. Long-term stable, liquid, ready-to-use monoreagent for the enzymatic assay of serum or plasma triglycerides (GPO-PAP method). *AACC Meeting Abstract* 34, *Clinical Chemistry* 39: 1127.
- SMITH, L. M., D. G. SHEELEY, AND D. B. WESTER. 1992. Condition models for wintering northern pintails in the southern high plains. *Great Basin Naturalist* 52: 226–231.
- TOTZKE, U., M. FENSKE, O. HÜPPOP, H. RAABE, AND N. SCHACH. 1999. The influence of fasting on blood and plasma composition of herring gulls (*Larus argentatus*). *Physiological and Biochemical Zoology* 72: 426–437.
- WHITE, G. C., AND W. R. CLARK. 1994. Microcomputer applications in wildlife management and research. In *Research and management techniques for wildlife and habitats*, 5th Edition, T. Bookhout (ed.). The Wildlife Society, Bethesda, Maryland, pp. 75–93.
- WHITTOW, C. G. 1986. Energy metabolism. In *Avian physiology*, 4th Edition, P. D. Sturkie (ed.). Springer-Verlag, New York, New York, pp. 253–268.
- WHYTE, R. J., AND E. G. BOLEN. 1984. Variation in winter fat depots and condition indices of mallards. *Journal of Wildlife Management* 48: 1370–1373.

Received for publication 13 March 2002.