



THE RISE AND FALL OF PSOROPTIC SCABIES IN BIGHORN SHEEP IN THE SAN ANDRES MOUNTAINS, NEW MEXICO

Authors: Boyce, Walter M., and Weisenberger, Mara E.

Source: Journal of Wildlife Diseases, 41(3) : 525-531

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-41.3.525>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

THE RISE AND FALL OF PSOROPTIC SCABIES IN BIGHORN SHEEP IN THE SAN ANDRES MOUNTAINS, NEW MEXICO

Walter M. Boyce^{1,3} and Mara E. Weisenberger²

¹ Wildlife Health Center, University of California, One Shields Avenue, Davis, California 95616, USA

² United States Fish and Wildlife Service, San Andres National Wildlife Refuge, PO Box 756, Las Cruces, New Mexico 88004, USA

³ Corresponding author (email: wmboyce@ucdavis.edu)

ABSTRACT: Between 1978 and 1997, a combination of psoroptic scabies (*Psoroptes* spp.), mountain lion (*Puma concolor*) predation, and periodic drought reduced a population of native desert bighorn sheep (*Ovis canadensis*) in the San Andres Mountains (SAM), New Mexico, from >200 individuals to a single ewe. In 1999, this ewe was captured, ensured to be *Psoroptes*-free, and released back into the SAM. Eleven radiocollared rams were translocated from the Red Rock Wildlife Area (RRWA) in New Mexico into the SAM range and monitored through 2002 to determine whether *Psoroptes* spp. mites were still in the environment. None of these sentinel rams acquired scabies during this period, and no additional native sheep were found to be present in the range. In 2002, 51 desert bighorn sheep were translocated into the SAM from the Kofa National Wildlife Refuge in Arizona ($n=20$) and the RRWA in New Mexico ($n=31$). Twenty-one bighorn sheep have died in the SAM since that time, but *Psoroptes* spp. mites have not been detected on any of these animals, nor have they been found on mule deer (*Odocoileus hemionus*) sampled since 2000. We conclude that psoroptic scabies is no longer present in the San Andres bighorn sheep population and that psoroptic scabies poses a minimal to nonexistent threat to the persistence of this population at this time.

Key words: Bighorn sheep, deer, *Ovis canadensis*, *Psoroptes*, scabies.

INTRODUCTION

The San Andres Mountains (SAM) once supported the largest population of native desert bighorn sheep (*Ovis canadensis*) in New Mexico. An epizootic of psoroptic scabies (*Psoroptes* spp.) that began in 1978 reduced this population from >200 to about 25 individuals by 1989 (Lange et al., 1980; Sandoval, 1980; Hoban, 1990). From 1985 to 1995, 62% (16/26) of the radiocollared bighorn sheep that died in the SAM had clinical scabies, and by 1997, the combined effects of scabies, prolonged drought, and mountain lion (*Puma concolor*) predation had reduced the population to a single ewe (Hoban, 1990; Clark and Jessup, 1992; Logan and Sweanor, 2001).

A fundamental challenge facing managers was how to re-establish a viable bighorn sheep population in the SAM, in the face of ongoing scabies infections. In 1997, it became feasible to eliminate all mites occurring on bighorn sheep in the SAM by capturing the single remaining ewe and then reintroducing mite-free bighorn sheep. However, morphologic, antigenic,

and molecular analyses indicated that *Psoroptes* spp. mites might not be host specific (Boyce et al., 1990; Boyce and Brown, 1991; Ramey et al., 2000). Therefore, the possibility existed that *Psoroptes* spp. mites could be reintroduced into the bighorn sheep population from other hosts in the SAM environment. Faced with this uncertainty, managers decided to reintroduce bighorn sheep into the SAM only after they had strong evidence that reintroduced bighorn sheep would not become infected.

This report describes the successful effort to restore a *Psoroptes*-free bighorn sheep population in the SAM. The major components of the project were to 1) identify and treat existing bighorn sheep, 2) conduct a sentinel ram study, 3) survey different host species for the presence of *Psoroptes* spp. mites, and 4) reintroduce and monitor mite-free bighorn sheep.

MATERIALS AND METHODS

Study area and animals

The 2,059-km² (205,900 ha) primary study area, located in south-central New Mexico

(32°40'33"N, 106°32'12"W), encompasses all of the SAM range. The San Andres National Wildlife Refuge (SANWR) comprises 23,154 ha, including the southern third of the SAM, and is surrounded by White Sands Missile Range (WSMR), which is under the jurisdiction of the US Army. The SAM range is long and narrow, measuring more than 129 km (80 mi) from north to south and 9–30 km (6–19 mi) from east to west. The SAM and SANWR have been relatively undisturbed by humans because access is restricted by WSMR. Elevations within the study area range from approximately 1,280 m (4,200 ft) along the east piedmont to 2,730 m (8,957 ft) at Salinas Peak. Fourteen major east–west canyons dissect the range and drain tributary canyons from the north and south. Natural springs with perennial water sources and seeps are scattered throughout the study area.

Three native ungulate species occur on the SAM: desert mule deer (*Odocoileus hemionus*), desert bighorn sheep, and pronghorn (*Antilocapra americana*). Mule deer are the most abundant ungulate and occur throughout the study area. Pronghorn infrequently range into the foothills in the northern portion of the SAM in small bands of <20 animals; however, they are more common in the surrounding flats and likely have little or no contact with bighorn sheep. Gemsbok (*Oryx gazella*), an exotic species, were first introduced to WSMR by the New Mexico Department of Game and Fish (NMDGF) in 1969. Although gemsbok prefer desert basins, the population has increased to >3,000 animals, and they are frequently observed in most areas of the SAM. Barbary sheep (*Ammotragus lervia*), which are native to northern Africa, were released along the Canadian River by the NMDGF beginning in 1950. By 1967, the herd was estimated to be between 175 and 225 animals (Findley et al., 1975). On WSMR, Barbary sheep are occasionally observed, primarily in the precipitous mountainous regions. Carnivores that frequent the study area include mountain lion, coyote (*Canis latrans*), gray fox (*Urocyon cinereoargenteus*), and bobcat (*Lynx rufus*). The SAM range has not been grazed by livestock since the mid-1950s, although cattle (*Bos taurus*) occasionally stray into the range from adjoining areas.

Testing and treating extant bighorn sheep

Before sentinel rams could be released into the SAM, it was critical that the known remaining bighorn ewe, SAE-067, was mite-free. She was captured by helicopter net gun on 25 September 1999 for evaluation. Blood was col-

lected for serological screening for *Psoroptes* (Boyce et al., 1991), and her ears were examined for the presence of *Psoroptes* spp. mites. She was given 4.5 cc ivermectin (IVOMEC®, 1% sterile solution, Merck & Co, Inc, Rahway, New Jersey, USA), fitted with a very high frequency (VHF) radiocollar (Telonics Inc, Mesa, Arizona, USA), and quarantined in an enclosure on the SANWR pending test results.

Sentinel ram study

In November 1999, six desert bighorn rams were captured from the RRWA captive breeding facility with a helicopter net gun. Blood and ear swab samples were collected for testing to ensure that each animal was free of mites. Upon capture, the rams were fitted with VHF radiocollars (Telonics Inc.) and transported to the SAM, where they were released individually throughout the range on the basis of historical bighorn habitat locations. Four of the radiocollars had platform transmitter terminals (PTT), as well as VHF beacons. The PTT radiocollars can track animals through ARGOS satellites. The ewe and six rams were intensively monitored by telemetry, air and ground sighting, and satellite tracking over a 2-yr period. Recaptures were scheduled every 4 to 6 mo so that blood could be drawn for *Psoroptes* spp. serology, and ear swab samples were collected to check for the presence of mites. After each capture, the rams were redistributed throughout the SAM to cover as much of the mountain range as possible. Bighorn are gregarious animals and it was expected that the sentinel rams would associate with extant bighorn in the SAM, should any exist. In November 2001, the number of sentinel rams had been reduced to three as a result of three mortalities (one from undetermined natural causes and two lion kills), and to bolster the number of sentinel animals, an additional five rams were captured from the RRWA, screened, radiocollared, and released into the SAM.

Psoroptes spp. survey

From 1988 through 2003, ear swabs and crusts, whole ears, and crusty lesions were opportunistically collected from live or dead mule deer, bighorn sheep, pronghorn, gemsbok, and Barbary sheep and examined for mites. Animals were primarily sampled in the SAM, but other regions were also included to gain an understanding of the distribution of *Psoroptes* spp. mites throughout the state. Samples were examined at the University of California, Davis, for mites with the aid of a dissecting microscope; the bases of whole ears were dissected and spread flat to view all inner surfaces for the

presence of mites. Many of the mites collected during this period were used in morphometric (Boyce et al., 1990), antigenic (Boyce and Brown, 1991), and molecular (Ramey et al., 2000) analyses of host specificity.

Reintroduced bighorn sheep

In December 2002, 51 bighorn sheep were captured by helicopter net gun and translocated into the SAM from the Kofa National Wildlife Refuge (KNWR) in Arizona ($n=20$) and the RRWA captive breeding facility in New Mexico ($n=31$). At the time of capture, each animal was physically examined for mites and scabies lesions, and ear swabs were collected for microscopic evaluation. Each animal was given a SQ injection of 1% ivermectin (KNWR animals received 2–3 cc; RRWA animals received 4–6 cc), and all KNWR bighorn sheep were fitted with a Cutter Gold Insecticide cattle ear tag (10% cyfluthrin, Bayer Healthcare LLC, Shawnee Mission, Kansas, USA) in each ear. A VHF radiocollar (Telonics, Inc.) with mortality sensor was placed on each animal to facilitate monitoring after release into the SAM.

RESULTS

Testing and treating extant sheep

During two previous captures in 1993 and 1997, SAE-067 tested positive for mites, and she was treated with ivermectin each time. In 1999, although both ears were occluded with waxy crusts, there was no evidence of an active *Psoroptes* spp. infection. Serological results were also negative for circulating antibodies against *Psoroptes*. The ewe was released from the enclosure back into the SAM on 5 October 1999.

Sentinel ram study

The ewe and first set of six sentinel rams were intensively monitored from 20 November 1999 to 30 November 2001. During this time, 161 visual observations and 595 satellite telemetry locations were obtained, revealing that the sentinel rams traversed the entire SAM, migrating up and down the range numerous times. The ewe remained within the vicinity of Bennett Mountain and Black Brushy Mountain. Two rams were killed by lions in April 2000 and April 2001, and one ram died of undetermined causes in September 2001.

The original goal was to capture each bighorn every 4–6 mo for sampling, but this proved difficult. In 2000, the ewe and five of the rams were each captured at least once during two capture attempts. In 2001, the ewe and all remaining rams were recaptured between one and three times during four capture attempts. In all events, ear swabs, serology, or both were negative for *Psoroptes* spp. mites (Table 1), and no other extant bighorn were observed during the course of the study.

Psoroptes spp. survey

Most of the native bighorn sheep sampled in the SAM from November 1988 through June 1997 were infested with *Psoroptes* spp. mites (30/32 samples; Table 1). Mites were not found on any samples from the lone remaining ewe (SAE-067) after June 1997, or any of the sentinel rams in this study. Mites were never found on bighorn sheep at the RRWA breeding facility or in free-ranging populations in New Mexico, other than the SAM (Table 1).

From 1989 to October 1997, mites were found in the ears of mule deer at several locations in and around the SAM, as well as at more distant locations such as the Bosque del Apache National Wildlife Refuge, New Mexico (Table 2). No mites were found in deer ear swabs examined after 1997, but samples were limited because of a substantial statewide decline in mule deer numbers. No mites were found on ear swabs from gemsbok ($n=387$), pronghorn ($n=18$), or Barbary sheep ($n=1$; Table 2).

Reintroduced bighorn sheep

No mites were found on 50 of the 51 bighorn sheep captured and translocated to the SAM in December 2002. However, ear crust and live mites were present on one ewe (SAE-092) from KNWR. The ears of this ewe were thoroughly cleaned to remove crust and mites, and she was given 2 cc of 1% ivermectin and fitted with cyfluthrin-impregnated ear tags before

TABLE 1. Prevalence of *Psoroptes* in bighorn sheep in New Mexico.^a

| Date | Location | Sample | Prevalence (%) |
|-----------|------------------------|------------|----------------|
| Nov 1988 | San Andres Mountains | Crust | 9/10 (90) |
| Nov 1989 | San Andres Mountains | Crust | 8/9 (88.8) |
| Oct 1992 | Red Rock Wildlife Area | Swabs | 0/22 (0) |
| Oct 1993 | San Andres Mountains | Swabs | 11/11 (100) |
| Oct 1993 | Red Rock Wildlife Area | Swabs | 0/20 (0) |
| Jun 1997 | San Andres Mountains | Swabs | 2/2 (100) |
| Nov 1997 | San Francisco River | Swabs | 0/12 (0) |
| Nov 1997 | Turkey Creek | Swabs | 0/7 (0) |
| Aug 1998 | Pecos Wilderness | Swabs | 0/30 (0) |
| Jan 1999 | Manzano Mountains | Swabs | 0/9 (0) |
| Sept 1999 | San Andres Mountains | Swabs | 0/1 (0) |
| Nov 1999 | Fra Cristobal | Swabs | 0/16 (0) |
| Nov 1999 | Red Rock Wildlife Area | Swabs | 0/27 (0) |
| Nov 1999 | Peloncillos | Swabs | 0/6 (0) |
| Nov 1999 | Big Hatchets | Swabs | 0/3 (0) |
| Nov 1999 | Animas | Swabs | 0/2 (0) |
| Nov 1999 | Little Hatchets | Swabs | 0/5 (0) |
| Nov 1999 | Ladrones | Swabs | 0/1 (0) |
| Apr 2000 | San Andres Mountains | Swabs | 0/4 (0) |
| May 2000 | Ladrones | Swabs | 0/4 (0) |
| Sept 2000 | San Andres Mountains | Swabs | 0/5 (0) |
| Feb 2001 | San Andres Mountains | Swabs | 0/3 (0) |
| Mar 2001 | San Andres Mountains | Swabs | 0/2 (0) |
| Jun 2001 | San Andres Mountains | Swabs | 0/5 (0) |
| Aug 2001 | Pecos Wilderness | Swabs | 0/56 (0) |
| Nov 2001 | San Andres Mountains | Swabs | 0/4 (0) |
| Nov 2001 | Red Rock Wildlife Area | Swabs | 0/15 (0) |
| Nov 2002 | Red Rock Wildlife Area | Swabs | 0/30 (0) |
| Dec 2003 | San Andres Mountains | Ears/swabs | 0/3 (0) |

^a All San Andres Mountains samples after 1999 are part of the sentinel study.

transport and release into the SAM. This ewe has not died or been captured since translocation, so it is not known whether she still has live mites present in her ears. However, skin lesions suggestive of psoroptic scabies have not been seen during monthly field observations of the reintroduced animals. Likewise, mites have not been found on any bighorn sheep that have died ($n=21$) in the SAM since December 2002.

DISCUSSION

The epizootic of psoroptic scabies that was first detected in bighorn sheep in the SAM in 1978 persisted until 1997. Despite attempts to treat the herd with insecticidal dust bags and dips, ballistic implants, and injectable ivermectin (Sandoval, 1980; Boyce et al., 1992; Clark and Jessup,

1992), lesions and mites were consistently found in the bighorn sheep population for that 20-yr period (Lange et al., 1980; Sandoval, 1980; Hoban, 1990; Clark and Jessup, 1992; Logan and Sweanor, 2001; Table 1).

In September 1999, the last remaining native ewe was found to be free of mites, and the sentinel ram study from 1999–2001 confirmed that newly introduced bighorn sheep did not become infected with mites. Thus, the near extinction of the native bighorn sheep population finally accomplished what humans could not—the apparent elimination of *Psoroptes* spp. mites from the SAM herd.

The origin of *Psoroptes* spp. mites infecting bighorn sheep in the SAM will likely never be known. The presence of mites on mule deer in and around the

TABLE 2. Prevalence of *Psoroptes* in mule deer and other ungulates in New Mexico.

| Date | Location | Sample | Prevalence (%) |
|---------------|---------------------------------|------------|----------------|
| Mule deer | | | |
| Nov 1989 | Oscura Mountains | Ears | 2/2 (100) |
| Nov 1989 | San Andres Mountains | Swabs | 2/24 (8.3) |
| Nov 1989 | South of North Oscura Peak | Swabs | 11/34 (32.3) |
| Jan 1990 | San Andres NWR | Ears | 1/1 (100) |
| Mar 1990 | Bosque del Apache NWR | Ears | 2/2 (100) |
| Mar 1990 | Bosque del Apache NWR | Ears | 1/1 (100) |
| Apr 1990 | Bosque del Apache NWR | Ears | 2/2 (100) |
| Apr 1990 | San Andres Mountains | Swabs | 4/28 (14.3) |
| May 1990 | McGregor Range | Swabs | 0/31 (0) |
| Oct 1990 | Salinas Peak | Swabs | 11/38 (28.9) |
| Oct 1990 | San Andres Mountains | Swabs | 6/25 (24) |
| April 1991 | San Andres Mountains | Swabs | 1/37 (2.7) |
| May 1991 | McGregor Range | Swabs | 0/32 (0) |
| Oct 1991 | Oscura Mountains | Swabs | 4/17 (23.5) |
| Jan 1992 | San Augustine Pass | Ears | 1/1 (100) |
| May 1992 | San Andres Mountains | Swabs | 6/20 (30) |
| Fall 1992 | Carson National Forest | Ears | 0/1 (0) |
| Oct 1993 | San Andres Mountains | Swabs | 5/15 (33.3) |
| Jan 1996 | Bosque del Apache NWR | Ears | 1/1 (100) |
| Jan 1997 | Oscura Mountains | Swabs | 12/20 (60) |
| Sept 1997 | Oscura Mountains | Swabs | 5/16 (31.2) |
| Oct 1997 | Oscura Mountains | Swabs | 6/16 (37.5) |
| Oct 1997 | San Augustine Pass | Swabs | 3/3 (100) |
| Jun 2000 | San Augustine Pass | Swabs | 0/1 (0) |
| Oct/Nov 2002 | WSMR/Main Range Post | Ears+Swabs | 0/6 (0) |
| Dec 2003 | WSMR | Swabs | 0/21 (0) |
| Dec 2003 | Organ Mountains | Swabs | 0/6 (0) |
| Dec 2003 | WSMR/Wood Ranch | Swabs | 0/2 (0) |
| Dec 2003 | San Andres NWR | Swabs | 0/1 (0) |
| Apr 2004 | San Andres Mountains | Swabs | 0/31 (0) |
| Barbary sheep | | | |
| Apr 2000 | San Andres Mountains | Swabs | 0/1 (0) |
| Elk | | | |
| Apr 2004 | San Andres Mountains | Swabs | 0/1 (0) |
| Oryx | | | |
| Jan 1997 | Tularosa Basin | Swabs | 0/23 (0) |
| Feb 1997 | Tularosa Basin | Swabs | 0/18 (0) |
| Sept 2000 | WSMR/Stallion Range Center | Swabs | 0/56 (0) |
| Oct 2000 | WSMR/South of 70 | Swabs | 0/9 (0) |
| Nov 2000 | WSMR/Rhodes Canyon Range Center | Swabs | 0/50 (0) |
| Dec 2000 | San Andres NWR | Swabs | 0/4 (0) |
| Dec 2000 | WSMR | Swabs | 0/73 (0) |
| Jan 2001 | San Andres NWR | Swabs | 0/5 (0) |
| Jan 2001 | WSMR | Swabs | 0/89 (0) |
| Feb–Apr 2002 | San Andres NWR | Swabs | 0/34 (0) |
| May 2002 | WSMR/Stallion Range Center | Swabs | 0/26 (0) |
| Pronghorn | | | |
| Sept 2000 | WSMR/Stallion Range Center | Swabs | 0/18 (0) |

^a WSMR = White Sands Missile Range; NWR = National Wildlife Refuge.

SAM (Table 2) demonstrated that *Psoroptes* spp. mites were not limited to bighorn sheep. A key question addressed by the sentinel ram study was: do mule deer serve as a reservoir of mites that will infect bighorn sheep? The answer appears to be, "No." One caveat is that our sample size for mule deer was very low after 1997 because mule deer numbers declined markedly during the prolonged drought in the 1990s (Logan and Sweaner, 2001). Additionally, bighorn sheep and deer numbers were both very low during the sentinel study, perhaps reducing the opportunity for mites to be shared between the two host species.

The reason mite infections caused such serious clinical disease in SAM bighorn sheep is not known. Although psoroptic scabies has been linked to historical population declines of bighorn sheep in the western United States (Beuchner, 1960), most current infections seem to be subclinical (primarily confined to the ears) and not associated with any population-level effects (Sandoval, 1980; Mazet et al., 1992). One hypothesis is that mites in the SAM were more virulent than those found in other areas. However, mites from different bighorn sheep populations could not be distinguished from those found on SAM bighorn sheep by morphologic (Boyce et al., 1990), antigenic (Boyce and Brown, 1991), or molecular methods (Ramey et al., 2000). Another hypothesis is that bighorn sheep in the SAM were exceptionally susceptible to scabies. Genetic analysis of bighorn sheep from the SAM and RRWA (which were originally derived from SAM breeding stock) revealed that these populations had far less genetic variability than other bighorn sheep populations (Boyce et al., 1997; Gutierrez-Espeleta et al., 2001). Although no evidence links low genetic variability with enhanced susceptibility to psoroptic scabies, decreased heterozygosity (as was found in the SAM and RRWA bighorn sheep) has been associated with decreased survival in Soay sheep (Coltman et al., 1999).

The results of the sentinel study (no new mite infections), coupled with failure to find mites on mule deer after 1997, provided strong support for the decision to reintroduce bighorn sheep to the SAM in 2002. Animals were obtained from two locations (KNWR and RRWA) because of genetic considerations and the need to introduce a substantial number of animals. Bighorn sheep from the KNWR in Arizona were identified as suitable genetic reintroduction stock because they were highly variable compared with RRWA and SAM animals (Gutierrez-Espeleta et al., 2001). Despite the occasional sighting of *Psoroptes* spp. mites in the ears of KNWR bighorn (Sandoval, 1980), this population had proven to be a productive source of animals for several other bighorn sheep transplants in Arizona and other western states (Cunningham et al., 1993). Because only a limited number of animals were available from KNWR, 31 animals were also translocated from the RRWA. This ensured that at least some descendants of native SAM bighorn sheep were reintroduced, and it increased the starting size of the population.

During the capture and translocation effort in December 2002, one ewe from KNWR was found to have *Psoroptes* spp. mites and crusty material in both ears. After considerable discussion, the decision was made to transport her to the SAM rather than release her back into KNWR. She was treated for mites before transport, but it is possible that live mites were translocated with her into the SAM. It is not known whether this ewe is still infected with mites because she has not died or been re-examined since her release in 2002. However, since the translocation effort, scabies lesions have not been seen on live bighorn sheep during monthly field observations, and mites have not been found on examination of dead sheep.

The decision to relocate the *Psoroptes*-infected ewe from KNWR was controversial. If psoroptic scabies reoccurs in the SAM bighorn sheep population, it will not be possible to determine whether mites

were reintroduced with the KNWR ewe or originated from some other infected host already present in the SAM. Because KNWR bighorn sheep have evidently coexisted with *Psoroptes* spp. mites for some period of time, it is also possible that any mites introduced by the KNWR ewe are less pathogenic than those responsible for the epizootic that began in the 1970s.

We conclude that the epizootic of psoroptic scabies in the SAM bighorn sheep population has ended and that psoroptic scabies poses a minimal to nonexistent threat to the persistence of this population at this time. Current management efforts are focused on predator control (lion removal), habitat enhancement (prescribed fire), and herd health monitoring. It remains to be seen whether or not sheep numbers will increase to those seen in the 1970s (>200) before the onset of the scabies epizootic.

ACKNOWLEDGMENTS

This project was supported by the New Mexico Department of Game and Fish (NMDGF), White Sands Missile Range, National Fish and Wildlife Foundation, and the US Fish and Wildlife Service. We especially appreciate the support and cooperation of Eric Rominger, Darrel Weybright, Brock Hoenes, and Amy Fisher from NMDGF; Grace Lee from the University of California, Davis; and personnel from the Arizona Department of Game and Fish, Arizona Desert Bighorn Sheep Society, and New Mexico Chapter of Foundation for North American Wild Sheep.

LITERATURE CITED

- BEUCHNER, H. K. 1960. The bighorn sheep in the United States, its past, present, and future. *Wildlife Monographs* 4, pp. 63–64.
- BOYCE, W. M., AND R. N. BROWN. 1991. Antigenic characterization of *Psoroptes* spp. (Acari: Psoroptidae) mites from different hosts. *Journal of Parasitology* 77: 675–679.
- , L. ELLIOTT, R. CLARK, AND D. JESSUP. 1990. Morphometric analysis of *Psoroptes* spp. mites from bighorn sheep, mule deer, cattle, and rabbits. *Journal of Parasitology* 76: 823–828.
- , J. A. K. MAZET, J. MELLIES, I. GARDNER, R. K. CLARK, AND D. A. JESSUP. 1991. Kinetic ELISA for detection of antibodies to *Psoroptes* sp. (Acari: Psoroptidae) in bighorn sheep (*Ovis canadensis*). *Journal of Parasitology* 77: 692–696.
- , J. A. MILLER, D. A. JESSUP, AND R. K. CLARK. 1992. Use of ivermectin implants for the treatment of psoroptic scabies in free-ranging bighorn sheep. *Journal of Zoo and Wildlife Medicine* 23: 211–213.
- , P. W. HEDRICK, N. E. MUGGLI-COCKETT, S. KALINOWSKI, M. C. T. PENEDO, AND R. R. RAMEY II. 1997. Genetic variation of major histocompatibility complex and microsatellite loci: A comparison in bighorn sheep. *Genetics* 145: 421–433.
- CLARK, R. K., AND D. A. JESSUP. 1992. The health of mountain sheep in the San Andres Mountains, New Mexico. *Desert Bighorn Council Transactions* 36: 30–55.
- COLTMAN, D. W., J. G. PILKINGTON, J. A. SMITH, AND J. M. PEMBERTON. 1999. Parasite-mediated selection against inbred Soay sheep in a free-living, island population. *Evolution* 53: 1259–1267.
- CUNNINGHAM, S., N. DODD, AND R. OLDING. 1993. Arizona's bighorn sheep reintroduction program. In *The desert bighorn sheep in Arizona*, R. M. Lee (ed.). Arizona Game and Fish Department, Phoenix, Arizona. pp. 203–239.
- FINDLEY, J. S., A. H. HARRIS, D. E. WILSON, AND C. JONES. 1975. *Mammals of New Mexico*. University of New Mexico Press, Albuquerque, New Mexico. 360 pp.
- GUTIERREZ-ESPELETA, G. A., P. W. HEDRICK, S. T. KALINOWSKI, D. GARRIGAN, AND W. M. BOYCE. 2001. Is the decline of desert bighorn sheep from infectious disease the result of low MHC variation? *Heredity* 86: 439–450.
- HOBAN, P. 1990. A review of desert bighorn sheep in the San Andres Mountains, New Mexico. *Desert Bighorn Council Transactions* 34: 14–22.
- LANGE, R. E., A. V. SANDOVAL AND W. P. MELENEY. 1980. Psoroptic scabies in bighorn sheep (*Ovis canadensis mexicana*) in New Mexico. *Journal of Wildlife Diseases* 16: 77–82.
- LOGAN, K. A., AND L. L. SWEANOR. 2001. *Desert puma: Evolutionary ecology and conservation of an enduring carnivore*. Island Press, Covelo, California, 463 pp.
- MAZET, J. A. K., W. M. BOYCE, J. MELLIES, I. A. GARDNER, R. K. CLARK, AND D. A. JESSUP. 1992. Exposure to *Psoroptes* sp. mites is common among bighorn sheep (*Ovis canadensis*) populations in California. *Journal of Wildlife Diseases* 28: 542–547.
- RAMEY, R. R., S. T. KELLEY, W. M. BOYCE, AND B. D. FARRELL. 2000. Phylogeny and host specificity of psoroptic mange mites (Acarina: Psoroptidae) as indicated by ITS sequence data. *Journal of Medical Entomology* 37: 791–796.
- SANDOVAL, A. V. 1980. Management of a psoroptic scabies epizootic in bighorn sheep (*Ovis canadensis mexicana*) in New Mexico. *Desert Bighorn Council Transactions* 24: 21–28.

Received for publication 27 September 2004.