

Isolation of Salmonella Enteritidis Phage Type 21b from a Eurasian Eagle-Owl (*Bubo bubo*)

Authors: Kocabiyik, A. Levent, Cangul, I. Taci, Alasonyalilar, Aylin, Dedicova, Daniela, and Karpiskova, Renata

Source: Journal of Wildlife Diseases, 42(3) : 696-698

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-42.3.696>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Isolation of *Salmonella* Enteritidis Phage Type 21b from a Eurasian Eagle-Owl (*Bubo bubo*)

A. Levent Kocabiyik,^{1,5} I. Taci Cangul,² Aylin Alasonyalilar,² Daniela Dedicova,³ and Renata Karpiskova⁴

¹ Microbiology Department, Faculty of Veterinary Medicine, Uludag University, 16059, Gorukle Campus, Bursa, Turkey; ² Pathology Department, Faculty of Veterinary Medicine, Uludag University, 16059, Gorukle Campus, Bursa, Turkey; ³ National Reference Laboratory for Salmonella, National Institute of Public Health, Srobarova 48, 10042 Prague 10, Czech Republic; ⁴ Centre for the Hygiene of Food Chains, National Institute of Public Health, Palackeho 3a, 61242 Brno, Czech Republic; ⁵ Corresponding author: (email: klevent@uludag.edu.tr)

ABSTRACT: A case of fatal salmonellosis in a Eurasian eagle-owl (*Bubo bubo*) from Bursa Province (northwestern Turkey) is described. The organs of the bird were examined histopathologically and microbiologically. Macroscopic and microscopic findings were consistent with a *Salmonella* infection. *Salmonella enterica* subspecies *enterica* serovar Enteritidis (*S. Enteritidis*) phage type (PT) 21b was isolated from the liver and spleen in pure culture and from the intestine. The isolate was susceptible to amoxicillin/clavulanic acid, ampicillin, chloramphenicol, gentamicin, kanamycin, tetracycline, and trimethoprim/sulphamethoxazole. This is the first report of an isolation of salmonellae from a wild bird species from Turkey and the first time *S. Enteritidis* PT21b has been reported from Turkey.

Key words: Antimicrobial susceptibility, Eurasian eagle owl, *Salmonella* Enteritidis phage type 21b, wild bird.

A male Eurasian eagle-owl (*Bubo bubo*) found dead in a private garden in Bursa Province in June 2005 was submitted to the Faculty of Veterinary Medicine, Uludag University, Bursa, Turkey, for diagnostic evaluation. At postmortem examination, small whitish nodules of 1–3 mm in diameter were observed on the liver surface. The proventriculus mucosa and the passage into the ventriculus were covered with hemorrhagic exudate.

Tissue samples from various organs were routinely processed, embedded in paraffin, sectioned at 5 µm, and stained with hematoxylin and eosin. Smears of tissues were stained using Gram and modified Ziehl-Neelsen methods. Tissue samples also were inoculated onto xylose-lysine-desoxycholate agar (Oxoid Ltd., Basingstoke, UK) as well as other culture media routinely used for pathogenic bac-

teria and fungi. Presumptive *Salmonella* colonies were chosen and subjected to biochemical tests. Species and subspecies differentiation of the isolate were performed according to the method described by Popoff (2001). Serotyping and phage typing were performed according to the Kauffman-White scheme (Rowe and Hall, 1989) and Ward et al. (1987), respectively. The susceptibility of the *Salmonella* isolate to various antibiotics was examined by the Kirby-Bauer disc diffusion method (Bauer et al., 1966) on Mueller-Hinton agar (Oxoid).

Microscopic examination of tissues revealed areas of coagulation necrosis in the hepatic parenchyma, together with mononuclear cell infiltrations in the subcapsular and periportal areas. Walls of branches of hepatic artery and portal vein showed signs of degeneration, and the lumina contained thrombi. Coccidian merozoites and oocysts were observed within the epithelial lining of the small intestine together with moderate inflammation. Focal mononuclear cell infiltrations, to a lesser extent, also were observed in the lamina propria and submucosa of the proventriculus, small intestine, and in the heart (together with focal myocyte degeneration). Thrombi were present in the middle-sized vessels in the lung.

Microscopically, gram-negative bacilli were observed in large numbers on liver smears and in lesser numbers on spleen smears. *Salmonella* was isolated from the liver and spleen in pure culture, and also from the intestine. The isolate was identified as *Salmonella enterica* subspecies *enterica* according to biochemical reaction

results. In serotyping, the strain had the following antigenic formula: 9, 12: g, m: —, corresponding to serovar Enteritidis. The phage type of the serovar was identified to be 21b (PT21b). The isolated agent was susceptible to amoxicillin/clavulanic acid, ampicillin, chloramphenicol, gentamicin, kanamycin, tetracycline, and trimethoprim/sulphamethoxazole, but resistant to cephalothin and nalidixic acid.

Serovar Typhimurium is a well-known cause of salmonellosis in many wild bird species (Pennycott et al., 1998; Refsum et al., 2002, 2003). In this report we describe a case of fatal salmonellosis in an owl caused by *S. Enteritidis* PT21b. Isolation of the bacteria from the liver, spleen, and intestine indicates the presence of a septicemic infection in the bird. *Coccidia* observed in the intestine could have enhanced the severity of *Salmonella* infection in this bird. Death associated with salmonellosis caused by nontyphoidal serovars of *S. enterica* usually occurs in birds coinfecting with other agents, especially with *coccidia*, presumably due to the increased ability of *Salmonella* to colonize the intestinal tract (Qin et al., 1995).

In the last few decades, *S. Enteritidis* has emerged as a major cause of foodborne illness worldwide (Agron et al., 2001). The serovar has been most commonly isolated from humans (Erdem et al., 2005) and chickens (Carli et al., 2001) in Turkey as well. However, there are no data regarding the existence of the agent in wild birds of the Turkish fauna, which may also act as an important reservoir for transmission of salmonellae to humans (Kapperud et al., 1998; Hudson et al., 2000). Kapperud et al. (1998) reported that sporadic cases of domestically acquired human salmonellosis and cases of fatal salmonellosis among small passerines in Norway were most often observed at the same time of year, thus indicating an epidemiologic link. In addition, a molecular epidemiologic study has shown that 32% of the isolates recovered from human patients belonged to clones detected in

the small passerine fauna in Norway (Heir et al., 2002).

This is the first report on the isolation of salmonellae from a wild bird species of Turkish fauna, and the first time this phage type was documented in Turkey. The study indicates that there is strong need to clarify the distribution of salmonellae in wild bird populations and the potential epidemiological relationship between salmonellae in wild birds and humans in Turkey.

LITERATURE CITED

- AGRON, P. G., R. L. WALKER, H. KINDE, S. J. SAWYER, D. C. HAYES, J. WOLLARD, AND G. L. ANDERSEN. 2001. Identification by subtractive hybridization of sequences specific for *Salmonella enterica* serovar Enteritidis. *Applied and Environmental Microbiology* 67: 4984–4991.
- BAUER, A. W., W. M. KIRBY, J. C. SHERRIS, AND M. TURCK. 1966. Antibiotic susceptibility testing by a standardized single disc method. *American Journal of Clinical Pathology* 45: 493–496.
- CARLI, K. T., A. EYGOR, AND V. CANER. 2001. Prevalence of *Salmonella* serovars in chickens in Turkey. *Journal of Food Protection* 64: 1832–1835.
- ERDEM, B., S. ERCIS, G. HASCELİK, D. GUR, S. GEDİKOĞLU, A. D. AYSEV, B. SUMERKAN, M. TATMAN-OTKUN, AND I. TUNCER. 2005. Antimicrobial resistance patterns and serotype distribution among *Salmonella enterica* strains in Turkey, 2000–2002. *European Journal of Clinical Microbiology and Infectious Diseases* 24: 220–225.
- HEIR, E., B. A. LINDSTEDT, I. NYGARD, T. VARDUND, V. HASSELTVEDT, AND G. KAPPERUD. 2002. Molecular epidemiology of *Salmonella* Typhimurium isolates from human sporadic and outbreak cases. *Epidemiology and Infection* 128: 373–382.
- HUDSON, C. R., C. QUIST, M. D. LEE, K. KEYES, S. V. DODSON, C. MORALES, S. SANCHEZ, D. G. WHITE, AND J. J. MAURER. 2000. Genetic relatedness of *Salmonella* isolates from nondomestic birds in Southeastern United States. *Journal of Clinical Microbiology* 38: 1860–1865.
- KAPPERUD, G., H. STENWIG, AND J. LASSEN. 1998. Epidemiology of *Salmonella typhimurium* O:4–12 infection in Norway: Evidence of transmission from an avian wildlife reservoir. *American Journal of Epidemiology* 147: 774–782.
- PENNYCOTT, T. W., H. M. ROSS, I. M. McLAREN, A. PARK, G. F. HOPKINS, AND G. FOSTER. 1998. Causes of death of wild birds of the family Fringillidae in Britain. *Veterinary Record* 143: 155–158.

- POPOFF, M. Y. 2001. Antigenic formulas of the *Salmonella* serovars. 8th Edition, WHO Collaborating Centre for Reference and Research on Salmonella, Institut Pasteur, Paris, France.
- QIN, Z. R., T. FUKATA, E. BABA, AND A. ARAKAWA. 1995. Effect of *Eimeria tenella* infection on *Salmonella enteritidis* infection in chickens. Poultry Science 74: 1–7.
- REFSUM, T., K. HANDELAND, D. L. BAGGESEN, G. HOLSTAD, AND G. KAPPERUD. 2002. Salmonellae in avian wildlife in Norway from 1969 to 2000. Applied and Environmental Microbiology 68: 5595–5599.
- , T. VIKOREN, K. HANDELAND, G. KAPPERUD, AND G. HOLSTAD. 2003. Epidemiologic and pathologic aspects of *Salmonella* Typhimurium infection in passerine birds in Norway. Journal of Wildlife Diseases 39: 64–72.
- ROWE, B., AND M. L. M. HALL. 1989. Kauffman-White Scheme., Public Health Laboratory Service, London, UK, 77 pp.
- WARD, L. R., J. D. DE SA, AND B. ROWE. 1987. A phage-typing scheme for *Salmonella enteritidis*. Epidemiology and Infection 99: 291–294.

Received for publication 30 September 2005.