

Disseminated Bovine Tuberculosis in a Wild Red Fox (*Vulpes vulpes*) in Southern Spain

Authors: Millán, Javier, Jiménez, María Ángeles, Viota, María, Candela, Mónica G., Peña, Laura, et al.

Source: Journal of Wildlife Diseases, 44(3) : 701-706

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-44.3.701>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Disseminated Bovine Tuberculosis in a Wild Red Fox (*Vulpes vulpes*) in Southern Spain

Javier Millán,^{1,4,5} María Ángeles Jiménez,² María Viota,¹ Mónica G. Candela,³ Laura Peña,² and Luis León-Vizcaino³ ¹Department of Conservation Biology, Estación Biológica de Doñana (CSIC), Pabellón del Perú, Avda. María Luisa s/n, 41013-Sevilla, Spain; ²Department of Pathology, Veterinary School, Complutense University, Madrid, Spain; ³Infectious Diseases Area, Department of Animal Health, Faculty of Veterinary Medicine, University of Murcia, Murcia, Spain; ⁴Present address: Ul. Piłsudskiego 13/11, 69-100, Ślubice, Poland; ⁵Corresponding author (email: syngamustrachea@hotmail.com)

ABSTRACT: A wild-caught, adult red fox (*Vulpes vulpes*) from Doñana National Park (southern Spain), in very poor condition, died during anesthesia. At necropsy, the submandibular, retropharyngeal, mediastinal, axillary, mesenteric, and popliteal lymph nodes were enlarged, and the right submandibular and mesenteric lymph nodes had hard, white-colored concretions (calcifications). Multiple white pinpoint foci were observed in the lungs, and abscesses were present in the left dorsal lung lobes. No lesions were seen in tonsils, liver, or spleen. On histopathology multiple tuberculous granulomas, with Ziehl-Neelsen-positive bacilli, were observed in the lung, and multifocal necrotic areas with calcification were present in the submandibular and mesenteric lymph nodes. Nucleic acid amplification from lymph node samples demonstrated the presence of mycobacteria belonging to the *Mycobacterium tuberculosis* complex. One strain was isolated by culture and identified as *Mycobacterium bovis*. The suspected route of infection was oral, probably after repeated scavenging of infected wild ungulate carcasses. This is the first report of generalized bovine tuberculosis (bTB) in a wild canid. This finding raises concerns about bTB as a disease risk for protected species, livestock, and humans in Mediterranean Spain.

Key words: Andalusia, Canidae, carnivore, *Mycobacterium bovis*, scavenger, tuberculosis.

Mycobacterium bovis, the etiological agent of bovine tuberculosis (bTB), has one of the broadest host ranges of all pathogens (de Lisle et al., 2001), including carnivorous mammals. However, canids appear less susceptible to the disease, especially when compared with felids or mustelids. For example, in a 30-yr-long survey in New Zealand, two isolates of *M. bovis* were obtained in dogs in contrast with 76 isolates in cats (de Lisle et al., 1993). Gay et al. (2000) stated that dogs

are naturally resistant to *M. bovis*. This may be speculative, but there are few reports of *M. bovis* infection among wild canids (Table 1), even though the family Canidae include scavenging species inhabiting areas where *M. bovis* is endemic (e.g. the red fox [*Vulpes vulpes*] in Europe or North America, or the wild dog [*Lycan ppyctus*] in Africa). Bruning-Fann et al. (2001) found bTB-suggestive gross or histologic lesions in five coyotes (*Canis latrans*) and one red fox; acid-fast bacilli were observed in only two of these coyotes. Delahay et al. (2006) reported gross lesions in one out of 24 bTB culture-positive red foxes. Carbyn (1982) reported enlargement of lymph nodes in two wolves (*Canis lupus*) infected with *M. bovis* and bTB lesions have been reported in captive fennec foxes (*Fennecus zerda*; Himes et al., 1980). In 2005, *M. bovis* was isolated for the first time in culture from a red fox in Spain. This animal came from the Doñana National Park (DNP) and did not have macroscopic or histologic lesions (Martín-Atance et al., 2005).

In the present case, we describe a disseminated disease caused by *M. bovis* in a wild red fox from the Reserva Biológica, in the DNP (southern Spain: 37°0'N, 6°30'W). The fox was an old male captured with a box trap in February 2005, as part of an ecologic research program. The animal was emaciated with poor pelage, skin ulcers, dyspnea, and pale mucosa. The fox was anesthetized with a combination of ketamine (Imalgène®, Merial, France) plus medetomidine (Domtor®, Salud Animal-Pfizer, Spain), and died about 20 min after induction.

TABLE 1. Summary of reports of *Mycobacterium bovis* infecting wild canids. This summary does not include captive wildlife.

Species	Origin	Prevalence	Lesions ^a	Suspected route of infection ^b	Reference
Coyote	USA	1/23	No	Ingestion of elk (<i>Cervus elaphus</i>)	Rhyan et al. 1995
<i>Canis latrans</i>	USA	6/110	M (1), H (3), both (1)	Ingestion of white-tailed deer (<i>Odocoileus virginianus</i>)	Bruning-Fann et al. 2001
Wolf	Canada	2/21	M (2)	n.s.	Carbyn 1982
<i>Canis lupus</i>					
Gray fox	USA	1/?	n.s.	n.s.	Patrick 2003
<i>Urocyon cinereo-argenteus</i>					
Red fox	USA	1/6	No	Ingestion of white-tailed deer	Bruning-Fann et al. 2001
<i>Vulpes vulpes</i>	Spain	1/7	No	n.s.	Martín-Atance et al. 2005
	UK	24/756	M (1)	Ingestion of infectious carcasses of cattle or cervids	Delahay et al. 2006
	Spain	2/24	H (1), both (1)	Ingestion of infectious carcasses of wild ungulates	Present study

^a M=macroscopic, H=histologic, (n)=number of cases; unspecific lesions such as “enlargement of lymph nodes” are not included.

^b n.s.=not specified.

Whole blood was obtained from jugular veins just before death, collected in serum separator tubes, and allowed to clot. The sample was centrifuged and serum was removed and frozen at -20°C until analyzed. The carcass was necropsied and samples from the major viscera were taken. Samples for histology were routinely fixed in buffered formalin and processed in 4-mm paraffin-embedded cassettes; 4- μm cuts were obtained and stained with hematoxylin-eosin and Ziehl-Neelsen (ZN). Polymerase chain reaction (PCR) was used for detection of *Mycobacterium tuberculosis* complex (MTC). Nucleic acid corresponding to MTC in microbiologic culture and in mandibular or retropharyngeal lymph nodes was performed with the use of primers TB1-F (GAA CAA TCC GCA GTT GAC AA) and TB1-R (AGC ACG CTG TCA ATC ATG TA), as described previously (Bödinghaus et al., 1990; Liébana et al., 1996), which amplify a 372-base-pair (bp) fragment of the gene that coded the MPB20 antigen specific for MTC (Cousins et al., 1991). Culture was undertaken with the use of macerates of lymph node

tissue suspended in 0.2% buffered bovine albumin, and decontaminated with a benzalkonium substrate (Kubica’s method modified by Krasnow) (Casal, 1983). Centrifuged deposits were resuspended in 10% buffered bovine serum and cultured on Middlebrook’s 7H11 oleic acid albumin agar medium (Gallagher and Horwill, 1977) and Lowenstein-Jensen with pyruvate and without glycerol (Corner and Nicolacopoulos, 1988). The microbiologic identification of mycobacteria was based on tests selected by Wayne and Kubica (1986) in accordance with the methods reported previously (León-Vizcaíno et al., 1990). Antibodies against *M. bovis* were detected in serum with the use of the indirect competition enzyme-linked immunosorbent assay (ELISA) technique. This was developed according to the method described by Harboe et al. (1990) with modifications (Acosta et al., 2000). Antigen consisted of MPB70 protein (Harboe and Nagai, 1984) and an anti-*M. bovis* goat serum was used to measure antigen captured by test and control sera. Positive- and negative-control serum samples were from red fox;



FIGURE 1. Bovine tuberculosis lesions in red fox (*Vulpes vulpes*). Calcification (arrows) of mesenteric lymph node (upper left), and in a section of the submandibular lymph node (upper right); and an abscess (arrowhead) in the left dorsal lobe of the lung (below). Bars=1 cm.

positive serum was obtained from a fox that was experimentally vaccinated with *M. bovis* BCG (see Martín-Atance et al., 2006 for further details).

External examination indicated that this animal was in very poor body condition. Deep ulcers were noted bilaterally in the skin over both ischia. The right ear was torn and the tip was missing. The muzzle was scarred and many teeth were worn down. Submandibular, retropharyngeal, mediastinal, axillary, mesenteric, and popliteal lymph nodes were enlarged. Hard, white concretions (calcifications) were detected in the right submandibular and mesenteric lymph nodes when cut (Fig. 1). No lesions were seen in the tonsils. The heart was enlarged and pale. In the lung, abscesses in the dorsal aspect of the left caudal lobe were present (Fig. 1). Multiple white pinpoint foci were

seen in both lungs. Both types of lesions were attributable to *M. bovis* infection. Bronchopulmonary nematodes were observed in the right lung. The stomach was empty of feedstuff. Blood vessels appeared engorged in the intestines. The gall bladder was full and the liver was grossly normal. The spleen was pale. Perirenal adipose tissue was absent and the adrenal glands were enlarged.

In histopathologic examination, the lung contained multifocal nodular accumulations of lymphocytes, macrophages, and some plasma cells, occasionally surrounding necrotic cellular debris or sections of metazoan parasites (granulomas). Acid-fast bacilli were detected in the cytoplasm of some macrophages (two–four bacilli per granuloma). The submandibular lymph node contained scattered giant multinucleated cells and multifocal dark basophil-

ic concretions (calcifications). The mesenteric lymph nodes contained similar calcifications and multifocal necrotic areas surrounded by lymphocytes, macrophages, and plasma cells. The liver had generalized necrosis. Antibodies against *M. bovis* were detected in sera. Nucleic acid amplification from lymph node samples demonstrated the presence of mycobacteria belonging to the *M. tuberculosis* complex. One strain was isolated by means of culture and identified by biochemical test as *M. bovis*. A concomitant infection with Canine Adenovirus-1 was also detected by means of immunofluorescence [210-34-CAV FITC Conjugate (Anti-Canine Adenovirus polyclonal antiserum conjugated to fluorescein isothiocyanate), VMRD Inc., Pullman, Washington, USA], and competitive ELISA (Canine Hepatitis IgG ELISA, Eurovet Veterinaria Daganzo, Madrid, Spain).

From our review of the literature, this is the first report of disseminated bTB in a wild canid. In the domesticated dog, it appears that natural *M. bovis* infection is one of wide anatomic lesion distribution (Francis, 1958). The generalized distribution of lesions of the disease in the present case may have been due to the advanced age and general debility of the fox, which may have increased the susceptibility of the animal to generalized infection. Tuberculosis may have a clinical course of many months (Kaneene and Thoen, 2004). In agreement with our findings, the six infected coyotes studied by Bruning-Fann et al. (2001) were also adults. Conversely, the cases of bTB in wolves reported by Carbyn (1982) were two littermate pups of about 6 mo of age, both found emaciated because of the *M. bovis* infection. The presence of concurrent infections, such as infectious canine hepatitis (which may have been the cause of the hepatic necrosis), may also have weakened the immune system of the fox and as a consequence predisposed it to generalized bTB.

The distribution of the lesions suggested that the primary route of infection was

the digestive tract, probably after repeated scavenging of infected wild ungulate carcasses. This agrees with other reports of bTB in canids that are often associated with high prevalences of bTB among cattle or wild cervid populations (Table 1). Red deer (*Cervus elaphus*), fallow deer (*Dama dama*), and wild boar (*Sus scrofa*) in southern, Mediterranean Spain have been recorded with a high prevalence of bTB (up to 100% in some wild boar populations; Vicente et al., 2006). These species are probably bTB reservoirs in this region of Spain (Vicente et al., 2006). Ungulates from DNP, where the fox was caught, are not an exception (Aranaz et al., 2004). As previously indicated, the infected fox reported by Martín-Atance et al. (2005) also belonged to DNP. In addition, antibodies against *M. bovis* were found in 23% of badgers (*Meles meles*) and 3% of endangered Iberian lynxes (*Lynx pardinus*) in DNP (Martín-Atance et al., 2006). Moreover, in this location three lynxes were found to be *M. bovis* positive, and the isolates showed an identical spoligotyping pattern to the predominant pattern found in domesticated and wild ungulate species in Southern Spain (Aranaz et al., 2004).

Feeding ecology may also explain the differences in the etiologic agent of TB between wild canids and the domestic dog. Although dogs are equally sensitive to *M. bovis* and *M. tuberculosis* experimentally (Francis, 1958), approximately 75% of cases of tuberculosis reported in dogs are caused by *M. tuberculosis* (Forster et al., 1986). Therefore, whereas dogs usually become infected with *M. tuberculosis* by indirect association with humans (Hackendahl et al., 2004), wild canids may become infected by *M. bovis* by the ingestion of infected carcasses.

Carnivores may serve as bTB reservoirs in other countries, for example, badgers in the UK (Delahay et al., 2006), or feral ferrets (*Mustela furo*) in some situations in Australia (Corner, 2006). However, the fox is usually a spillover host, at least in the

UK (Delahay et al., 2006), and probably in the Iberian Peninsula. So far, another 24 foxes from DNP have been necropsied and only in one other case was a Z-N-positive individual detected (unpubl. data).

The finding of a positive animal of a species with little reported susceptibility to *M. bovis*, such as the red fox, with generalized bTB lesions, suggests that opportunities for infection with *M. bovis* are present in the natural environment in the southern Iberian Peninsula, and that bTB is a threat for the conservation of wild species including the extremely endangered Iberian lynx, that inhabits DNP. In addition, biologists and other professionals working with wild carnivores must take into account the risks of acquiring bTB infection from infected samples and tissues, or from bite wounds.

This is a contribution to the “Programa de Actuaciones para la Conservación del Lince Ibérico en Andalucía II” (Consejería de Medio Ambiente, Junta de Andalucía). We wish to thank M. Delibes, F. Palomares, and S. Conradi. R. Ballesteros kindly helped in the composition of Figure 1. P. Duff was very helpful reviewing the English.

LITERATURE CITED

- ACOSTA, B., F. REAL, L. LEÓN, S. DÉNIZ, O. FERRER, I. ROSARIO, AND A. RAMÍREZ. 2000. ELISA for anti-MPB70: An option for the diagnosis of goat tuberculosis caused by *Mycobacterium bovis*. *Australian Veterinary Journal* 78: 423–424.
- ARANAZ, A., L. DE JUAN, N. MONTERO, C. SÁNCHEZ, M. GÁLKA, C. DELSO, J. ÁLVAREZ, B. ROMERO, J. BEZOS, A. I. VELA, V. BRIONES, A. MATEOS, AND L. DOMÍNGUEZ. 2004. Bovine tuberculosis (*Mycobacterium bovis*) in wildlife in Spain. *Journal of Clinical Microbiology* 42: 2602–2608.
- BÖDDINGHAUSS, B., T. ROGALL, T. FLOHR, H. BLÜCKER, AND E. C. BÜTTGER. 1990. Detection and identification of mycobacteria by amplification of rRNA. *Journal of Clinical Microbiology* 28: 1751–1759.
- BRUNING-FANN, C. S., S. M. SCHMITT, S. D. FITZGERALD, J. S. FIERKE, P. D. FRIEDRICH, J. B. KANEENE, K. A. CLARKE, K. L. BUTLER, J. B. PAYEUR, D. L. WHIPPLE, T. M. COOLEY, J. M. MILLER, AND D. P. MUZO. 2001. Bovine tuberculosis in free-ranging carnivores from Michigan. *Journal of Wildlife Diseases* 37: 58–64.
- CARBYN, L. N. 1982. Incidence of disease and its potential role in the population of wolves in Riding Mountain National Park, Manitoba. In *Wolves of the world. Perspectives of behavior, ecology and conservation*, F. H. Harrington and P. C. Paquet (eds.). Noyes Publications, Park Ridge, New Jersey, pp. 106–116.
- CASAL, M. 1983. *Bacteriología de la tuberculosis y micobacterias*. Editorial AC, Madrid, 125 pp.
- CORNER, L. A. L. 2006. The role of wild animal populations in the epidemiology of tuberculosis in domestic animals: How to assess the risk. *Veterinary Microbiology* 112: 303–312.
- , AND C. NICOLACOPOULOS. 1988. Comparison of media used for the primary isolation of *Mycobacterium bovis* by veterinary and medical diagnostic laboratories. *Australian Veterinary Journal* 65: 202–205.
- COUSINS, D. V., S. D. WILTON, B. R. FRANCIS, AND B. L. GOW. 1991. Use of polymerase chain reaction for rapid diagnosis of tuberculosis. *Journal of Clinical Microbiology* 30: 255–258.
- DELAHAY, R. J., G. C. SMITH, A. M. BARLOW, N. WALKER, A. HARRIS, R. S. CLIFTON-HADLEY, AND C. L. CHEESEMAN. 2006. Bovine tuberculosis infection in wild mammals in the south-west region of England: A survey of prevalence and a semi-quantitative assessment of the relative risks to cattle. *Veterinary Journal* 164: 90–105.
- DE LISLE, G. W., K. CREWS, J. DE ZWART, R. JACKSON, G. J. E. KNOWLES, K. D. PATTERSON, R. W. MACKENZIE, K. A. WALDRUP, AND R. WALKER. 1993. *Mycobacterium bovis* infections in wild ferrets. *New Zealand Veterinary Journal* 41: 148–149.
- , C. G. MACKENZIE, AND R. G. BENGIS. 2001. *Mycobacterium bovis* in free-living and captive wildlife, including farmed deer. *Revue Scientifique et Technique—Office International des Epizooties* 20: 86–111.
- FORSTER, E. S., T. D. SCAVELLI, P. G. GREENLEE, AND S. R. GILBERTSON. 1986. Cutaneous lesion caused by *Mycobacterium tuberculosis* in a dog. *Journal of the American Veterinary Medical Association* 188: 1188–1190.
- FRANCIS, J. 1958. *Tuberculosis in animals and man: A study in comparative pathology*. Cassell and Company Limited, London, UK, 357 pp.
- GALLAGER, J., AND D. M. HORWILL. 1977. A selective oleic acid albumin agar medium for the cultivation of *Mycobacterium bovis*. *Journal of Hygiene (Cambridge)* 79: 155–160.
- GAY, G., H. M. BURBIDGE, P. BENNETT, S. G. FENWICK, C. DUPONT, A. MURRAY, AND M. R. ALLEY. 2000. Pulmonary *Mycobacterium bovis* infection in a dog. *New Zealand Veterinary Journal* 48: 78–81.
- HACKENDAHL, N. C., D. I. MAWBY, D. A. BEMIS, AND

- S. L. BEAZLEY. 2004. Putative transmission of *Mycobacterium tuberculosis* infection from a human to a dog. *Journal of the American Veterinary Medical Association* 225: 1573–1577.
- HARBOE, M., AND S. NAGAI. 1984. MPB70, an unique antigen of *Mycobacterium bovis* BCG. *American Journal of Respiratory Diseases* 129: 444–452.
- , H. G. WIKER, J. R. DUNCAN, M. GARCÍA, T. W. DUKES, B. W. BROOKS, C. TURCOTTE, AND S. NAGAI. 1990. Protein G-based enzyme-linked immunosorbent assay for antiMPB70 antibodies in bovine tuberculosis. *Journal of Clinical Microbiology* 28: 913–921.
- HIMES, E. M., D. W. LUCHSINGER, J. L. JARNAGIN, C. O. THOEN, H. B. HOOD, AND D. A. FERRIN. 1980. Tuberculosis in fennec foxes. *Journal of the American Veterinary Medical Association* 177: 825–826.
- KANEENE, J. B., AND C. O. THOEN. 2004. Tuberculosis. *Journal of the American Veterinary Medical Association* 224: 685–691.
- LEÓN-VIZCAÍNO, L., A. BERNABÉ, A. CONTRERAS, M. J. CUBERO, S. GÓMEZ, AND R. ASTORGA. 1990. Outbreak of tuberculosis caused by *Mycobacterium bovis* in wild boars (*Sus scrofa*). In *Proceedings: 32 Internationalen Symposiums Erkrankungen der Zoo- und Wildtiere*. ESKILSRUNA. Akademie-Verlag, Berlin, pp. 185–190.
- LIÉBANA, E., A. ARANAZ, B. FRANCIS, AND D. COUSINS. 1996. Assessment of genetic markers for species differentiation within the *Mycobacterium tuberculosis* complex. *Journal of Clinical Microbiology* 34: 933–938.
- MARTÍN-ATANCE, P., F. PALOMARES, M. GONZÁLEZ-CANDELA, E. REVILLA, M. J. CUBERO, J. CALZADA, AND L. LEÓN-VIZCAÍNO. 2005. Bovine tuberculosis in a free-ranging red fox (*Vulpes vulpes*) from Doñana National Park (Spain). *Journal of Wildlife Diseases* 41: 435–436.
- , L. LEÓN-VIZCAÍNO, F. PALOMARES, E. REVILLA, M. GONZÁLEZ-CANDELA, M. J. CUBERO, AND M. DELIBES. 2006. Antibodies to *Mycobacterium bovis* in wild carnivores of Doñana National Park (Spain). *Journal of Wildlife Diseases* 42: 704–708.
- PATRICK, B. 2003. Michigan bovine tuberculosis eradication project activities report 2003. http://www.michigan.gov/documents/BTB_Activities_Report_87802_7.pdf. Accessed 6 May 2007.
- RHYAN, J. C., K. AUNE, R. CLARKE, R. MEYER, C. SIROKY, AND L. STACKHOUSE. 1995. Discovery of bovine tuberculosis in free-ranging mule deer and results of continued wildlife surveillance in the area. In *Proceedings of the Veterinary Epidemiology and Economics Symposium*, US Department of Agriculture, Animal and Plant Health Inspection Service, Centers for Epidemiology and Animal Health, Fort Collins, Colorado, pp. 182–184.
- VICENTE, J., U. HÖFLE, J. M. GARRIDO, I. G. FERNÁNDEZ-DE-MERA, R. JUSTE, M. BARRAL, AND C. GORTAZAR. 2006. Wild boar and red deer display high prevalences of tuberculosis-like lesions in Spain. *Veterinary Research* 37: 107–119.
- WAYNE, L. G., AND G. P. KUBICA. 1986. Family Mycobacteriaceae. In *Bergey's manual of systematic bacteriology*, Vol. 2, 9th Edition, P. H. Sneath, N. S. Mair, M. E. Sharpe, and J. G. Holt (eds.). Williams & Wilkins, Baltimore, Maryland, pp. 1326–1457.

Received for publication 9 May 2007.