

Late Quaternary Glacial and Environmental History of the Burstall Pass Area, Kananaskis Country, Alberta, Canada

Authors: Beierle, Brandon D., Smith, Derald G., and Hills, Leonard V.

Source: Arctic, Antarctic, and Alpine Research, 35(3) : 391-398

Published By: Institute of Arctic and Alpine Research (INSTAAR),
University of Colorado

URL: [https://doi.org/10.1657/1523-0430\(2003\)035\[0391:LQGAEH\]2.0.CO;2](https://doi.org/10.1657/1523-0430(2003)035[0391:LQGAEH]2.0.CO;2)

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Late Quaternary Glacial and Environmental History of the Burstall Pass Area, Kananaskis Country, Alberta, Canada

Brandon D. Beierle,*
Derald G. Smith,† and
Leonard V. Hills‡

*Department of Geography,
Queen's University, Kingston,
Ontario, Canada K7L 3N6.
brandon@lake.geog.queensu.ca

†Department of Geography,
The University of Calgary,
2500 University Dr. N.W., Calgary,
Alberta, Canada T2N 1N4.

‡Department of Geology and Geophysics,
The University of Calgary,
2500 University Dr. N.W., Calgary,
Alberta, Canada T2N 1N4.

Abstract

We used integrated multiproxy analysis of a lake sediment core and glacial geomorphology to reconstruct the late Pleistocene and Holocene climate and geomorphic evolution of the Burstall Pass area, Kananaskis Country, Alberta, Canada. Analysis of macrofossils, pollen, sedimentology, and sediment geochemistry from a lake sediment core and geomorphology and tephrochronology of glacial moraines provide evidence for multiple modes of climate during the last ca. 11,000 yr. An advance of the Robertson Glacier prior to ca. 9200 ^{14}C BP is correlated to the Crowfoot advance and was the largest of the postglacial period. Immediately following this event, increased lake productivity and the deposition of marl as well as increased arboreal/nonarboreal pollen (AP/NAP) ratios suggest that the climate warmed, possibly accompanied by increased aridity. Decreased turbidity and clastic sediment flux in Lower Burstall Lake during the early Holocene suggest reduced glacial runoff and may indicate the complete ablation of the Robertson Glacier shortly after 10,000 ^{14}C BP. Clastic sediment flux to Lower Burstall Lake remained minimal until after ca. 3500 ^{14}C BP, when decreasing LOI (loss-on-ignition) organic carbon levels in lake sediments signaled the return of glacial runoff to the lake system. The largest Neoglacial advance in the Burstall Pass area appears to have been the most recent and was followed by rapid recession during the 20th century.

Introduction

Previous paleoclimatic studies examining the late Pleistocene and Holocene glacial history of the front and middle ranges of the Southern Canadian Rocky Mountains have identified three major periods of increased glacial activity since the late Wisconsinan. On the basis of stratigraphic relationships between moraines and tephrochronology, the earliest of these advances, the Crowfoot, was initially thought to be of early Holocene or late Wisconsinan age (Luckman and Osborn, 1979). More recent analysis of lake sediment cores and moraine stratigraphy has suggested that this advance occurred during the Younger Dryas (Osborn and Luckman, 1988; Reasoner et al., 1994). Neoglacial activity in the Canadian Rockies is thought to have begun with the Peyto advance at ca. 3000 ^{14}C BP (Luckman et al., 1993; Leonard and Reasoner, 1999), which was followed by early and later Little Ice Age advances beginning at ca. 800 ^{14}C BP and lasting until the beginning of the 20th century (Luckman, 2000). Pollen and diatom studies from lakes in the same area suggest that treeline elevations were higher during the early Holocene and have decreased through the past 4000 yr, with cooler, wetter conditions becoming predominant (MacDonald, 1982; Kearney and Luckman, 1983; Luckman and Kearney, 1986; Hickman and Reasoner, 1998; Reasoner and Huber, 1999).

The paleoenvironmental chronology presented in this study combines sedimentological and palynological analysis of a lake sediment core with geomorphic reconstructions of past glacial advances. This approach provides a direct link between vegetation, sedimentary, and glacial responses to past climatic changes, allowing more detailed reconstruction. This work also provides an additional minimum age for the Crowfoot advance, strengthening previous evidence suggesting that the type section Crowfoot moraine is of late Pleistocene age and possibly correlative to the Younger Dryas (Reasoner et al., 1994). Paleo-equilibrium-line altitude (ELA) reconstructions provide an estimate for the magnitude of climatic change at the termination of the Crowfoot advance.

Regional Setting

The Burstall Valley is a tributary to the Spray River valley in Kananaskis Country, Alberta, approximately 100 km west of Calgary (Fig. 1). The hydrology of the valley is dominated by meltwater from the Robertson Glacier, which lies between 2500 and 2750 m a.s.l., on the north side of a col at the head of a deeply incised glacial valley southwest of Burstall Pass (Fig. 2). The ELA of the Robertson Glacier was estimated at 2600 m a.s.l., based on aerial photographs taken 3 October 1993. Downvalley 1.5 km from the modern glacier terminus, a pair of moraines record two post-Late Wisconsinan advances of the Robertson Glacier (Figs. 2 and 3a).

Lower Burstall Lake is fed by meltwater from the Robertson Glacier via two more proximal basins (upper and middle Burstall Lakes; Fig. 2). Because the upper two lakes act as sediment traps, both the total influx and particle size of sediment entering Lower Burstall Lake are restricted, resulting in a relatively shallow sedimentary fill that is uninterrupted by density flows or other sedimentary events associated with environments having high sedimentation rates. Meltwater from the Robertson Glacier (via the upper two lakes) is the only major inflow to Lower Burstall Lake (Fig. 2). Lower Burstall Lake (1970 m a.s.l.) is currently below treeline (2450 m a.s.l.) and is surrounded by a mixed *Picea engelmannii* and *Abies lasiocarpa* forest.

Methods

We took sediment cores from Lower Burstall Lake with a percussion coring system (Reasoner, 1993) in 8-m water depth, using the frozen lake surface as a platform. The core LBSL1 penetrated 290 cm into the lake bed before being stopped by dense, clay-rich sediment. The cores were frozen and cut in half longitudinally with a diamond rock saw. Moraines and drainage were mapped, both in the field and by analysis of aerial photographs.




FIGURE 1. Location of Burstall Pass/Lower Burstall Lake area (1.) and locations of other lakes mentioned in text (2. Upper Kananaskis Lake, 3. Lake O'Hara, 4. Lake Louise, 5. Hector Lake, 6. Crowfoot Lake, 7. Bow Lake).

Analysis of the lake sediment cores included total organic carbon (TOC) and total inorganic carbon (TIC) by loss-on-ignition (Bengtsson and Enull, 1986) as well as lithostratigraphic, macrofossil, and pollen analyses. Pollen samples were taken at 10-cm intervals, prepared using standard treatment (Faegri and Iversen, 1989), and spiked with lycopodium spores, with an average of 300 grains counted per sample. Macrofossils and sand grains ($>175\ \mu\text{m}$) were counted at 10-cm intervals by wet sieving 1-cm-thick slices of core ($\sim 45\ \text{cm}^3$). Raw counts were then normalized to $1\ \text{cm}^3$. An accelerator mass spectrometer (AMS) radiocarbon date was obtained from a single conifer needle (*Picea* sp.) sieved from the center of the core at 242-cm depth. The Mazama (ca. 6730 ^{14}C BP) and Bridge River (ca. 2400 ^{14}C BP) tephra provided additional chronological control (Clague et al., 1995; Hallett et al., 1997), and allowed the creation of an age-depth model and calculation of sedimentation rates by linear interpolation (Fig. 4). Tephra in both the lake core and soil pit were identified on the basis of petrographic morphology according to the criteria suggested by Reasoner and Healy (1986), stratigraphic position where appropriate, and laboratory comparison to known reference samples of Mazama, Bridge River, and St. Helen's Y tephra. All dates are quoted in radiocarbon years BP (^{14}C BP), and the thickness of the Mazama tephra was removed from the age-depth model (Fig. 4). Core zonation was done primarily on the basis of visual lithostratigraphy as well as the pollen, TOC, and TIC data (Figs. 5 and 6).

Results

GEOMORPHOLOGY

The outermost of the two moraines downvalley from the Robertson Glacier (Moraine 1) is rounded in profile and well vegetated (Fig. 3a), with both arboreal and nonarboreal species established on the crest. Mazama tephra was found near the base of a 20-cm soil profile (Fig. 3b) atop the crest of Moraine 1. The location of the tephra on the apex of the moraine precludes secondary deposition, thereby providing a minimum age for Moraine 1 of ca. 6730 ^{14}C BP. Although Mazama is the only tephra plume mapped as crossing this area, Bridge River tephra and St. Helen's Y tephra were identified in a section 15 km northeast of the Burstall Pass Area (Limbird, 1984). Despite the presence of Bridge River tephra in the core from Lower Burstall Lake, neither it nor St. Helen's Y tephra were found on either moraine.

The proximal lateral moraine (Moraine 2) is sharp-crested and pushed into the flank of Moraine 1. Soil is mostly absent from the surface, with vegetation limited to scattered willows, and subalpine fir seedlings growing in small pockets of fine-grained sediment (Fig. 3a). The flanks of Moraine 2 are oversteepened and incised with small rills and gullies. Small recessional ridges within the perimeter of Moraine 2, in the glacier forefield, are rounded, less than 2 m high in profile, and mostly devoid of vegetation. Because there are no other terminal moraines located inside these recessional features, we assumed that the




FIGURE 2. Contour map of the Burstall Pass area showing the Robertson Glacier, lateral and terminal moraines, and the locations of the soil pit in Moraine 1 and the coring site in Lower Burstall Lake.

Moraine 2 represents the most recent Holocene advance of the Robertson Glacier. Following this advance, the Robertson Glacier retreated approximately 1.5 km to its current position (Fig. 2).

LAKE CORE

The 290-cm-long sediment core obtained from Lower Burstall Lake contained three prominent lithostratigraphic zones, consisting of a basal inorganic silt and clay unit (Zone 1), an organic-rich marl and silt unit (Zone 2), and a transitional organic-rich to inorganic silt unit (Zones 3a and b, respectively) (Fig. 5). Analysis of LOI organic and inorganic carbon content, pollen, and macrofossils (Figs. 5 and 6) indicate that zonation based on visual sedimentology matches well with geochemical and biological changes in the sediments.

Zone 1: 290–246 cm

Sedimentation in Zone 1 was dominated by massive to laminated sand, silt, and clay. Sieving Zone 1 revealed small concentrations of coarse sand in several intervals. TOC in Zone 1 was consistently below

5% (Fig. 5), and aquatic or terrestrial macrofossils were absent. Pollen was very sparse in Zone 1 sediments as compared to Zone 2, requiring as many as five slides (compared to less than one for Zone 2) to count ~300 pollen grains. The pollen assemblage in Zone 1 was dominated by xeric herbaceous and shrub taxa, with very little arboreal representation (Fig. 6).

Zone 2: 246–160 cm

An abrupt, conformably bedded change to organic-rich marl at 246 cm marked the boundary between Zones 1 and 2, suggesting a rapid change in sedimentary regime. The marl, 4 cm thick, ended in a sharp transition to organic-rich silt. TOC in Zone 2 increased to 10% and remained above 5% throughout the zone (Fig. 5). Molluscs of the genus *Pisidium* were found in concentrations of 55 valves/100 cc within the marl but declined rapidly to 7 valves/100 cc within the first few centimeters of the organic-rich silt before disappearing altogether by the 225-cm interval. Also present within the marl were very high concentrations of charophyte oogonia. The AP/NAP ratio increased from less than 1 in Zone 1 to 3 at the base of Zone 2, with *Pinus*




FIGURE 3. (a) The paired lateral moraines downvalley from Robertson Glacier showing the difference in vegetation cover between the older Moraine 1 (left) and younger Moraine 2. (b) Soil pit in older (left) moraine.

increasing to more than 60% relative abundance (Fig. 6). *Alnus* pollen decreased sharply across this boundary, as did herbaceous pollen and pollen from several other shrubs. A brief decrease in the AP/NAP ratio and a concentration of sub millimeter charcoal macrofossils at 230 cm corresponded to the sedimentary change from marl to organic-rich silt (Figs. 5 and 6). An AMS radiocarbon date of 9180 ± 60 ^{14}C BP (CAMS-20358) was obtained from a *Picea* needle fragment found 4 cm above the Zone 1–2 boundary. The Mazama tephra (ca. 6730 ^{14}C BP; Hallett et al., 1997), between 210 and 218 cm, also provided chronological control for this zone (Fig. 4). The average sedimentation rate between the radiocarbon date and Mazama tephra was the lowest in the core at 0.9 mm yr^{-1} .




FIGURE 4. Age depth curve based on linear interpolation between points of known age in the core LBSL2 from Lower Burstall Lake. Note that the thickness of the Mazama tephra was removed from the calculations.

Zone 3: 160–0 cm

The transition from Zone 2 to Zone 3 was gradational and characterized by decreased TOC and a sharp decline in the AP/NAP ratio. Sedimentation in Zone 3 was dominated by carbonate silt and clay, becoming less organic and more carbonate rich up core. Zone 3 contained no macrofossils and averaged less than 5% TOC throughout (Fig. 5). On the basis of TOC and AP/NAP ratios, Zone 3 was divided into two subzones, Zones 3a (160–90 cm) and 3b (90–0 cm) (Figs. 5 and 6). The Bridge River tephra, dated at ca. 2400 ^{14}C BP, was found at the 146-cm interval of Zone 3a. Sedimentation rates in Zone 3 were 1.7 mm yr^{-1} below the Bridge River tephra and 6 mm yr^{-1} above it. The abrupt nature and precise timing of this change in sedimentation rate are a function of the location of the dating control point.

Zone 3a: 160–90 cm

Beginning at about 160 cm, *Pinus* pollen declined abruptly and was replaced by increasing amounts of *Alnus*, *Salix*, *Shepherdia*, and several herbaceous pollen taxa. These changes resulted in decreased AP/NAP ratios, with levels similar to those in Zone 1 at 120 cm (Fig. 6). LOI organic carbon declined steadily from the base of Zone 3a, stabilizing at $\sim 5\%$ above the 120-cm interval. Carbonate values increased from the base of Zone 3a until 130 cm, reaching $\sim 20\%$ at the transition to Zone 3b.

Zone 3b: 90–0 cm

At the base of Zone 3b, the AP/NAP ratio increased as *Pinus* pollen became more dominant and herb and shrub pollen, which dominated Zone 3a, declined significantly (Fig. 6). Following a peak at 80 cm, the AP/NAP ratio declined steadily with herb and shrub pollen becoming more dominant and *Pinus* pollen decreasing by 20%. Herbaceous pollen remained at increased levels until the 20 cm interval of the core, where the AP/NAP ratio increased, reflecting decreased importance of *Alnus* and herbaceous pollen taxa (Fig. 6). LOI organic carbon continued to decrease above 90 cm, reaching the lowest values of the Holocene in the upper 10 cm of the core. Carbonate increased steadily after 90 cm, reaching a Holocene maximum value of $\sim 25\%$ in the topmost sample of the core.

Discussion

LATE PLEISTOCENE (>10,000 ¹⁴C BP)

The presence of the Mazama tephra, which is found within the soil profile on the surface of Moraine 1, shows that Moraine 1 is older than ca. 6730 ¹⁴C BP (Fig. 3b). This date indicates that it is likely of similar age to a pre-Mazama aged moraine at the Crowfoot Glacier, identified by Luckman and Osborn (1979) and named the Crowfoot advance. Because the only period of clastic sedimentation stratigraphically below the Mazama tephra in the core from Lower Burstall Lake occurs in Zone 1, below 246-cm depth, the radiocarbon date at 242-cm depth provides a minimum age for the advance of 9180 ± 60 ¹⁴C BP (Zone 1; Fig. 5). Because this date comes from 4 cm above the onset of organic sedimentation, it underestimates the age of the basal clastic sediments, likely by several hundred years. On this basis, it is probable that the transition from clastic to organic sediments in Lower Burstall Lake is correlative to similar transitions in numerous lakes in the Southern Canadian Rocky Mountains that occurred at ca. 10,000 ¹⁴C BP (e.g., Reasoner and Hickman, 1989; Beierle, 1997; Beierle and Smith, 1998; Reasoner and Huber, 1999).

The age of this basal inorganic sediment package is also similar to basal inorganic sediments from Crowfoot Lake that were deposited between 10,100 ¹⁴C BP and 11,300 ¹⁴C BP and have been correlated to the Younger Dryas cold period (Reasoner et al., 1994). The dominantly herbaceous pollen assemblage in Zone 1 supports this inference and suggests that cold, dry conditions prevailed during this interval. Although recent work has cast some doubt on the validity of the correlation between the Crowfoot type section moraine and the basal sediments from Crowfoot Lake (Leonard, 1998), the apparent synchrony of the Crowfoot event in the Rocky Mountains (this paper; Reasoner et al., 1994; Reasoner and Jodry, 2000) and the Younger Dryas in the Greenland ice cores (Alley, 2000) indicates that the two may represent the same event. Evidence for a significant Younger Dryas climatic event in western North America is also supported by recent work from both New Mexico (Armour et al., 2002) and British Columbia (Kovanen and Easterbrook, 2002) which also suggest the occurrence of major glacial advances during the late Pleistocene.

EARLY TO MIDDLE HOLOCENE (CA. 10,000 ¹⁴C BP–3500 ¹⁴C BP)

Clastic sediment flux decreased abruptly in Lower Burstall Lake at ca. 10,000 ¹⁴C BP and remained low until ca. 3500 ¹⁴C BP (Zone 2; Fig. 5). Clastic sediment was replaced by organic-rich silt and marl during this interval, suggesting a major change in depositional environment. Marl deposition in this and other glacier-fed lakes in the southern Canadian Rocky Mountains between ca. 10,000 and 9000 ¹⁴C BP (e.g., Reasoner et al., 1994; Beierle, 1997) suggests that lake water was warmer and may have been thermally unstratified, because marl accumulated on the floor of the lake, rather than redissolving at depth due to increased acidity often associated with anoxic bottom conditions and stratified lake waters. Higher rates of algal photosynthesis inferred by increased LOI organic carbon levels suggest decreased water turbidity during the early Holocene, as does the presence of gill-breathing benthic molluscs (Fig. 5). Gill-breathing benthic fauna also show that oxic conditions likely prevailed in bottom waters during this interval. Marl deposition terminated abruptly at ca. 9000 ¹⁴C BP, after which molluscs also disappeared from the sedimentary record. This change may indicate seasonal stratification of lake waters, with anoxic conditions below the thermocline resulting in redissolution of carbonate solids and the extinction of some benthic fauna. Continued organic sedimentation above the marl suggests that turbidity and clastic sediment influx remained low until ca. 3500 ¹⁴C BP. Increased arboreal pollen percentages, particularly *Pinus* (Fig. 6),




FIGURE 5. Lithostratigraphy, LOI carbonate and organic carbon, and *Pisidium* macrofossils from Lower Burstall Lake.

may indicate that treeline was higher during the early Holocene, probably as a result of a warmer climate. Many other records from this region also indicate that treeline was higher than at present during the early Holocene (e.g., Kearney and Luckman, 1983; Luckman and Kearney, 1986; Reasoner and Hickman, 1989), although this record is unique in that there are several dramatic drops in the AP/NAP ratio in Zone 2, which may indicate the establishment of more open valley-bottom forests as treeline moved to even higher elevations. This interpretation is corroborated by increased abundances of xerophytic pollen taxa during these intervals at ca. 8000 ¹⁴C BP and between 5000 and 6000 ¹⁴C BP.

Increased water temperature and decreased water column turbidity suggest that glacial influence in Lower Burstall Lake was substantially decreased during the early Holocene. Similar early Holocene records from Crowfoot Lake, Upper Kananaskis Lake, Lake Louise, Lake O'Hara, and Bow Lake (Reasoner and Hickman, 1989; Reasoner et al., 1994; Beierle, 1997; Beierle and Smith, 1998; Leonard and Reasoner, 1999) also contain evidence for significantly decreased glacial sediment input, with the presence of organic sediments, marl, and molluscs suggesting that glacial ice may have completely disappeared in some locations. Sediments from Hector Lake also suggest decreased glacial activity but indicate that the Waputik Ice Field, although reduced in area, continued to exist through the early Holocene (Leonard and Reasoner, 1999). This may have been the case because the Waputik Ice Field is one of the highest glaciers in this area of the




FIGURE 6. Lithostratigraphy and relative pollen percentages from Lower Burstall Lake.

Canadian Rockies, at 3240 m a.s.l. The regional picture of decreased glacierization is also supported by detrital wood dating to the early Holocene discovered at the terminus of the Athabasca and Dome Glaciers (Luckman, 1988; Luckman et al., 1993), indicating that these glaciers were substantially less extensive during the early Holocene.

LATE HOLOCENE (CA. 3500 ¹⁴C BP TO PRESENT)

Whereas the Zone 1–2 boundary is sharp, the sedimentological change from Zone 2 to Zone 3 is gradual (Fig. 5). LOI organic carbon values began to decline at ca. 3500 ¹⁴C BP and decreased steadily through the late Holocene. AP/NAP values dropped rapidly just prior to ca. 3500 ¹⁴C BP and continued to decline through ca. 2000 ¹⁴C BP, remaining at almost pre-Holocene levels until ca. 1300 ¹⁴C BP (Fig. 6). Zone 3a likely represents the onset of the Neoglacial, with decreased AP/NAP values suggesting either reduced treeline elevations or disruption of local forest communities by reactivation of the Robertson Glacier outwash plain, which occupies much of the valley bottom area upstream of Lower Burstall Lake (Fig. 2). Either of these hypotheses is consistent with decreased LOI organic carbon, increased LOI carbonate values, and increased sedimentation rates, which suggest renewed glacial activity in the catchment (Fig. 5).

AP/NAP values remained low until ca. 1300 ¹⁴C BP, when increased arboreal pollen percentages suggest that spruce and pine increased in abundance (Zone 3b; Fig. 6). This interval had the highest

arboreal pollen content of the core, suggesting improved growing conditions over those in Zone 3a (Fig. 6). After this initial increase, arboreal pollen again declined in importance, although more slowly this time, reaching a minimum at ca. 400 ¹⁴C BP (Fig. 6). After ca. 400 ¹⁴C BP, the AP/NAP ratio increased until ca. 200 ¹⁴C BP before declining into modern times, likely in response to renewed glacial activity during the Little Ice Age (Fig. 6). Although it is uncertain whether these changes in AP/NAP ratio indicate movement of treeline or changes in local vegetation dynamics, periods with decreased arboreal pollen abundances during the last 1300 yr may correspond with Neoglacial and Little Ice Age advances proposed by Luckman et al. (1993), Smith et al. (1995), and Luckman (2000), although the chronology through this portion of the core lacks sufficient resolution for these correlations to be more than speculative.

Because LOI carbon and carbonate remained stable through the late Holocene, we suggest that the lacustrine sedimentary record in Lower Burstall Lake is not as sensitive to small-scale fluctuations in climate as the pollen record (Figs. 5 and 6). Increased clastic flux to the basin after ca. 3500 ¹⁴C BP is likely indicative of renewed glacierization in a general sense, as opposed to recording specific glacial events. Thus, the multiple advances thought to have occurred during the Neoglacial were likely blended together within the clastic sedimentary record.

Despite the minimal resolution in the lacustrine sedimentary record during the late Holocene, the lack of smaller terminal moraines within Moraine 2 at the Robertson Glacier suggest that it was most

likely produced during the Little Ice Age. Although this moraine is undated, its morphology and size are consistent with known Little Ice Age moraines in the area. This moraine is sharp-crested, with oversteepened flanks, similar to the Little Ice Age moraine described by Smith et al. (1995) at the adjacent French Glacier. In addition, the lack of turf or large trees makes it likely that Moraine 2 is relatively recent compared to older Neoglacial moraines at the nearby French Glacier that have been dated to ca. 500 ^{14}C BP (Smith et al., 1995) and at other locations in the Southern Canadian Rockies (Luckman and Osborn, 1979). Because the small ridges in the forefield of the Robertson Glacier are poorly formed and lack the characteristic sharp crest common in terminal moraines, they are interpreted as recessional features deposited during ice recession following the advance responsible for depositing the inner moraine (Moraine 2). McCarthy and Smith (1994) used historical photographs to document the retreat of many glaciers in the Southern Canadian Rocky Mountains, including the Robertson, and showed that it has retreated more than 1.2 km since 1916, losing 35% of its area.

Conclusion

We suggest that Holocene climatic and landscape evolution in the Burstall Pass area occurred in three stages, beginning with an advance of the Robertson Glacier prior to ca. 10,000 ^{14}C BP, which is correlated to the Crowfoot Advance and may represent cooling during the Younger Dryas. This advance deposited lateral and end moraines about 1.5 km downvalley from the modern glacier terminus correlating to an ELA depression of ~ 300 m relative to the current ELA. The magnitude of the advances depositing Moraines 1 and 2 was almost equal, suggesting either that the ELA depression for both the Younger Dryas and Little Ice Age may have been similar in this region or that the glacier may not have had time to equilibrate to changes in ELA during one or both advances.

Following this advance, the termination of clastic sediment flux to Lower Burstall Lake suggests that abrupt climatic warming at ca. 10,000 ^{14}C BP resulted in rapid retreat of the Robertson Glacier, and possibly its complete ablation. Treeline may also have been higher after ca. 10,000 ^{14}C BP, and reconstructions of paleo-ELA using the 70% area method (Bradley, 1999) indicate that the ELA of the Robertson Glacier may have risen by as much as 440 m at ca. 10,000 ^{14}C BP.

This period of limited glacial activity lasted until ca. 3500 ^{14}C BP, when arboreal pollen levels suggest that treeline elevation began to decrease and declining LOI organic carbon values suggest declining lake productivity and increased glacial sediment flux. Palynological analysis indicates that the period between 3500 and 1300 ^{14}C BP was colder, with renewed glacial activity resulting in increased clastic sediment flux to the lake. Although the clastic sediment record from Lower Burstall Lake cannot resolve multiple Neoglacial advances, fluctuations in arboreal pollen levels through the last 2000 radiocarbon yr may correlate to Late Holocene glacial events noted by Luckman and Osborn (1979) and Smith et al. (1995). Warmer climate after ca. 1300 ^{14}C BP is suggested by increased representation of arboreal pollen, although the almost steady decline in LOI organic carbon and steady increase in LOI carbonate after ca. 3500 ^{14}C BP suggest that climatic fluctuations after this time were also too subtle to be reflected in the bulk physical properties of the sediment. The younger moraine at the Robertson Glacier is most likely associated with the last advance of the Little Ice Age during the mid- to late 19th century, but it remains undated.

Acknowledgments

The authors would like to thank David Mazzucchi and Natalie Veldhoen for their assistance in the field. Mel Reasoner is also thanked

for his field assistance and for providing the radiocarbon date. Laboratory facilities were provided by the Departments of Geography, The University of Calgary, and Queen's University. Julian Szeicz provided laboratory space, equipment, and assistance with the pollen analysis. Markus Heinrichs, Robert Gilbert, Scott Lamoureux, Eric Leonard, Mel Reasoner, and an anonymous reviewer provided valuable critical reviews of this manuscript.

References Cited

- Alley, R. B., 2000: The Younger Dryas cold interval as viewed from central Greenland: past global changes and their significance for the future. *Quaternary Science Reviews*, 19: 213–226.
- Armour, J., Fawcett, P.J. and Geissman, J.W., 2002: 15 k.y. paleoclimatic and glacial record from northern New Mexico. *Geology*, 30: 723–726.
- Beierle, B. D., 1997: Early Holocene climate of southwestern Alberta, Canada, reconstructed from lake sediment cores. Master's thesis, Department of Geography, The University of Calgary, Calgary, Alberta, Canada. 115 pp.
- Beierle, B., and Smith, D. G., 1998: Severe drought in the early Holocene (10,000–6800 BP) interpreted from lake sediment cores, southwestern Alberta, Canada. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 140: 75–83.
- Bengtsson, L., and Ennall, M., 1986: Chemical Analysis. In Berglund, B. E. (ed.), *Handbook of Holocene Palaeoecology and Palaeohydrology*. Chichester: John Wiley and Sons Ltd., 423–451.
- Bradley, R. S., 1999: *Paleoclimatology*, 2nd ed. San Diego: Academic Press. 613 pp.
- Clague, J. J., Evans, S. G., Rampton, V. N., and Woodsworth, G. J., 1995: Improved age estimates for the White River and Bridge River tephra, western Canada. *Canadian Journal of Earth Sciences*, 32: 1172–1179.
- Fægri, K., and Iversen, J., 1989: *Textbook of Pollen Analysis*, 4th ed. Chichester: John Wiley and Sons. 328 pp.
- Hallett, D. J., Hills, L. V., and Clague, J. J., 1997: New accelerator mass spectrometry radiocarbon ages for the Mazama tephra layer from Kootenay National Park, British Columbia, Canada. *Canadian Journal of Earth Sciences*, 34: 1202–1209.
- Hickman, M., and Reasoner, M. A., 1998: Late Quaternary diatom response to vegetation and climate change in a subalpine lake in Banff National Park, Alberta. *Journal of Paleolimnology*, 20: 253–265.
- Kearney, M. S., and Luckman, B. H., 1983: Postglacial vegetational history of Tonquin Pass, British Columbia. *Canadian Journal of Earth Sciences*, 20: 776–786.
- Kovnen, D.J. and Easterbrook, D.J., 2002: Timing and extent of Allerød and Younger Dryas Age (ca. 12,500–10,000 ^{14}C YR B.P.) Oscillations of the cordilleran Ice Sheet in the Fraser Lowland, western North America. *Quaternary Research*: 57: 208–224.
- Leonard, E. M., 1998: Is the type-Crowfoot Moraine, Alberta, really a Younger Dryas equivalent? *Geological Society of America, 1998 Annual Meeting*, 30: 66.
- Leonard, E. M., and Reasoner, M. A., 1999: A continuous Holocene glacial record inferred from proglacial lake sediments in Banff National Park, Alberta, Canada. *Quaternary Research*, 51: 1–13.
- Limbird, A., 1984: Genesis of soils affected by discrete volcanic ash inclusions, Alberta, Canada. *Catena*, Selected Papers of the Congreso Internacional de Seulos Volcanicos: 119–130.
- Luckman, B. H., 1988: 8000 year old wood from the Athabasca Glacier, Alberta. *Canadian Journal of Earth Sciences*, 25: 148–151.
- Luckman, B. H., 2000: The Little Ice Age in the Canadian Rockies. *Geomorphology*, 32: 357–384.
- Luckman, B. H., and Kearney, M. S., 1986: Reconstruction of Holocene changes in alpine vegetation and climate in the Maligne Range, Jasper National Park, Alberta. *Quaternary Research*, 26: 244–261.

- Luckman, B. H., and Osborn, G. D., 1979: Holocene glacier fluctuations in the middle Canadian Rocky Mountains. *Quaternary Research*, 11: 52–77.
- Luckman, B. H., Holdsworth, G., and Osborn, G. D., 1993: Neoglacial glacier fluctuations in the Canadian Rockies. *Quaternary Research*, 39: 144–153.
- MacDonald, G. M., 1982: Late Quaternary paleoenvironments of the Morley Flats and Kananaskis Valley of southwestern Alberta. *Canadian Journal of Earth Sciences*, 19: 23–25.
- McCarthy, D. P., and Smith, D. J., 1994: Historical glacier activity in the vicinity of Peter Lougheed Provincial Park, Canadian Rocky Mountains. *Western Geography*, 4: 94–109.
- Osborn, G. D., and Luckman, B. H., 1988: Holocene glacier fluctuations in the Canadian Cordillera (Alberta and British Columbia). *Quaternary Science Reviews*, 7: 115–128.
- Reasoner, M. A., and Healy, R. E., 1986: Identification and significance of tephra encountered in a core from Mary Lake, Yoho National Park, British Columbia. *Canadian Journal of Earth Sciences*, 23: 1991–1999.
- Reasoner, M. A., 1993: Equipment and procedure improvements for a lightweight, inexpensive, percussion core sampling system. *Journal of Paleolimnology*, 8: 273–281.
- Reasoner, M. A., and Hickman, M., 1989: Late Quaternary environmental change in the Lake O'Hara region, Yoho National Park, British Columbia. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 72: 291–316.
- Reasoner, M. A., and Huber, U. M., 1999: Postglacial palaeoenvironments of the upper Bow Valley, Banff National Park, Alberta, Canada. *Quaternary Science Reviews*, 18: 475–492.
- Reasoner, M. A., and Jodry, M. A., 2000: Rapid response of alpine timberline vegetation to the Younger Dryas climate oscillation in the Colorado Rocky Mountains, U.S.A. *Geology*, 28: 51–54.
- Reasoner, M. A., Osborn, G. D., and Rutter, N. W., 1994: Age of the Crowfoot advance in the Canadian Rocky Mountains: a glacial event coeval with the Younger Dryas oscillation. *Geology*, 22: 439–442.
- Smith, D.J., McCarthy, D.P., and Colenutt, M.E., 1995: Little Ice Age glacial activity in Peter Lougheed and Elk Lakes provincial parks, Canadian Rocky Mountains. *Canadian Journal of Earth Sciences*, 32: 579–589.

Ms submitted January 2002

Revised ms submitted September 2002