

## **Stable Carbon Isotope Compositions of Eastern Beringian Grasses and Sedges: Investigating Their Potential as Paleoenvironmental Indicators**

Authors: Wooller, Matthew J., Zazula, Grant D., Edwards, Mary, Froese, Duane G., Boone, Richard D., et al.

Source: Arctic, Antarctic, and Alpine Research, 39(2) : 318-331

Published By: Institute of Arctic and Alpine Research (INSTAAR), University of Colorado

URL: [https://doi.org/10.1657/1523-0430\(2007\)39\[318:SCICOE\]2.0.CO;2](https://doi.org/10.1657/1523-0430(2007)39[318:SCICOE]2.0.CO;2)

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# Stable Carbon Isotope Compositions of Eastern Beringian Grasses and Sedges: Investigating Their Potential as Paleoenvironmental Indicators

Matthew J. Wooller\*

Grant D. Zazula†

Mary Edwards‡

Duane G. Froese§

Richard D. Boone#

Carolyn Parker\$ and

Bruce Bennett@

\*Alaska Stable Isotope Facility, Water and Environmental Research Center and School of Fisheries and Ocean Sciences, Duckering Building, University of Alaska, Fairbanks, Alaska 99775-5860, U.S.A.

ffmjw@uaf.edu

†Department of Biological Sciences, Simon Fraser University, 8888 University Drive, Burnaby, British Columbia, V5A 1S6, Canada

‡School of Geography, University of Southampton, Room 2019 Shackleton Building, Highfield, Southampton, SO17 1BJ, U.K.

§Department of Earth and Atmospheric Sciences, University of Alberta,

Edmonton, Alberta, T6G 2E3, Canada

#Institute of Arctic Biology, University of Alaska, Fairbanks, Alaska 99775-7000, U.S.A.

\$Herbarium, University of Alaska

Museum of the North, University of Alaska, Fairbanks, Alaska 99775-6960, U.S.A.

@Nature Serve Yukon, V5N #10, Burns Road, Whitehorse, Yukon Territory, Y1A 2C6, Canada

## Abstract

The nature of vegetation cover present in Beringia during the last glaciation remains unclear. Uncertainty rests partly with the limitations of conventional paleoecological methods. A lack of sufficient taxonomic resolution most notably associated with the grasses and sedges restricts the paleoecological inferences that can be made. Stable isotope measurements of subfossil plants are frequently used to enhance paleoenvironmental reconstructions. We present an investigation of the stable carbon isotope composition ( $\delta^{13}\text{C}$ ) of modern and subfossil grasses and sedges (graminoids) from Eastern Beringia. Modern grasses from wet habitats had a mean  $\delta^{13}\text{C}$  of  $-29.1\text{‰}$  (standard deviation [SD] =  $2.1\text{‰}$ ,  $n = 75$ ), while those from dry habitats had a mean of  $-26.9\text{‰}$  (SD =  $1.19$ ,  $n = 27$ ). Sedges ( $n = \sim 50$ ) from dry, wet, marsh, and sand dune habitats had specific habitat ranges. Four modern  $\text{C}_4$  grasses had  $\delta^{13}\text{C}$  values typical of  $\text{C}_4$  plants. Analyses were also conducted using subfossil graminoid remains from several sedimentary paleoecological contexts (e.g., arctic ground squirrel nests, loess, permafrost, and paleosols) in Eastern Beringia. Results from these subfossil samples, ranging in age from  $>40,000$  to ca.  $11,000$  cal. yr BP, illustrate that the  $\delta^{13}\text{C}$  of graminoid remains has altered during the past. The range of variation in the subfossil samples is within the range from modern graminoid specimens from dry and wet habitats. The results indicate that stable isotopes could contribute to a comprehensive and multiproxy reconstruction of Beringian paleoenvironments.

## Introduction

An accurate reconstruction of the glacial-age vegetation of Beringia (the largely unglaciated region lying between the Lena River in the west and the Mackenzie River in the east) (for a recent discussion of the extent and definitions of Beringia, see Harington, 2005) (Fig. 1) is a key element in several areas of scientific investigation. This region was a subcontinental Pleistocene refugium for northern biota (Hultén, 1937) and the exposed Bering land bridge allowed migrations between northeastern Asia and North America. The structure, composition, and productivity of the vegetation of the region would have influenced animal population dynamics, species distributions, and the timing and routes of continental biotic exchange (Hopkins et al., 1982). In addition, land-atmosphere interactions likely contributed (via feedbacks) to the climate changes of the late Quaternary. The challenge of modeling land-cover feedbacks in the context of

potential future climate change is partly addressed by testing the capacity of climate models to simulate past conditions that differ markedly from those at present (Paleoclimate Modelling Intercomparison Project, <http://www-lsce.cea.fr/pmip/>; Kaplan et al., 2003). A critical time for such tests is the height of the Last Glacial Maximum (LGM), ca.  $21,000$  cal. yr BP, referred to as Marine Isotope Stage (MIS) 2 (Kageyama et al., 2001). However, the nature of the vegetation across Beringia during the last glaciation is unclear, both in terms of large-scale gradients related to temperature and moisture and in the range of variability associated with the topographic and topoclimatic mosaic. The continuing uncertainty rests partly with the logistical challenges of studying this remote region. The uncertainty also stems from the limitations of conventional paleoecological methods to resolve the vegetation taxonomically to the degree necessary to infer its bioclimatic requirements and ecological characteristics.

During Pleistocene glaciations, Eastern Beringia (unglaciated Yukon Territory and Alaska) was isolated from North America by continental ice sheets and became biogeographically continuous with northeast Asia (Hopkins et al., 1982; Shapiro et al., 2004). However, there exists a long-running and incompletely resolved dilemma: the so-called “steppe-tundra debate,” or “productivity paradox,” which is by now well rehearsed in the literature (e.g., Cwynar and Ritchie, 1980; Hopkins et al., 1982; Guthrie, 1990; Colinvaux, 1996a; Kozhevnikov and Ukraintseva, 1999; Zazula et al., 2003). Logic suggests that a region spanning ~60–75°N, when LGM global temperatures were colder than today, was most probably covered by some form of tundra (i.e., vegetation occurring where low-growing season temperatures do not permit tree growth; Ritchie, 1984; Viereck et al., 1992). However, numerous dated faunal remains (Guthrie, 1968, 1990; Harington, 2003), and even ancient DNA (e.g., Shapiro and Cooper, 2003; Willerslev et al., 2003), attest to the presence of a diversity of herbivorous and carnivorous mammals throughout the last glaciation. While tundra can take many forms, most contemporary tundra types are unpromising analogs for vegetation that could have supported populations of large grazers, including mammoths, horses, and bison. Thus, Guthrie (1968, 1990, 2001) argued that the ancient Beringian vegetation was more akin to a modern northern steppe (i.e., grassland vegetation where it is too dry for trees) such as that found today in portions of northern and central Asia. Hence there arose the concept of steppe-tundra (tundra-steppe in the Russian literature), which importantly indicates that the environment was both dry and cold (e.g., Yurtsev, 1982). However, steppe-tundra is a flexible enough concept to allow the continuance of a range of interpretations (Colinvaux, 1980a, 1980b, 1996a, 1996b; Colinvaux and West, 1984; Guthrie, 1990). Regional pollen records from the height of the last glaciation show that grasses (Poaceae) and sedges (Cyperaceae) were dominant components of the vegetation that was almost entirely herbaceous (e.g., Anderson and Brubaker, 1994; Edwards et al., 2000). However, the taxonomic resolution for these taxa achievable via pollen analysis is less than that needed to characterize the paleovegetation. This deficiency has resulted in a range of conflicting vegetation reconstructions.

There are a number of sedimentary contexts (in addition to lake sediments) and paleoecological resources (in addition to pollen) found in Eastern Beringia that have considerable potential for enhancing paleoecological reconstructions. Plant macrofossils are somewhat rare, but previous studies demonstrate that they can provide detailed paleoenvironmental information where they have been found (Elias, 1992; Elias et al., 1996, 1997; Goetcheus and Birks, 2001; Zazula et al., 2003, 2005). Sites with *in situ* fossil vegetation and paleosols during the last glaciation are even more rare (Goetcheus and Birks, 2001; Froese et al., 2006; Zazula et al., 2006), though they provide direct information about Pleistocene plant community composition and structure. More recently, subfossil middens (e.g., nests and seed caches) within burrows of arctic ground squirrels (*Spermophilus parryi*) and other rodents from ice-rich silt deposits of interior Yukon Territory provide archives of local Pleistocene flora (Zazula et al., 2003, 2005).

Much of the graminoid macro-remains (leaves and stems) from fossil peat, buried vegetation, and arctic ground squirrel nests cannot be identified reliably based on macro-morphological characteristics. However, these samples could serve as ideal material to conduct stable carbon isotope analyses. The stable carbon isotope composition ( $\delta^{13}\text{C}$ ) of plants is influenced by a number of environmental conditions including soil moisture, salinity, humidity, and the stable isotope composition of atmospheric  $\text{CO}_2$  (O’Leary, 1988; Ehleringer et al., 1991; Lin

and Sternberg, 1992; Arens et al., 2000). The modern vegetation of Eastern Beringia is almost exclusively dominated by plants using the  $\text{C}_3$  photosynthetic pathway (e.g., Hultén, 1968; Welsh, 1974; Sage et al., 1999).  $\text{C}_4$  plants are rare in modern Eastern Beringia (e.g., Welsh, 1974; Sage et al., 1999) and have distinctive  $\delta^{13}\text{C}$  values between  $-10$  and  $-14\text{‰}$  (Ehleringer et al., 1991). By comparison, the total range of  $\delta^{13}\text{C}$  values exhibited by  $\text{C}_3$  plants is approximately  $-35\text{‰}$  (plants living in moist habitats) to  $-21\text{‰}$  (plants living in dry habitats), with an average value of  $-27\text{‰}$  (Ehleringer et al., 1991). Under drier conditions, plants using the  $\text{C}_3$  photosynthetic pathway tend to become more water-use-efficient (e.g., Ehleringer and Monson, 1993). In an attempt to reduce water losses through transpiration, these plants discriminate less against the heavier stable isotope of carbon ( $^{13}\text{C}$ ) in  $\text{CO}_2$ . Plant biomass subsequently becomes enriched in the heavier stable isotope of carbon, and the resulting  $\delta^{13}\text{C}$  derived from an analysis of the biomass is higher compared with  $\text{C}_3$  plant biomass from a wet environment (O’Leary, 1988). Biomass generated under different environmental conditions tends to retain a unique stable isotopic signature that is subsequently retained in subfossil material of a plant (e.g., Wooller et al., 2003a, 2004). In some instances these signatures can be generated from plants dating from millions of years in the past (Schweizer et al., 2006).

Based on this relationship, we began our research on the stable isotope composition of graminoids from Eastern Beringia by addressing several research questions. Do the stable isotope compositions of modern grasses and sedges reflect the environmental conditions (i.e., wet or dry) of the habitat in which the plants grew? Can we measure the  $\delta^{13}\text{C}$  values of subfossil graminoid material from sites in Eastern Beringia, and how do these values compare with the modern range of data for plants growing in the same region today? In our initial study, we examined the stable isotopic composition of a collection of modern and subfossil graminoids taken from several sites and sedimentary contexts in Eastern Beringia.

## Study Sites

Samples of modern grasses and sedges collected in Alaska and Yukon Territory were selected from the collection of plants archived in the Herbarium at the University of Alaska Museum of the North (ALA) in Fairbanks, Alaska. Exact locations of the samples, related to the sample codes in Table 1 and 2, are recorded in the database of specimens at the Herbarium (<http://arctos.database.museum>).

Subfossil graminoid material was taken from three separate locations with differing sedimentary contexts (Table 1). The first location is the Fox Permafrost Tunnel (<http://www.crrrel.usace.army.mil/permafrosttunnel/>) located roughly 10 km northeast of Fairbanks, off the Steese Highway (Fig. 1). The tunnel, excavated by the U.S. Army from 1963 to 1969, passes through loess deposits that span 10,000 to  $>40,000$   $^{14}\text{C}$  yr BP and contain the remains of subfossil plants (in many instances attached to paleosols—e.g., Figs. 2a and 2b) and animals (Hamilton et al., 1988). The second site is the Quartz Creek locality, a large north-facing placer-mining exposure in the Klondike goldfields of west-central Yukon Territory (Fig. 1). Three rodent nests (e.g., Fig. 2c) directly associated with the late Pleistocene Dawson tephra were excavated in the summer of 2002 from an exposure of ice-rich silt (Zazula et al., 2005). Two of the nests (GZ.02.07.01.31, GZ.02.07.01.33) are those of Pleistocene arctic ground squirrels, while the third (GZ.27.08 #8) is likely that of a microtine rodent. The age of the nests is similar to the age of the Dawson tephra, ca. 25,300  $^{14}\text{C}$

**TABLE 1**  
**The locations and descriptions of sites used in this investigation.**

Location	Sample code	Sample type	Age range ( $^{14}\text{C}$ yr BP)	References
Eastern Beringia	See Table 2 and 3	Leaves from herbarium specimens of modern grasses and sedges	Modern	Collections at the University of Alaska Museum of the North Herbarium
Fox Permafrost Tunnel	NA	Graminoid leaves	10,000 to >40,000	Hamilton et al. (1988)
Quartz Creek, Dawson	GZ02.07.01.31	Graminoid leaves from arctic ground squirrel nest	~25,300	Zazula et al. (2005)
	GZ02.07.01.33	Graminoid leaves from arctic ground squirrel nest	~25,300	Zazula et al. (2005)
	GZ27.08 #8	Graminoid leaves microtine from rodent nest	~25,300	Zazula et al. (2005)
Goldbottom Creek, Dawson		Graminoid leaves from paleo turf	~25,300	Zazula et al. (2005)

yr BP (Froese et al., 2002; Zazula et al., 2006). Nearly 12 m of ice-rich silt composed of both primary and colluviated loess overlies the Dawson tephra (for site details see Site 4 in Froese et al., 2002). The final paleoecological site is an *in situ* vegetated surface, or paleoturf (Fig. 2d), buried by Dawson tephra at Goldbottom Creek, approximately 30 km away from the Quartz Creek locality in the Klondike goldfields (Fig. 1). The buried fossil vegetation is a riparian plant community consisting of dwarf willow (*Salix* cf. *arctica*), abundant tufted hair grass (*Deschampsia caespitosa*), sedges (*Carex* spp.) with interspersed horsetail (*Equisetum* cf. *palustre*), mosses, and limited forbs or other grasses (Zazula et al., 2006; Froese et al., 2006).

## Methods

### MODERN AND SUBFOSSIL GRAMINOID ANALYSES

Leaf samples of modern sedges and grasses (~200 taxa) from Alaska and Yukon Territory were collected from specimens held at the University of Alaska Museum of the North Herbarium. Caution was taken to avoid any incorporation of material with herbarium glue attached. Four grass specimens from genera known to be  $\text{C}_4$  (*Muhlenbergia* and *Setaria*; Watson and Dallwitz, 1994) that were from sites within Eastern Beringia were also analyzed to test whether they were indeed  $\text{C}_4$ . Species were grouped (using information in Hultén, 1968, and habitat information taken from specimen herbarium labels) according to their habitat (wet or dry). An aliquot (~300 to 500  $\mu\text{g}$ ) of each freeze-dried sample of leaf fragments was weighed into a tin capsule, which was sealed and introduced into the autosampler (A2100) of a CE Instruments, NA 2500 series, Elemental Analyzer (EA) at the Alaska Stable Isotope Facility, University of Alaska Fairbanks (UAF). The stable carbon isotope ratio of the combustion gas ( $\text{CO}_2$ ) from each sample was analyzed using continuous-flow, stable isotope ratio mass spectrometry (IRMS, Finnigan MAT, Delta plus XP). The results are presented in standard delta ( $\delta$ ) notation.  $\text{CO}_2$  samples were analyzed relative to internal, working gas standards. Carbon isotope compositions ( $\delta^{13}\text{C}$ ) are expressed relative to Vienna Pee Dee Belemnite (V-PDB). Peptone was analyzed as a check on the analytical precision associated with the EA-IRMS. Analytical precision (SD) for  $\delta^{13}\text{C}$  was <0.1‰ (‰C = 1.8 and ‰N = 0.5). Precisions were typically lower than this for replicate ( $n = \geq 3$ ) analyses of subfossil and modern graminoid specimens.

*In situ* graminoid samples were removed from the loess composing the walls of the Fox Permafrost Tunnel in December 2003. Samples were taken in proximity to a series of radiocarbon-

dated points along the tunnel, which are the results of previous investigations and are marked in the tunnel (Hamilton et al., 1988). The radiocarbon years before present for each of five points in the tunnel were reported in Hamilton et al. (1988) and are  $11,300 \pm 160$ ,  $11,910 \pm 180$ ,  $14,280 \pm 230$ ,  $31,200 \pm 3000$ , and  $>40,000$   $^{14}\text{C}$  yr BP. Between four and seven separate plants were removed at each of the five locations. Loess was removed from the surface of the samples with a fine-haired brush. The samples were then freeze dried, weighed, and analyzed for their  $\delta^{13}\text{C}$  (as described above). A scanning electron microscope image of the surface of a graminoid leaf from the Fox Permafrost Tunnel was taken using an Electroscan E2020 environmental scanning electron microscope (ESEM) (Department of Geology and Geophysics, UAF) to examine the preservation state of the samples (e.g., Fig. 2e). Between 5 and 9 samples of separate graminoid remains were removed from each of the Quartz Creek middens and were freeze dried, weighed, and analyzed (as above) for their  $\delta^{13}\text{C}$ . Ten samples of separate graminoid remains were removed from the Goldbottom Creek paleoturf. The surfaces of these samples were brushed with a fine-haired brush to remove dirt. Subsamples of these samples were freeze dried, weighed, and analyzed for their  $\delta^{13}\text{C}$  (as described above).

### DATA ANALYSES

Each of our modern graminoid specimens was classified as representing a wet or dry environment based on herbarium sheet data and ecological information in Hultén (1968). Student's *t*-test was used to examine whether there were statistical differences between different groups of graminoid samples (e.g., plant specimens listed on their herbarium labels as having come from either a dry or wet habitat).

The  $\delta^{13}\text{C}$  of a plant is influenced by the  $\delta^{13}\text{C}$  of the surrounding atmospheric  $\text{CO}_2$  (e.g., Arens et al., 2000), and the  $\delta^{13}\text{C}$  of atmospheric  $\text{CO}_2$  has changed in the past (Friedli et al., 1986; Leuenberger et al., 1992). To be able to compare the  $\delta^{13}\text{C}$  of subfossil plant samples and modern plant samples we took into account the difference ( $\sim 1 \pm 0.2\text{‰}$ ) in  $\delta^{13}\text{C}$  of atmospheric  $\text{CO}_2$  of modern (Friedli et al., 1986) and paleo (late Quaternary) (Leuenberger et al., 1992) atmospheres. Given that the  $\delta^{13}\text{C}$  of past atmospheric  $\text{CO}_2$  for the date range of our subfossil samples was  $\sim 1\text{‰}$  higher than that of today's atmospheric  $\text{CO}_2$  (Friedli et al., 1986; Leuenberger et al., 1992), we added 1‰ to the  $\delta^{13}\text{C}$  of the modern plants (only when comparing modern and subfossil samples, i.e., Figs. 4 and 5). The  $\delta^{13}\text{C}$  values reported in the tables herein are unaltered  $\delta^{13}\text{C}$  values to allow their use in future studies.


TABLE 2

The stable carbon isotope composition and percent C of modern sedge (*Carex*) species from Alaska and the Yukon Territory.

Species	ALA accession	Ecology*	%C	$\delta^{13}\text{C}$
Dry sites				
<i>C. albo-nigra</i>	43466	Dry mountain sides	45.9	-25.6
<i>C. bigelowii</i>	10939	Dry places/solifluction soil/mountains	43.5	-26.9
<i>C. crawfordii</i>	V131128	Dry grasslands/roadsides	46.0	-28.7
<i>C. eburnea</i>	86872	Dry sand or rocky places/calcareous soils	43.5	-29.2
<i>C. filifolia</i>	43450	Dry ridges	45.0	-26.4
<i>C. foenea</i>	94961	Woods/river banks/sandy soils	43.1	-26.9
<i>C. glacialis</i>	V1302292	Dry calcareous soils	46.1	-25.3
Mean			44.7	-27.0
Standard deviation			1.3	1.5
Bogs and marshes				
<i>C. capitata</i>	V132883	Bogs/marshes	42.5	-29.4
<i>C. chordorrhiza</i>	44161	Bogs/lake margins/quagmires	45.9	-26.8
<i>C. diandra</i>	38521	Bogs/swamps/mires/borders of ponds	43.2	-28.1
<i>C. diandra</i>	V103152	Bogs/swamps/mires/borders of ponds	41.1	-26.8
<i>C. gynocrates</i>	60125	Wet meadows/marshes	44.6	-24.2
<i>C. heleonastes</i>	n.d.	Peat bogs/swamps	45.6	-28.4
<i>C. holostoma</i>	V134376	Marshes and mountain slopes	45.4	-27.2
<i>C. lasiocarpa</i>	V85249	Bogs/shallow water	46.0	-25.9
<i>C. leptalea</i>	V141806	Bogs/meadows and shores	46.6	-24.2
<i>C. limosa</i>	V143375	Bogs/quagmires/shallow water	47.7	-25.8
<i>C. loliacea</i>	V138710	Moist places/peat soil/muskeg	47.9	-26.2
Mean			45.1	-26.6
Standard deviation			2.1	1.6
Coastal/saline				
<i>C. glareosa</i>	V134815	Brackish marshes	45.3	-27.1
<i>C. gmelinii</i>	V144041	Sandy saline shores	45.9	-29.0
<i>C. lyngbyei</i>	V143833	Coastal salt marshes	46.6	-28.0
<i>C. mackenziei</i>	V143215	Wet places along seashores	42.3	-26.9
<i>C. maritima</i>	80996	shores/sandy soils	46.3	-28.1
<i>C. maritima</i>	90630	shores/sandy soils	42.8	-27.4
<i>C. macrocephala</i>	V04023	Sandy seashores	45.9	-27.4
Mean			45.2	-27.6
Standard deviation			1.7	0.8
Wet/moist places				
<i>C. adelostoma</i>	55014	Wet places	40.4	-28.3
<i>C. aquatilis</i>	V79576	Shallow water/marshes/along rivers	42.8	-28.3
<i>C. arcta</i>	V131115	Wet places	42.4	-28.1
<i>C. atherodes</i>	V117447	Wet meadows	36.1	-29.7
<i>C. atratiformis</i>	V13117	Moist places	42.5	-29.9
<i>C. aurea</i>	V131119	Meadows/moist places	37.3	-29.0
<i>C. bicolor</i>	V134906	Wet gravelly places in mountains	40.1	-30.6
<i>C. capillaris</i>	V145250	Moist (or dry) places	43.8	-32.3
<i>C. disperma</i>	V143683	Moist places	44.0	-30.3
<i>C. eleusinoides</i>	24985	Wet places/gravel bars	29.3	-27.6
<i>C. flava</i>	V94575	Moist places/calcareous soils	45.4	-29.1
<i>C. flava</i>	V94575	Moist places/calcareous soils	47.3	-29.2
<i>C. garberi</i>	V080696	Wet places	45.9	-29.1
<i>C. interior</i>	V141199	Wet meadows	46.6	-29.7
<i>C. kelloggii</i>	V117726	Wet places	41.9	-28.1
<i>C. krausei</i>	V139461	Moist places	38.6	-29.9
<i>C. lachenalii</i>	V143687	Alpine tundra/snow beds/above treeline	50.7	-28.9
<i>C. laeviculmis</i>	V135797	Wet meadows	46.2	-27.7
<i>C. lapponica</i>	V141611	Wet grassy places	46.7	-27.7
<i>C. laxa</i>	V141915	Wet places/mostly in woods	40.5	-27.1
<i>C. livida</i>	V143369	Wet places	44.6	-27.9
<i>C. macloviana</i>	96553	Meadows/gravelly shores	45.2	-28.2
<i>C. media</i>	V143394	Moist places in lowlands	44.9	-29.8
<i>C. membranacea</i>	V139569	Wet places	47.0	-27.8
<i>C. mertensii</i>	V143693	Wet rocky slopes	46.0	-29.3
Mean			43.0	-28.9
Standard deviation			4.5	1.2

\* Taken from the ecological descriptions given on the herbarium sheets.





**FIGURE 1.** The location of study sites used in this investigation. Fox tunnel provided subfossil graminoid specimens from paleosols in permafrost/loess. Goldbottom Creek and Quartz Creek provided subfossil graminoid remains preserved in a paleoturf and paleoarctic ground squirrel nests, respectively.

## Results

### MODERN GRASSES AND SEDGES

The total analyzed  $\delta^{13}\text{C}$  range for the sedges by habitat from Alaska and the Yukon is  $-32.3\text{‰}$  (listed as from wet habitats) to  $-24.2\text{‰}$  (listed as from bogs and marshes—permanently inundated with water) (Fig. 3, Table 2). The  $\delta^{13}\text{C}$  range for sedges from dry habitats was  $-29.2\text{‰}$  to  $-25.3\text{‰}$  with a mean of  $-27.0\text{‰}$  ( $\text{SD} = 1.5\text{‰}$ ), while the mean  $\delta^{13}\text{C}$  for sedges from bogs and marshes is  $-26.63\text{‰}$  ( $\text{SD} = 1.6\text{‰}$ ). Sedges from coastal and saline habitats have  $\delta^{13}\text{C}$  with a mean of  $-27.58\text{‰}$  ( $\text{SD} = 0.8\text{‰}$ ). The sedges from wet places have the most negative  $\delta^{13}\text{C}$  values with a mean of  $-28.91\text{‰}$  ( $\text{SD} = 1.2\text{‰}$ ). The  $\delta^{13}\text{C}$  values of sedges from wet habitats are significantly lower ( $p < 0.005$ ) than those of a dry habitat. However, there is no significant difference ( $p > 0.5$ ) in the mean  $\%C$  composition of the sedges from the different habitat types, and the mean  $\%C$  for sedges was  $\sim 45\%$ .

Table 3 presents the  $\delta^{13}\text{C}$ ,  $\%C$ ,  $\%N$ , and C:N data by habitat for modern grasses from Alaska and the Yukon. There is a significant difference ( $p < 0.001$ ) in  $\delta^{13}\text{C}$  by grass habitat (dry vs. wet) (Fig. 3). The grasses from wet habitats have a lower mean  $\delta^{13}\text{C}$  value ( $-29.1\text{‰}$ ,  $\text{SD} = 2.1$ ) compared with those from dry habitats ( $-26.4\text{‰}$ ,  $\text{SD} = 1.2\text{‰}$ ). No grasses from dry habitats have a  $\delta^{13}\text{C}$  value below approximately  $-29\text{‰}$ . Our preliminary analyses show that sedges and grasses from dry habitats have significantly less negative  $\delta^{13}\text{C}$  values compared with those from wet habitats, which is consistent with theory (O'Leary, 1988; Ehleringer et al., 1991). All four of the grasses suspected to be  $C_4$  have  $\delta^{13}\text{C}$  values within the range expected for  $C_4$  plants ( $-10$  to  $-14\text{‰}$ ) (Table 4).

### PLEISTOCENE GRASSES AND SEDGES

A number of features, including stomata, prickles, and long cells, showing a high degree of leaf preservation, are evident from the ESEM image (Fig. 2e) taken of a subfossil graminoid leaf

surface. In some instances, as in Figures 2a, 2b, and 2c, the graminoid remains are well preserved as whole plants attached to a paleosol. In the case of arctic ground squirrel nests, the collection of plant remains, including grasses, are tangled together. The  $\delta^{13}\text{C}$  and  $\%C$  values of the graminoid samples taken from the Fox Permafrost Tunnel are presented in Table 5. The  $\%C$  values of the samples range from 52 to 32 and are not significantly different compared with modern grasses and sedges ( $p > 0.2$ ). The  $\delta^{13}\text{C}$  values of these subfossil specimens range from  $-28.9$  to  $-24.3\text{‰}$  and vary significantly ( $p < 0.05$ ) among time points (Fig. 4), with the lowest mean  $\delta^{13}\text{C}$  value ( $-28\text{‰}$ ) seen from an area of the tunnel dated as ca.  $11,300 \pm 160$   $^{14}\text{C}$  yr BP and the highest mean  $\delta^{13}\text{C}$  value (approximately  $-25.5\text{‰}$ ) evident in a portion of the tunnel dated as between  $11,910 \pm 180$  and  $14,280 \pm 230$   $^{14}\text{C}$  yr BP.


The mean  $\%N$  of the samples was  $1.3 \pm 0.5$ , which is significantly smaller than that of the modern grasses (mean = 2.7,  $\text{SD} = 1.0$ ) ( $p < 0.001$ ) but within the range of values by species (Table 5). The  $\%C$  values (mean = 37,  $\text{SD} = 9$ ) are also significantly lower compared with the modern grasses ( $p < 0.001$ ). The Goldbottom Creek subfossils had the lowest  $\delta^{13}\text{C}$  values (mean =  $-27\text{‰}$ ) compared with the subfossil plants from Quartz Creek. The mean C:N values from Quartz Creek nests 1 and 2 (Table 4) were noticeably smaller (18 and 23) compared with the other Quartz Creek nest (46) and the Goldbottom Creek paleoturf (49).

## Discussion

### STABLE ISOTOPE AND ELEMENTAL COMPOSITIONS OF MODERN GRAMINOIDS

The  $\delta^{13}\text{C}$  values from analyses of grasses from Alaska and Yukon Territory (Fig. 3) illustrate that, as expected from theory (O'Leary, 1988), grasses from wet habitats had  $\delta^{13}\text{C}$  values that were more negative than those from dry habitats. Although there was considerable overlap among the data from these two habitats, the mean of samples from each habitat were found to be statistically different. Grasses from supposedly wet habitats were found to have  $\delta^{13}\text{C}$  values up to the maximum extent of the range shown by dry grasses. However, no grasses from dry habitats were found with  $\delta^{13}\text{C}$  values below approximately  $-29\text{‰}$ . The  $\delta^{13}\text{C}$  values of grasses from wet habitats ranged down to approximately  $-34\text{‰}$ . It seems, therefore, that  $\delta^{13}\text{C}$  values from an unknown grass would be better at typing the specimen as having come from a wet environment when  $\delta^{13}\text{C}$  are less than  $-29\text{‰}$ .

Whereas the  $\delta^{13}\text{C}$  patterns for wet and dry grasses seem relatively clear, this is not the case for modern sedges (Fig. 3). Like the grasses, sedges from moist habitats had significantly more negative  $\delta^{13}\text{C}$  values compared with those from dry habitats. However, sedges listed as from bogs or marshes had relatively high  $\delta^{13}\text{C}$  values, not significantly different from those for sedges in dry habitats ( $p > 0.1$ ). In fact, the highest  $\delta^{13}\text{C}$ , which is supposedly a marker of dry habitats, was found in a sedge from a bog/marsh (*Carex gynocrates*; Table 2). However, the photosynthetic physiology of plants can be influenced by additional environmental stressors (e.g., Lin and Sternberg, 1992; Dawson and Bliss, 1989). This is why we grouped sedges found in coastal or marine habitats to examine whether sedges potentially exposed to more saline conditions had relatively less negative  $\delta^{13}\text{C}$  values, a trend that has been demonstrated for other plants (Lin and Sternberg, 1992). In fact, the  $\delta^{13}\text{C}$  values of sedges found in coastal and saline conditions were significantly different compared with sedges from moist habitats ( $p < 0.005$ ). Salinity is unlikely to be the driving




**FIGURE 2.** (a and b) Plan and side view (respectively) of a paleosol removed from the Fox Permafrost Tunnel with well-preserved grasses and sedges still attached. (c) A subfossil arctic ground squirrel nest above the Dawson tephra (redrawn from Froese et al., 2002). (d) A coin (for scale) lying on the paleoturf at Goldbottom Creek. (e) An environmental scanning electron microscope image of a fossil graminoid cuticle (ca. 31,000  $^{14}\text{C}$  yr BP) from the Fox Permafrost Tunnel, Alaska.

factor of the relatively high  $\delta^{13}\text{C}$  values of sedges found in marshes and bogs (Fig. 3, Table 2). Although we cannot explain the relatively high  $\delta^{13}\text{C}$  values for sedges listed as from bogs and marshes, other researchers (Keeley and Sandquist, 1992) have reported similar results for some  $\text{C}_3$  plants growing in waterlogged conditions.

In terms of interpreting the paleoenvironmental record, the modern sedge data suggest a difficulty in separating dry sedges from marsh or bog sedges using their  $\delta^{13}\text{C}$  values. The answer might lie in additional lines of paleoenvironmental evidence, such as the sedimentary context of the sample, i.e., whether from a nest, paleosol, or paleoturf. This would assume that the paleograminoid

sample was local to the lithology. This might be a safer assumption when considering an *in situ* paleosol or paleoturf (e.g., Goetcheus and Birks, 2001; Froese et al., 2006; Zazula et al., 2006) compared with a loess sequence in which some material may be detrital. Moreover, when interpreting  $\delta^{13}\text{C}$  from subfossil remains, it is necessary to consider the role diagenesis might have played in altering the isotopic composition of a sample. Some processes associated with the decomposition of organic matter can slightly alter the stable isotopic composition of plant material (e.g., Macko et al., 1993; Fogel and Tuross, 1999; Wooller et al., 2003a), but this would generally be insufficient to shift the signature from one indicating high water-use efficiency (e.g.,





**FIGURE 3.** The stable carbon isotope composition of modern grasses and sedges from different habitats types in Alaska and Yukon Territory. The square symbols represent the mean of these groups with 1 standard deviation of the means shown. V-PDB = Vienna Pee Dee Belemnite.

$\delta^{13}\text{C} = -22\text{‰}$ ) to one of low water-use efficiency (e.g.,  $\delta^{13}\text{C} = -32\text{‰}$ ).

#### *C<sub>4</sub> GRASSES WITHIN THE EXTENT OF MODERN EASTERN BERINGIA*

Stable carbon isotopes provide a clear way of identifying grasses as either  $\text{C}_3$  or  $\text{C}_4$  (Tables 3 and 4). Four modern specimens of  $\text{C}_4$  grasses representing two species were identified from sites within Alaska and Yukon Territory (Table 4). This implies that  $\text{C}_4$  grasses are able to grow under modern subarctic environmental and atmospheric conditions (e.g., temperature, precipitation, and atmospheric  $\text{CO}_2$  concentrations). Some of these conditions are atypical of the wider “natural” environment. For instance, the areas where they grow were glaciated during the late Pleistocene (Duk-Rodkin, 1999). *Setaria viridis* is a Eurasian plant that has only been found as a nonpersistent casual introduction in Yukon, and *Muhlenbergia glomerata* is a plant sometimes associated with hot springs. Nevertheless, these  $\text{C}_4$  grasses were growing at relatively high latitudes for  $\text{C}_4$  plants (Sage et al., 1999).

The presence of these modern grasses at relatively high latitudes in the modern Eastern Beringia begs the question of whether  $\text{C}_4$  grasses could have been present during Pleistocene glacials. There is some literature to suggest that some  $\text{C}_4$  plant species existed in glacial refugia during the LGM (StreetPerrott et al., 1997; Boom et al., 2001; Wooller et al., 2001, 2003b). For instance, remains of the  $\text{C}_4$  grass genus *Bouteloua* based on phytolith morphology were reported from a Pleistocene fossil arctic ground squirrel nest in Alaska (Guthrie, 1982), though, unfortunately, this report was not confirmed with carbon isotope data.

Environmental change and variation (e.g., in atmospheric  $\text{CO}_2$  concentrations, salinity, disturbance or fire frequency, temperature, precipitation amounts, and seasonality) have been found to drive modern (e.g., Hattersley, 1992; Wooller et al., 2005) and past (e.g., Ehleringer et al., 1991; Ehleringer and Monson, 1993; StreetPerrott et al., 1997) proportions of  $\text{C}_4$  plants in vegetation. Are there any environmental factors of the LGM that

might have favored  $\text{C}_4$  plants in Eastern Beringia? Previous research indicates that some areas of Beringia, notably the interior, experienced relatively drier conditions in the past (Hopkins et al., 1982; Barnosky et al., 1987; Kutzbach, 1987; Anderson and Brubaker, 1994). Atmospheric  $\text{CO}_2$  concentrations were also lower during the LGM and could have favored  $\text{C}_4$  plants in some other locations (StreetPerrott et al., 1997). Although temperature, which is suggested to have been lower during the LGM (although there is limited information on seasonality and data are not quantitative or geographically widespread), favors  $\text{C}_3$  photosynthesis over  $\text{C}_4$  photosynthesis (Sage et al., 1999), there are some  $\text{C}_4$  plants that can survive at relatively lower temperatures (Ehleringer et al., 1991; Ehleringer and Monson, 1993). Our point here is not to suggest that  $\text{C}_4$  plants expanded during the past in Eastern Beringia, but rather, stating that conditions for  $\text{C}_4$  plant occurrence *may have* been present within the Beringian glacial refugium; and if they were, the presence of  $\text{C}_4$  plants in samples such as subfossil ground squirrel nests, loess, or paleoturfs would be obvious if the stable carbon isotopic composition of the graminoid material were analyzed.

#### *STABLE CARBON ISOTOPE MEASUREMENTS OF PALEOSAMPLES*

None of the  $\delta^{13}\text{C}$  values of subfossil graminoids taken from the Fox Permafrost Tunnel, Quartz Creek, or Goldbottom Creek sites were within the range of  $\text{C}_4$  grasses. The  $\delta^{13}\text{C}$  values were all within the range of  $\text{C}_3$  plants and within the total range of the modern graminoids analyzed in this study. There were significant differences between ages (Fig. 4) and between sites (Fig. 5), which implies that the past graminoids grew under different environmental conditions. Once the difference ( $\sim 1\text{‰}$ ) in the  $\delta^{13}\text{C}$  of  $\text{CO}_2$  in the modern and past atmospheres has been taken into account, the  $\delta^{13}\text{C}$  of modern and subfossil graminoids can be compared (Figs. 4 and 5). This approach is consistent with the procedure described by Arens et al. (2000) and allows  $\delta^{13}\text{C}$  values from plants of different ages to be compared. Having taken this into account, Figure 4 shows that the graminoid specimens from the oldest part of Fox Permafrost Tunnel likely grew in relatively mesic habitats. The ages of these samples suggest these graminoids inhabited wetter, pre-LGM environmental conditions that are characteristic of the Middle Wisconsinan interstadial (Anderson and Lozhkin, 2001). However, the means of the two sample sets from an area of the tunnel dated as between  $11,910 \pm 180$  and  $14,280 \pm 230$   $^{14}\text{C}$  yr BP lay within the range of the modern graminoids from dry habitats. In this respect the  $\delta^{13}\text{C}$  data imply that the graminoids grew in a relatively dry habitat that characterizes full glacial conditions during the LGM (Anderson and Brubaker, 1994; Kaplan et al., 2003). Analyses of the graminoids from the youngest portion of the tunnel ( $11,300 \pm 160$   $^{14}\text{C}$  yr BP) imply a marked change toward wetter conditions, where the  $\delta^{13}\text{C}$  of the subfossils are within the 1 SD of the mean of wet graminoids and outside the 1 SD of the mean of dry graminoids (Fig. 4). This marked shift toward wetter conditions at this time would be consistent with multiple lines of other evidence from Eastern Beringia that document the rise of the mesic-hydric taiga and tundra vegetation (Guthrie, 2006). Figure 4 does not include the  $\delta^{13}\text{C}$  range of coastal sedges, given the sedimentary context of the samples (paleosols in loess). Figure 4 also assumes that the subfossil plant remains were not derived from sedges that were once living in bogs and marshes, which were inundated with water. This is also a relatively safe assumption given the sedimentary context of the subfossils.



TABLE 3

The stable carbon isotope and elemental (C and N) composition of modern grasses (all C<sub>3</sub>) by habitat from Alaska and the Yukon Territory.

University of Alaska Museum of the North Herbarium accession						
Species	number	Habitat source*	%N	%C	δ <sup>13</sup> C	C:N
Wet sites						
<i>Agrostis alaskana</i>	V130401	2	2.6	47.1	−33.5	18.1
<i>Agrostis mertensii</i>	V104130	2	1.8	45.3	−31.1	24.6
<i>Agrostis clavata</i>	V133576	2	3.3	39.3	−32.9	12.1
<i>Agrostis exarata</i>	87242	3	3.8	47.3	−32.2	12.6
<i>Agrostis geminata</i>	V101618	1	1.3	39.6	−28.5	30.7
<i>Agrostis scabra</i>	V136251	1	1.6	43.5	−30.6	27.1
<i>Agrostis stolonifera</i>	87184	1	1.9	50.1	−32.1	25.8
<i>Agrostis trinii</i>	32710	2	2.6	48.9	−26.3	18.9
<i>Alopecurus aequalis</i>	V145495	1,2	3.4	40.2	−29.0	11.9
<i>Alopecurus alpinus</i>	V08050	2	5.0	52.3	−27.2	10.5
<i>Alopecurus alpinus</i>	V145923	2	4.8	49.9	−26.4	10.3
<i>Alopecurus geniculatus</i>	V90444	1,2	3.9	42.9	−30.0	11.0
<i>Alopecurus pratensis</i>	V084708	1	2.4	43.1	−27.7	17.6
<i>Arctagrostis latifolia</i>	V150486	1,2	4.2	50.2	−27.8	11.8
<i>Arctophila fulva</i>	V124235	1	1.7	44.4	−29.3	25.4
<i>Bromus ciliatus</i>	V98021	1	2.6	49.1	−28.7	18.6
<i>Bromus inermis</i>	V132817	3	1.9	44.0	−28.7	23.3
<i>Bromus pacificus</i>	87314	1	3.3	47.3	−30.5	14.5
<i>Bromus sitchensis</i>	V130430	1	2.4	44.7	−29.4	18.4
<i>Calamagrostis canadensis</i>	37341	1	3.6	45.8	−28.4	12.7
<i>Calamagrostis deschampsoides</i>	V150221	1,2	2.1	42.5	−28.6	20.1
<i>Calamagrostis inexpansa</i>	V98652	1,2	1.4	41.7	−27.9	29.6
<i>Calamagrostis lapponica</i>	V148854	1	1.8	40.0	−27.8	22.6
<i>Calamagrostis nutkaënsis</i>	V135176	2	2.4	51.0	−28.9	21.2
<i>Calamagrostis stricta</i>	V148979	3	2.6	44.9	−29.6	17.6
<i>Catabrosa aquatica</i>	V126356	1,2	2.8	42.7	−30.0	15.5
<i>Colpodium</i> sp.	V126356	1	2.3	49.9	−28.3	21.4
<i>Colpodium wrightii</i>	V134571	1	2.5	47.9	−28.4	19.4
<i>Deschampsia beringensis</i>	V144794	1,2	1.5	51.3	−27.2	33.1
<i>Deschampsia brevifolia</i>	VC73070	2	3.1	52.6	−28.0	17.2
<i>Elytrigia repens</i>	79377	1	2.5	45.2	−30.8	17.8
<i>Elymus glaucus</i>	V120708	1	2.8	50.6	−28.2	18.3
<i>Elymus glaucus</i>	V134801	1	3.4	47.2	−27.6	13.8
<i>Elymus hirsutus</i>	10446	1	3.0	45.4	−23.7	15.0
<i>Elymus macrourus</i>	V145983	1	4.8	53.0	−27.3	11.0
<i>Elymus subsecundus</i>	V134753	1	2.9	49.9	−33.3	17.4
<i>Elymus trachycaulus</i>	V134563	1	3.2	45.8	−29.4	14.2
<i>Festuca vahlmanum</i>	V150495	1	2.8	57.0	−28.4	20.5
<i>Glyceria borealis</i>	V147874	1	3.5	47.8	−28.7	13.6
<i>Glyceria maxima</i>	3901	1	3.2	44.5	−25.5	13.9
<i>Glyceria pulchella</i>	3929	1	3.4	45.0	−26.9	13.1
<i>Hierochloë odorata</i>	V144344	1,2	2.6	48.8	−27.5	18.5
<i>Hierochloë pauciflora</i>	V120083	1,2	3.7	49.8	−26.4	13.5
<i>Hordeum brachyantherum</i>	V134804	1,2	2.2	45.5	−31.0	20.7
<i>Hordeum caespitosum (jubatum)</i>	V102342	1	4.6	47.5	−33.4	10.4
<i>Koeleria gracilis</i>	V105345	1	2.6	48.4	−28.1	18.8
<i>Leymus mollis</i>	V109145	1	0.4	51.9	−25.7	126.5
<i>Leymus mollis</i>	V112482	1	1.6	45.4	−25.8	28.4
<i>Lolium multiflorum</i>	V120558	1	1.4	39.6	−30.9	27.4
<i>Lolium perenne</i>	V136111	3	3.5	41.5	−31.2	11.7
<i>Melica subulata</i>	V118033	1	5.5	50.9	−33.6	9.3
<i>Phalaris arundinacea</i>	V145470	1,2	3.1	50.4	−28.6	16.2
<i>Phippsia algida</i>	79368	2	2.1	41.9	−26.3	19.9
<i>Pleuropogon sabinei</i>	V90734	1,2	4.2	46.8	−29.4	11.1
<i>Poa annua</i>	V78642	1	2.1	47.7	−31.5	23.0
<i>Poa cusickii</i>	V133885	1	4.0	50.1	−27.5	12.4
<i>Poa eminens</i>	95349	1	1.4	38.1	−30.1	26.7
<i>Poa laxiflora</i>	V1491689	1	3.0	42.9	−29.8	14.3
<i>Poa macrocalyx</i>	V111930	1	2.6	42.3	−32.4	16.5
<i>Poa palustris</i>	V141408	1	2.0	43.5	−29.7	21.3
<i>Podagrostis aequalis</i>	V146059	2	1.8	48.0	−30.3	26.5

**TABLE 3**  
**Continued.**

Species	University of Alaska Museum of the North Herbarium accession		Habitat source*	%N	%C	$\delta^{13}\text{C}$	C:N
	number						
<i>Podagrostis thurberiana</i>	V146064		2	3.2	49.4	-29.7	15.5
<i>Puccinellia arctica</i>	V95031		1	1.6	51.2	-29.9	31.3
<i>Puccinellia deschampsoides</i>	V125157		1	4.2	52.5	-27.2	12.4
Mean				2.8	46.6	-29.1	19.9
Standard deviation				1.0	4.0	2.1	14.8
Dry sites							
<i>Avena hookeri</i>	V13072		2	2.2	45.2	-28.0	20.5
<i>Bromus carinatus</i>	V084995		1	1.4	44.9	-27.7	32.0
<i>Calamagrostis purpurascens</i>	V70363		1,2	3.1	50.9	-27.2	16.4
<i>Calamagrostis purpurascens</i>	V114008		1,2	2.5	45.8	-26.0	18.0
<i>Dactylis glomerata</i>	V112039		2	2.6	41.8	-28.6	16.1
<i>Danthonia intermedia</i>	V147827		1	1.4	46.8	-26.6	33.0
<i>Danthonia spicata</i>	V140898		1	1.9	47.6	-25.7	25.5
<i>Elymus alaskanum</i>	85269		1	1.4	45.6	-26.3	33.5
<i>Elymus spicata</i>	n.d.		1	3.2	52.3	-26.8	16.4
<i>Festuca baffinensis</i>	V143087		1	2.4	47.1	-24.6	20.0
<i>Festuca minutiflora</i>	80824		1	2.3	47.8	-25.7	20.9
<i>Festuca pratensis (Lolium pretense)</i>	10307		1	3.2	46.3	-28.8	14.4
<i>Festuca saximontana</i>	44197		1	3.7	44.3	-26.5	12.0
<i>Glyceria striata</i>	V70055		1	2.3	44.3	-25.8	19.4
<i>Hordeum vulgare</i>	V103674		1	0.4	34.1	-29.2	88.5
<i>Poa abbreviata</i>	44209		1	2.7	48.8	-26.2	18.3
<i>Poa alpina</i>	V144897		2	3.2	50.6	-28.7	15.7
<i>Poa arctosteporum</i>	V145360		1	1.0	48.8	-25.7	49.6
<i>Poa glauca</i>	280		2	2.3	49.9	-28.4	21.2
<i>Poa interior</i>	V136512		1	2.0	44.8	-28.1	22.8
<i>Poa pratensis</i>	289		1	3.3	44.8	-26.4	13.7
<i>Psathyrostachys juncea</i>	44311		3	3.2	47.9	-27.2	15.1
<i>Puccinellia pumila</i>	V116375		1	2.4	49.0	-26.0	20.0
<i>Schizachne purpurascens</i>	V137957		1	2.4	51.5	-27.3	21.2
<i>Secale cereale</i>	48653		1	2.1	51.1	-25.4	24.4
<i>Stipa comata</i>	V133110		2	3.8	51.5	-27.6	13.4
<i>Trisetum sibiricum</i>	V083666		1	3.3	51.2	-26.1	15.4
<i>Trisetum spicatum</i>	34532		1	2.6	49.1	-26.4	18.5
Mean				2.4	47.3	-26.9	23.4
Standard deviation				0.8	3.8	1.2	15.0

\* Taken from the ecological descriptions from (1) the herbarium sheets; (2) Hult n (1968); and (3) the Internet ([http://www.mun.ca/biology/delta/arctic/cf\\_ca/www/pocast.htm](http://www.mun.ca/biology/delta/arctic/cf_ca/www/pocast.htm), [http://www.hort.purdue.edu/newcrop/duke\\_energy/Bromus\\_inermis.html#Ecology](http://www.hort.purdue.edu/newcrop/duke_energy/Bromus_inermis.html#Ecology), [http://www.hort.purdue.edu/newcrop/duke\\_energy/Lolium\\_perenne.html#Ecology](http://www.hort.purdue.edu/newcrop/duke_energy/Lolium_perenne.html#Ecology), <http://www.fs.fed.us/database/feis/plants/graminoid/psajun/index.html>).

**TABLE 4**  
**The stable carbon isotope and elemental (C and N) composition of modern C<sub>4</sub> grasses from Eastern Beringia. n.d. = not determined.**

Species	University of Alaska Museum of the North Herbarium		Location	Habitat	%N	%C	$\delta^{13}\text{C}$	C:N
	accession number							
<i>Muhlenbergia glomerata</i>	20481		Liard River, Yukon Territory, 60�02'N, 128�35'W	n.d.	1.7	47.4	-12.0	28.5
<i>Muhlenbergia glomerata</i>	V128443		Beaver River Hot Springs, 60�22'N, 125�34'W	Growing on gravel bar	1.3	42.0	-13.2	32.9
<i>Muhlenbergia glomerata</i>	41089		Canada, NW Territories, 64�58'N, 123�40'W	Sandy cutbanks along river	1.9	40.5	-13.6	21.2
<i>Setaria viridis</i>	V128403*		Whitehorse, Yukon Territory	Growing in cracks in pavement	2.4	44.0	-14.0	18.0

\* Duplicates at University of Alaska Museum of the North Herbarium and B.A. Bennett Herbarium.

TABLE 5

The stable carbon isotope and elemental (C and N) composition of subfossil graminoid from Eastern Beringia. n.d. = not determined.

Age in <sup>14</sup> C years BP	%N	%C	δ <sup>13</sup> C	C:N
Fox Permafrost Tunnel				
11,300	n.d.	32.4	-27.4	n.d.
11,300	n.d.	35.5	-28.9	n.d.
11,300	n.d.	46.3	-27.6	n.d.
11,300	n.d.	46.1	-27.4	n.d.
11,300	n.d.	40.5	-28.9	n.d.
11,910	n.d.	41.1	-24.6	n.d.
11,910	n.d.	47.9	-26.8	n.d.
11,910	n.d.	54.8	-25.7	n.d.
11,910	n.d.	55.2	-25.6	n.d.
11,910	n.d.	31.7	-25.7	n.d.
14,280	n.d.	47.1	-25.0	n.d.
14,280	n.d.	51.2	-24.7	n.d.
14,280	n.d.	47.4	-28.0	n.d.
14,280	n.d.	43.9	-26.8	n.d.
14,280	n.d.	52.2	-24.3	n.d.
14,280	n.d.	48.2	-25.0	n.d.
14,280	n.d.	39.3	-24.7	n.d.
31,200	n.d.	52.2	-26.3	n.d.
31,200	n.d.	48.2	-27.2	n.d.
31,200	n.d.	39.3	-27.4	n.d.
31,200	n.d.	47.4	-28.2	n.d.
31,200	n.d.	46.4	-26.3	n.d.
≥40,000	n.d.	41.1	-26.7	n.d.
≥40,000	n.d.	48.8	-25.9	n.d.
≥40,000	n.d.	52.3	-25.7	n.d.
≥40,000	n.d.	50.3	-26.6	n.d.
Quartz Creek (nest GZ27.08 #8)				
~25,300	0.6	42.6	-25.4	74.6
~25,300	1.2	37.7	-25.5	30.6
~25,300	1.3	26.8	-27.3	20.7
~25,300	0.7	29.7	-24.7	45.0
~25,300	0.8	29.7	-25.1	35.8
~25,300	0.5	34.0	-25.7	69.4
Mean	0.9	33.4	-25.6	46.0
Standard deviation	0.3	5.9	0.9	21.7
Quartz Creek (nest GZ02.07.01.31)				
~25,300	2.4	45.1	-26.3	18.5
~25,300	1.9	30.8	-24.5	16.2
~25,300	1.8	37.2	-26.6	20.8
~25,300	1.6	28.0	-27.2	17.1
~25,300	2.1	32.7	-26.7	15.8
Mean	2.0	34.8	-26.3	17.7
Standard deviation	0.3	6.7	1.02	2.0
Quartz Creek (nest GZ02.07.01.33)				
~25,300	1.5	39.5	-26.8	26.3
~25,300	1.0	33.7	-24.6	34.6
~25,300	1.4	37.4	-24.8	27.1
~25,300	1.2	40.1	-25.9	33.0
~25,300	1.6	26.1	-27.3	16.2
~25,300	1.5	23.9	-27.3	16.5
~25,300	1.6	25.7	-27.3	15.9
~25,300	1.6	26.4	-27.2	16.9
~25,300	1.5	24.6	-27.5	16.1
Mean	1.4	30.8	-26.5	22.5
Standard deviation	0.2	6.8	1.2	7.8
Goldbottom Creek				
~25,300	0.9	48.8	-26.9	53.0
~25,300	0.9	47.2	-28.0	51.3
~25,300	0.9	34.4	-27.0	38.3
~25,300	1.0	45.9	-27.4	45.9
~25,300	0.7	47.3	-26.5	69.5
~25,300	1.0	48.1	-27.1	47.6
~25,300	1.1	47.7	-27.0	45.4


TABLE 5

Continued.

Age in <sup>14</sup> C years BP	%N	%C	δ <sup>13</sup> C	C:N
~25,300	1.2	46.4	-28.1	40.3
~25,300	0.8	47.0	-26.8	59.5
~25,300	1.1	43.7	-26.5	40.4
Mean	1.0	45.6	-27.1	49.1
Standard deviation	0.1	4.2	0.5	9.7

Unlike Fox Permafrost Tunnel, the sedimentary context of the graminoids from the arctic ground squirrel nests (Fig. 5) does not allow us to simply rule out the possibility that the graminoids originated from sedge-dominated bogs or marshes. The plants' remains in the nests may have been removed from a nearby habitat by arctic ground squirrels. In fact, the paleoecological reconstruction implies that the nest material likely came from a tundra/steppe upland plant community (Zazula et al., 2005, 2006). In this respect the paleoecological reconstruction is largely consistent with the δ<sup>13</sup>C analyses of the graminoids from Quartz Creek nest #8, where the mean δ<sup>13</sup>C value of the subfossil graminoids is within the 1 SD of the mean δ<sup>13</sup>C of modern grasses and sedges from dry habitats. There is some overlap in the 1 SD of the mean of the δ<sup>13</sup>C from subfossil graminoids from Quartz Creek nests 1 and 2 with the 1 SD of the δ<sup>13</sup>C of grasses and sedges from wet and dry habitats. Many paleoenvironmental data point to aridity in glacial Beringia, e.g., widespread sand dunes and sand sheets, loess, and sand wedges (Hopkins et al., 1982; Mann et al., 2002), and greatly lowered lake levels (Barber and Finney, 2000; Abbott et al., 2000; Edwards et al., 2000). GCM simulations of past conditions (e.g., Kaplan et al., 2003) also tend to indicate drier conditions but underestimate the degree of aridity compared with paleoecological data. In Kaplan et al. (2003), only the "driest" of four GCM simulations closely matched the proxy reconstructions for 21,000 cal. yr BP. Hultén (1937) hypothesized mesic environments for the Bering land bridge to explain the present distribution of Beringian plants. Mesic environments may have been more common on the central Bering land bridge than in continental areas of Eastern Beringia, as suggested by plant and insect macrofossils (Elias et al., 1996, 1997), mammalian faunal data (Guthrie, 2001), synthesis of regional pollen data (Anderson and Brubaker, 1994), and macrofossil data (Goetcheus and Birks, 2001). Furthermore, topography and landscape position greatly influence the distribution of arid and mesic habitats in local settings, such as constrained valleys with pronounced changes in relief over short distances. Evidence to support local-scale moisture variability is found in the diversity of vascular and bryophyte macrofossils within subfossil ground squirrel nests, with evidence for hydric, mesic, and xeric habitats all within a short distance (Zazula et al., 2005, 2006). In general, zonal habitats in the interior of Eastern Beringia were probably more arid, due perhaps to enhanced continentality and the influence of the nearby Laurentide ice sheet. A number of proxy- and model-based paleoclimatic reconstructions for glacial Eastern Beringia stress the role of cold, aridity, and clear conditions in the interior (Hopkins et al., 1982; Barnosky et al., 1987; Kutzbach, 1987; Anderson and Brubaker, 1994; Guthrie, 2001). Arid interior climates may have led to a topographically controlled vegetation mosaic with strong steppe-like elements. Modern examples of steppe habitats in Eastern Beringia are restricted to soils with insufficient moisture to support tree growth within the forest zone and are often associated with south-facing slopes with high net insolation and high evaporation (e.g., Edwards and Armbruster,






**FIGURE 4.** The stable carbon isotope composition of subfossil graminoid samples from the Fox Permafrost Tunnel along a timeline. Where a sample set has the same letter as another sample set, then this denotes no significant statistical difference between the data sets ( $p \geq 0.05$ ). Where sample sets have different letters, this denotes there is a significant statistical difference ( $p \leq 0.05$ ). The mean  $\delta^{13}\text{C}$  (and 1 standard deviation [std] shown) of the modern grasses from wet habitats (filled square) and dry habitats (square), and sedges from wet habitats (filled triangle) and dry habitats (triangle) have been corrected to account for the less negative  $\delta^{13}\text{C}$  of the atmosphere under which the subfossil plants grew. Given that the  $\delta^{13}\text{C}$  of past atmospheric  $\text{CO}_2$  for the date range of our subfossil samples was  $\sim 1\text{‰}$  higher than that of today's atmospheric  $\text{CO}_2$  (Friedli et al., 1986; Leuenberger et al., 1992), we added  $1\text{‰}$  to the  $\delta^{13}\text{C}$  of the modern plants.

1989; Lloyd et al., 1994; Vetter, 2000). Guthrie (2001) suggested that the effect of clear skies, exposed ground, and resultant increased absorption of solar radiation created warmer and better-drained soils than present. Plant remains and subfossil arctic ground squirrel nests from some sites from the Yukon (Zazula et al., 2003, 2005) also provide supporting evidence for this reconstruction.

The  $\delta^{13}\text{C}$  values of graminoids from the Goldbottom paleoturf yielded the lowest  $\delta^{13}\text{C}$  values, which were within the 1 SD of the mean of modern grasses and sedges from wet habitats. In this respect these data imply that the plants from the paleoturf were growing in a relatively wet habitat. This is consistent with the paleoecological reconstruction based on plant remains from the Goldbottom Creek paleoturf, which implies a riparian margin meadow (Froese et al., 2006; Zazula et al., 2006). Although the  $\delta^{13}\text{C}$  of plants from graminoids from the paleoturf imply a relatively wet habitat, the paleosol that these plant remains were attached to indicates that it was a fairly well-drained site and was not permanently inundated with water (Froese et al., 2006).

#### FUTURE DIRECTIONS: STABLE CARBON ISOTOPE MEASUREMENTS OF SUBFOSSIL GRAMINOID REMAINS AS A NOVEL PROXY WITHIN AN INTEGRATED MULTIPROXY APPROACH TO RECONSTRUCTING PAST BERINGIAN VEGETATION

Perhaps a better future classification of modern wet vs. dry grasses could be achieved if the classification of the habitat moisture status was based on quantitative (i.e., soil moisture content) rather than qualitative data (i.e., habitat descriptions on herbarium labels). This could subsequently be tested by taking specimens from a series of habitats where the percent soil moisture has been measured.



**FIGURE 5.** The stable carbon isotope composition of subfossil graminoid samples taken from paleoarctic ground squirrel nests from Goldbottom Creek and Quartz Creek sites in Yukon Territory. Where a sample set has the same letter as another sample set, then this denotes no significant statistical difference between the data sets ( $p \geq 0.05$ ). Where sample sets have different letters, this denotes there is a significant statistical difference ( $p \leq 0.05$ ). The mean  $\delta^{13}\text{C}$  (and 1 standard deviation shown) of the modern grasses from wet habitats (filled square) and dry habitats (open square) and sedges from wet habitats (filled triangle) and dry habitats (open triangle) have been corrected to account for the less negative  $\delta^{13}\text{C}$  of the atmosphere under which the subfossil plants grew. Given that the  $\delta^{13}\text{C}$  of past atmospheric  $\text{CO}_2$  for the date range of our subfossil samples was  $\sim 1\text{‰}$  higher than that of today's atmospheric  $\text{CO}_2$  (Friedli et al., 1986; Leuenberger et al., 1992), we added  $1\text{‰}$  to the  $\delta^{13}\text{C}$  of the modern plants.

Our current analysis of modern sedges was restricted to *Carex* specimens, and it would be interesting to analyze the  $\delta^{13}\text{C}$  values of other genera, e.g., *Eleocharis*, a taxon of which grows in Alaska (Hultén, 1968). Some species of this genus (e.g., *E. baldwinii*) are able to exist as either  $\text{C}_3$  or  $\text{C}_4$  according to habitat type (wet and dry, respectively; Uchino et al., 1995).

In the case of graminoids, the cuticle micromorphology of preserved leaves can potentially provide independent evidence to support or test a paleoecological reconstruction (Wooller et al., 2000, 2003b; Wooller, 2002; Wooller and Beuning, 2002). Figure 3 shows the well-preserved cuticular micromorphological features of a specimen dated at ca. 31,200  $^{14}\text{C}$  yr BP. The cuticle approach would be especially valuable where subfossil seeds and florets of graminoids are not preserved. The stable isotope data from a paleograminoid specimen could also be compared with a paleoecological reconstruction based on the ecological tolerances of macroscopically identifiable graminoid specimens recovered from the same sample.

We have presented an initial study investigating the application of  $\delta^{13}\text{C}$  analyses of subfossil graminoids in Eastern Beringian paleoecological research. Previous applications of  $\delta^{13}\text{C}$  analyses of subfossil plant material has largely been reserved to examine changes in the past proportions of  $\text{C}_3$  and  $\text{C}_4$  plants in the tropics and subtropics (e.g., Wooller et al. 2003b, 2005). Our data illustrate that  $\delta^{13}\text{C}$  of subfossil plant remains could also be used to reconstruct the environments of ecosystems dominated by  $\text{C}_3$  plants, such as the Arctic and subarctic.

## Conclusions

The  $\delta^{13}\text{C}$  of modern grasses species from Alaska and the Yukon Territory listed as from dry habitats on their herbarium labels is significantly higher than modern grasses from wet

habitats. This difference was also seen between modern sedges listed as from dry habitats compared with those from wet habitats. However, sedges listed as specifically from bogs or marshes were found to have a similar isotopic composition compared with dry sedges. To interpret  $\delta^{13}\text{C}$  from subfossil sedges it would be necessary to consider other lines of paleoenvironmental information, including the sedimentary context of a sample and plant macrofossil data.

Our analyses of subfossil graminoid remains from three separate sites (Fox Permafrost Tunnel, Quartz Creek, and Goldbottom Creek) show considerable variations in  $\delta^{13}\text{C}$ . Our investigation indicates that stable isotope analyses can be used to study past vegetation in Eastern Beringia during the late Quaternary. Information about plant growth conditions, particularly the occurrence of moisture stress, would aid in our understanding of arctic plant responses to climate change. Complementary ecological information can be derived from cuticle and macrofossil analyses of the same samples, and together these methods promise new insights into the paleoenvironments and paleoecology of Beringia.

### Acknowledgments

We thank Dr. Darren Grocke, Dr. Peter Sauer, and two anonymous reviewers for their constructive reviews and comments, which improved our manuscript. We thank the two diligent and enthusiastic high school students, Collin Stackhouse (Monroe School) and Robert Marcotte (West Valley School), for their hard work during the isotope survey of grasses and sedges from the University of Alaska Museum of the North Herbarium. We also thank Emily Molhoek for taking the ESEM images under the guidance of Ken Severin in the Department of Geology and Geophysics, UAF. We appreciated discussions with Tom Douglas from the U.S. Army Cold Regions Research and Engineering Laboratory. We also appreciated Ken Tape taking the time to facilitate our access to the Fox Permafrost Tunnel. We thank University of Alaska Museum of the North Herbarium for allowing us access to their herbarium specimens. We thank Tim Howe and Norma Haubenstock at the Alaska Stable Isotope Facility, UAF, for assisting with advising and supporting the high school students that worked on this project. Funding was provided by start-up funds awarded to M. Wooller by UAF. G. Zazula was funded by a Sir James Loughheed Scholarship (Alberta Provincial Government) and research grants from the Geological Society of America, Northern Scientific Training Program (DIAND, Canada), and a Natural Sciences and Engineering Research Council operating grant to Dr. R. Mathewes.

### References Cited

- Abbott, M. B., Finney, B. P., Edwards, M. E., and Kelts, K. R., 2000: Lake-level reconstructions and paleohydrology of Birch Lake, central Alaska, based on seismic reflection profiles and core transects. *Quaternary Research*, 53: 154–166.
- Anderson, P. M., and Brubaker, L. B., 1994: Vegetation history of northcentral Alaska: a mapped summary of late-Quaternary pollen data. *Quaternary Science Reviews*, 13: 71–92.
- Anderson, P. M., and Lozhkin, A. V., 2001: The stage 3 interstadial complex (Karginskii/middle Wisconsinan interval) of Beringia: variations in paleoenvironments and implications for paleoclimatic interpretations. *Quaternary Science Reviews*, 20: 93–126.
- Arens, N. C., Jahren, A. H., and Amundson, R., 2000: Can  $\text{C}_3$  plants faithfully record the carbon isotopic composition of atmospheric carbon dioxide? *Paleobiology*, 26: 137–164.
- Barber, V. A., and Finney, B. P., 2000: Late Quaternary paleoclimatic reconstructions for interior Alaska based on paleolake-level data and hydrologic models. *Journal of Paleolimnology*, 24: 29–41.
- Barnosky, A. G., Anderson, P. M., and Bartlein, P. J., 1987: The northwest US during deglaciation: vegetation history and paleoclimatic implications. In Ruddiman, W. F., and Wright, H. E., Jr. (eds.), *North America and adjacent oceans during the last deglaciation*. The Geology of North America series. Boulder: Geological Society of America, 289–321.
- Boom, A., Mora, G., Cleef, A., and Hooghiemstra, H., 2001: High altitude C-4 grasslands in the northern Andes: relicts from glacial conditions? *Review of Palaeobotany and Palynology*, 115: 147–160.
- Colinvaux, P. A., 1980a: Vegetation of the Bering land bridge revisited. *Quarterly Review of Archaeology*, 5: 10–16.
- Colinvaux, P. A., 1980b: Historical ecology of Beringia: the south land bridge coast at St. Paul Island. *Quaternary Research*, 16: 18–36.
- Colinvaux, P. A., 1996a: Low-down on a land bridge. *Nature*, 382: 21–22.
- Colinvaux, P. A., 1996b: Reconstructing the environment. In West, F. H. (ed.), *American beginnings, the prehistory and palaeoecology of Beringia*. Chicago: University of Chicago Press, 13–19.
- Colinvaux, P. H., and West, F. H., 1984: The Beringian ecosystem. *Quarterly Review of Archaeology*, 5: 10–16.
- Cwynar, L. C., and Ritchie, J. C., 1980: Arctic steppe-tundra: a Yukon perspective. *Science*, 208: 1375–1377.
- Dawson, T. E., and Bliss, L. C., 1989: Intraspecific variation in the water relations of *Salix arctica*, an arctic-alpine dwarf willow. *Oecologia*, 79(3): 322–331.
- Duk-Rodkin, A., 1999: *Glacial limits map of Yukon Territory*. Geological Survey of Canada, Open File 3694, Indian and Northern Affairs Canada Geoscience, Map 1999-2, scale 1:1,000,000.
- Edwards, M. E., and Armbruster, W. S., 1989: A tundra-steppe transition on Kathul Mountain, Alaska. *Arctic and Alpine Research*, 21: 521–554.
- Edwards, M. E., Bigelow, N. H., Finney, B. P., and Eisner, W. R., 2000: Records of aquatic pollen and sediment properties as indicators of late-Quaternary Alaskan lake levels. *Journal of Paleolimnology*, 24: 55–68.
- Ehleringer, J. R., and Monson, R. K., 1993: Evolutionary and ecological aspects of photosynthetic pathway variation. *Annual Review of Ecology and Systematics*, 24: 411–439.
- Ehleringer, J. R., Sage, R. F., Flanagan, L. B., and Pearcy, R. W., 1991: Climate change and the evolution of  $\text{C}_4$  photosynthesis. *Trends in Ecology & Evolution*, 6: 95–99.
- Elias, S. A., 1992: Late Quaternary beetle faunas of southwestern Alaska: evidence of a refugium for mesic hygrophilous species. *Arctic and Alpine Research*, 24: 133–144.
- Elias, S. A., Short, S. K., and Birks, H. H., 1996: Life and times of the Bering land bridge. *Nature*, 382: 60–63.
- Elias, S. A., Short, S. K., and Birks, H. H., 1997: Late Wisconsinan environments of the Bering land bridge. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 136: 293–308.
- Fogel, M. L., and Tuross, N., 1999: Transformation of plant biochemical to geological macromolecules during early diagenesis. *Oecologia*, 120: 336–346.
- Friedli, H., Lotscher, H., Oeschger, H., Siegenthaler, U., and Stauffer, B., 1986: Ice core record of the  $^{13}\text{C}/^{12}\text{C}$  ratio of atmospheric  $\text{CO}_2$  in the past two centuries. *Nature*, 324: 237–238.
- Froese, D. G., Westgate, J. A., Preece, S., and Storer, J. E., 2002: Age and significance of the late Pleistocene Dawson tephra in eastern Beringia. *Quaternary Science Reviews*, 21: 2137–2142.
- Froese, D. G., Zazula, G. D., and Reyes, A. V., 2006: Seasonality of the late Pleistocene Dawson tephra and exceptional preservation of a buried riparian surface in central Yukon Territory, Canada. *Quaternary Science Reviews*, 25: 1542–1551.

- Goetcheus, V. G., and Birks, H. H., 2001: Full-glacial upland tundra vegetation preserved under tephra in the Beringia National Park, Seward Peninsula, Alaska. *Quaternary Science Reviews*, 20: 135–147.
- Guthrie, R. D., 1968: Palaeoecology of the large-mammal community in interior Alaska during the late Pleistocene. *American Midland Naturalist*, 70: 346–363.
- Guthrie, R. D., 1982: Mammals of the mammoth steppe as paleoenvironmental indicators. In Hopkins, D. M., Matthews, J. V. Jr., Schweger, C. E., and Young, S. B. (eds.), *The paleoecology of Beringia*. New York: Academic Press, 307–326.
- Guthrie, R. D., 1990: *Frozen fauna of the mammoth steppe: the story of Blue Babe*. Chicago: University of Chicago Press, 323 pp.
- Guthrie, R. D., 2001: Origin and causes of the mammoth steppe: a story of cloud cover, woolly mammal tooth pits, buckles, and inside-out Beringia. *Quaternary Science Reviews*, 20: 549–574.
- Guthrie, R. D., 2006: New carbon dates link climatic change with human colonization and Pleistocene extinctions. *Nature*, 441: 207–209.
- Hamilton, T. D., Craig, J. L., and Sellmann, P. V., 1988: The Fox permafrost tunnel: a late Quaternary geologic record in central Alaska. *Geological Society of America Bulletin*, 100: 948–969.
- Harington, C. R., 2003: *Annotated bibliography of Quaternary vertebrates of northern North America—With radiocarbon dates*. Toronto: University of Toronto Press, 539 pp.
- Harington, C. R., 2005: The eastern limit of Beringia: mammoth remains from Banks and Melville Islands, Northwest Territories. *Arctic*, 58(4): 361–369.
- Hattersley, P. W., 1992: C<sub>4</sub> photosynthetic pathway variation in grasses (Poaceae): its significance for arid and semi-arid lands. In Chapman, G. P. (ed.), *Desertified grasslands their biology and management*. London: Academic Press, 181–212.
- Hopkins, D. M., Matthews, J. V. J., Schweger, C. E., and Young, S. B. (eds.), 1982: *Paleoecology of Beringia*. New York: Academic Press, 489 pp.
- Hultén, E., 1937: *Outline of the history of Arctic and Boreal biota during the Quaternary Period*. Stockholm: Bokforlags Akiebolaget Thule, 168 pp.
- Hultén, E., 1968: *Flora of Alaska and neighbouring territories: a manual of the vascular plants*. Stanford: Stanford University Press, 1008 pp.
- Kageyama, M., Peyron, O., Pinot, S., Tarasov, P., Guiot, J., Joussaume, S., and Ramstein, G., 2001: The Last Glacial Maximum climate over Europe and western Siberia: a PMIP comparison between models and data. *Climate Dynamics*, 17: 23–43.
- Kaplan, J. O., Bigelow, N. H., Prentice, I. C., Harrison, S. P., Bartlein, P. J., Christensen, T. R., Cramer, W., Matveyeva, N. V., McGuire, A. D., Murray, D. F., Razzhivin, V. Y., Smith, B., Walker, D. A., Anderson, P. M., Andreev, A. A., Brubaker, L. B., Edwards, M. E., and Lozhkin, A. V., 2003: Climate change and Arctic ecosystems: 2. Modeling, paleodata-model comparisons, and future projections. *Journal of Geophysical Research—Atmospheres*, 108: ALT12-1-17.
- Keeley, J. E., and Sandquist, D. R., 1992: Carbon—Fresh-water plants. *Plant Cell and Environment*, 15: 1021–1035.
- Kozhevnikov, J. P., and Ukraintseva, V. V., 1999: Pleistocene tundra-steppe: arguments pro and con. In Haynes, G., Klimowicz, J., and Reumer, J. W. F. (eds.), *Mammoths and the mammoth fauna: studies of an extinct ecosystem*. St. Petersburg, Russia: Proceedings of the First International Mammoth Conference, 1995: 199–210.
- Kutzbach, J. E., 1987: Model simulations of the climatic patterns during the deglaciation of North America. In Ruddiman, W. F., and Wright, H. E. Jr. (eds.), *North America and adjacent oceans during the last deglaciation*. The Geology of North America series. Boulder: Geological Society of America, 289–321.
- Leuenberger, M., Siegenthaler, U., and Langway, C. C., 1992: Carbon isotope composition of atmospheric CO<sub>2</sub> during the last ice age from an Antarctic ice core. *Nature*, 357: 488–490.
- Lin, G., and Sternberg, L. D. L., 1992: Effect of growth form, salinity, nutrient and sulfide on photosynthesis, carbon isotope discrimination and growth of red mangrove (*Rhizophora Mangle* L.). *Australian Journal of Plant Physiology*, 19: 509–517.
- Lloyd, A. H., Armbruster, W. S., and Edwards, M. E., 1994: Ecology of a steppe-tundra gradient in interior Alaska. *Journal of Vegetation Science*, 5: 897–912.
- Macko, S. A., Engel, M. H., and Parker, P. L., 1993: Early diagenesis of organic matter in sediments: assessment of mechanisms and preservation by the use of isotopic molecular approaches. In Engel, M. H., and Macko, S. A. (eds.), *Organic geochemistry*. New York: Plenum Press, 211–224.
- Mann, D. H., Peteet, D. M., Reanier, R. E., and Kunz, M. L., 2002: Responses of an arctic landscape to late glacial and early Holocene climatic changes: the importance of moisture. *Quaternary Science Reviews*, 21: 997–1021.
- O'Leary, M. H., 1988: Carbon isotopes in photosynthesis: fractionation techniques may reveal new aspects of carbon dynamics in plants. *Bioscience*, 38: 328–336.
- Ritchie, J. C., 1984: *Past and present vegetation of the far northwest of Canada*. Toronto: University of Toronto Press.
- Sage, R. F., Wedin, D. A., and Meirong, L., 1999: The biogeography of C<sub>4</sub> photosynthesis: patterns and controlling factors. In Sage, R. F., and Monson, R. K. (eds.), *C<sub>4</sub> plant biology*. San Diego: Academic Press, 596 pp.
- Schweizer, M. K., Wooller, M. J., Toporski, J., Fogel, M. L., and Steele, A., 2006: Examination of an Oligocene lacustrine ecosystem using C and N stable isotopes. *Palaeogeography Palaeoclimatology Palaeoecology*, 230: 335–351.
- Shapiro, B., and Cooper, A., 2003: Beringia as an Ice Age genetic museum. *Quaternary Research*, 60: 94–100.
- Shapiro, B., Drummond, A. J., Rambaut, A., Wilson, M. C., Matheus, P. E., Sher, A. V., Pybus, O. G., Gilbert, M. T. P., Barnes, I., Binladen, J., Willerslev, E., Hansen, A. J., Baryshnikov, G. F., Burns, J. A., Davydov, S., Driver, J. C., Froese, D. G., Harington, C. R., Keddie, G., Kosintsev, P., Kunz, M. L., Martin, L. D., Stephenson, R. O., Storer, J., Tedford, R., Zimov, S., and Cooper, A., 2004: Rise and fall of the Beringian steppe bison. *Science*, 306: 1561–1565.
- StreetPerrott, F. A., Huang, Y. S., Perrott, R. A., Eglinton, G., Barker, P., BenKhelifa, L., Harkness, D. D., and Olago, D. O., 1997: Impact of lower atmospheric carbon dioxide on tropical mountain ecosystems. *Science*, 278: 1422–1426.
- Uchino, A., Samejima, M., Ishii, R., and Ueno, O., 1995: Photosynthetic carbon metabolism in an amphibious sedge, *Eleocharis baldwinii* (Torr.) Chapman: modified expression of C<sub>4</sub> characteristics under submerged aquatic conditions. *Plant Cell Physiology*, 36: 229–238.
- Vetter, M. A., 2000: Grasslands of the Aishihik-Sekulmun Lakes area, Yukon Territory, Canada. *Arctic and Alpine Research*, 53: 165–173.
- Viereck, L. A., Dyrness, C. T., Batten, A. R., and Wenzlick, K. J., 1992: *The Alaska vegetation classification*. Portland: U.S. Department of Agriculture Forest Service.
- Watson, L., and Dallwitz, M. J., 1994: *Grass genera of the world*. Cambridge: CAB International.
- Welsh, S. L., 1974: *Anderson's flora of Alaska and adjacent parts of Canada*. Provo: Brigham Young University Press.
- Willerslev, E., Hansen, A. J., Binladen, J., Brand, T. B., Gilbert, M. T. P., Shapiro, B., Bunce, M., Wiuf, C., Gilichinsky, D. A., and Cooper, A., 2003: Diverse plant and animal genetic records from Holocene and Pleistocene sediments. *Science*, 300(5620): 791–795.
- Wooller, M. J., 2002: Fossil grass cuticles from lacustrine sediments: a review of methods applicable to the analysis of tropical African lake cores. *Holocene*, 12: 97–105.



- Wooller, M. J., and Beuning, K. R., 2002: Introduction to the reconstruction and modeling of grass-dominated ecosystems—Preface. *Palaeogeography Palaeoclimatology Palaeoecology*, 177: 1–3.
- Wooller, M. J., Street-Perrott, F. A., and Agnew, A. D. Q., 2000: Late Quaternary fires and grassland palaeoecology of Mount Kenya, East Africa: evidence from charred grass cuticles in lake sediments. *Palaeogeography Palaeoclimatology Palaeoecology*, 164: 207–230.
- Wooller, M. J., Agnew, A. D. Q., Mathai, S., Swain, D. L., and Street-Perrott, F. A., 2001: An altitudinal and stable carbon isotope survey of grasses on Mount Kenya, East Africa. *Journal of East African Natural History*, 90: 68–85.
- Wooller, M., Smallwood, B., Scharler, U., Jacobson, M., and Fogel, M., 2003a: A taphonomic study of  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values in *Rhizophora mangle* leaves for a multi-proxy approach to mangrove palaeoecology. *Organic Geochemistry*, 34: 1259–1275.
- Wooller, M. J., Swain, D. L., Ficken, K. J., Agnew, A. D. Q., Street-Perrott, F. A., and Eglinton, G., 2003b: Late Quaternary vegetation changes around Lake Rutundu, Mount Kenya, East Africa: evidence from grass cuticles, pollen and stable carbon isotopes. *Journal of Quaternary Science*, 18: 3–15.
- Wooller, M. J., Smallwood, B., Behling, H., and Fogel, M., 2004: Mangrove ecosystem dynamics and elemental cycling at Twin Cays, Belize during the Holocene. *Journal of Quaternary Science*, 19: 1–9.
- Wooller, M., Johnson, B., Wilkie, A., and Fogel, M., 2005: Stable isotope characteristics across narrow savanna/woodland ecotones in Wolfe Creek Meteorite Crater, Western Australia. *Oecologia*, 145: 100–112.
- Yurtsev, B. A., 1982: Relicts of the xerophytic vegetation of Beringia in northeast Asia. In Hopkins, D. M., Matthews, J. V. Jr., Schweger, C. E., and Young, S. B. (eds.), *Paleoecology of Beringia*. New York: Academic Press, 157–177.
- Zazula, G. D., Froese, D. G., Schweger, C. E., Mathewes, R. W., Beaudoin, A. B., Telka, A. M., Harington, C. R., and Westgate, J. A., 2003: Ice Age steppe vegetation in east Beringia. *Nature*, 423: 603.
- Zazula, G. D., Froese, D. G., Westgate, J. A., La Farge, C. A., and Mathewes, R. W., 2005: Paleoeecology of Beringian “packrat” middens from central Yukon Territory. *Quaternary Research*, 63: 189–198.
- Zazula, G. D., Froese, D. G., Elias, S. A., Kuzmina, S., La Farge, C., Reyes, A. V., Sanborn, P. T., Schweger, C. E., Smith, C. A. S., and Mathewes, R. W., 2006: Vegetation buried under Dawson tephra (25,300  $^{14}\text{C}$  yr BP) and locally diverse late Pleistocene paleoenvironments of Goldbottom Creek, Yukon, Canada. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 242: 253–286.

*Ms accepted July 2006*