

Vegetation Development on Deglaciaded Rock Outcrops from Glaciar Frías, Argentina

Authors: Garibotti, Irene A., Pissolito, Clara I., and Villalba, Ricardo

Source: Arctic, Antarctic, and Alpine Research, 43(1) : 35-45

Published By: Institute of Arctic and Alpine Research (INSTAAR),
University of Colorado

URL: <https://doi.org/10.1657/1938-4246-43.1.35>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Vegetation Development on Deglaciaded Rock Outcrops from Glaciar Frías, Argentina

Irene A. Garibotti*†

Clara I. Pissolito* and

Ricardo Villalba*

*Instituto Argentino de Nivología,
Glaciología y Ciencias Ambientales
(IANIGLA), CCT-CONICET
Mendoza, Av. Ruiz Leal s/n, C.C. 330,
Mendoza 5500, Argentina.

†Corresponding author:
ireneg@lab.cricyt.edu.ar

Abstract

The retreat of glaciers during past decades has led to the emergence of large rock outcrops in many glaciaded areas around the world. Primary succession of vegetation in glacier forelands has been described for many regions, but most studies have been conducted on glacial deposits, whereas deglaciaded rock outcrops have received little attention. This study assesses the pattern of primary succession on a chronosequence of five rock outcrops exposed during the past 140 years by the retreat of Glaciar Frías in the Patagonian Andes, Argentina. Data on floristic composition and species cover for algae, lichens, ferns, bryophytes, and vascular plants were recorded on sampling plots. Ordination and classification analyses discriminate three major successional stages, each dominated by a different species assemblage, suggesting directional replacement of species in the succession. The pioneer stage is dominated by the crustose lichen *Placopsis perrugosa*, the mid-successional stage by a lichen-moss mat dominated by the moss *Racomitrium lanuginosum*, and the late-successional stage by a large diversity of vascular plants. The low density of *Nothofagus dombeyi* saplings in the late-successional site indicates that plant succession is still in progress 140 years after deglaciation. Progress in succession appears to be influenced by species life-cycle traits and facilitative interactions among species. The comparison of the successional processes between rock outcrops and unconsolidated glacial deposits suggests that the vegetation sequence is similar, but the rate of succession is slower on rock outcrops. The development of a ground lichen-moss cover, previous to the widespread colonization by vascular plants, accounts for the slower succession progress on rock outcrops. The establishment of *Nothofagus* stands takes at least 100 yrs longer on the rock outcrops than on glacial deposits. Under predicted climate warming, most Patagonian Andes glaciers will continue the retreat along steep bedrock slopes, where similar, long-term vegetation successional patterns to those observed on Glaciar Frías foreland will eventually occur.

DOI: 10.1657/1938-4246-43.1.35

Introduction

Climatic warming in mountain ecosystems influences the dynamics of the vegetation through changes in ecophysiological processes and in disturbance regimes (Körner, 2005). In addition, climatic warming affects glaciaded-mountain ecosystems by determining glacier retreat that exposes extended areas of bare terrain to biological colonization (Matthews and Whittaker, 1987; Chapin et al., 1994). Studies of ecosystem development on recently deglaciaded terrains in Europe and North America provide comprehensive information on successional vegetation changes and the mechanisms driving succession (Matthews, 1992). The relative importance of vegetation traits, biological interactions, and environmental forces driving the processes of species colonization and replacement are well known (Svoboda and Henry, 1987; Walker and Chapin, 1987; Chapin et al., 1994). In addition, it has been shown that landscape characteristics, stochastic processes, and disturbance events have a large influence on vegetation successional patterns (del Moral et al., 1995; Matthews, 1999).

Large rock outcrops have emerged during past decades from beneath glaciers in many glaciaded areas, such as Patagonia, the

Alps, and North America (Rivera and Casassa, 2004; Paul et al., 2007; Pelto, 2009). Most studies of primary succession on deglaciaded terrains have been performed on unconsolidated glacial sediments, whereas there is little empirical data on vegetation development on bedrock outcrops (Matthews, 1992). Rock outcrops are stressful environments subjected to high thermal contrasts, drought due to low water holding capacity, substrate instability due to intense water runoff, and limiting soil formation (Shure and Ragsdale, 1977; Sarthou et al., 2009). Our knowledge about the primary succession on rock outcrops is mainly based on studies conducted within forested landscapes. In general, the vegetation establishment on outcrops is initiated by the early development of a lichen cover and an organic matter layer, and the succession progress is frequently associated with the deepening of the soil layer (Burbanck and Phillips, 1983; Uno and Collins, 1987; Asselin et al., 2006). The development of a mature forest community on rock outcrops is slow and periods of ca. 1000 years have been reported (Asselin et al., 2006). A similar slow vegetation development occurs on lava flows, where the pioneer cryptogam colonizers are replaced by higher plants ca. 600 years after the lava emplacement (Cutler et al., 2008). Hence,




FIGURE 1. Location map of Glaciar Frías foreland in the north Patagonian Andes showing the position of the frontal moraines (M) and the sampling sites (S). Estimated dates of moraines and sampling sites exposition are given in Table 1.

successional processes involving interactions between different vegetation groups are expected to occur on long-term temporal scales on deglaciated rock outcrops. Additional analyses on deglaciated rock outcrops are required to properly understand the landscape changes related to glacier retreat along mountain bedrock slopes, and to predict short- and long-term effects of changing climate on glaciated-mountain ecosystems.

The retreat of glaciers is a conspicuous sign of climate changes during the last century in the Patagonian Andes (Luckman and Villalba, 2001; Masiokas et al., 2010), yet investigations of primary succession on recently deglaciated terrains have been relatively rare in this area. Some studies have analyzed the vegetation on glacier moraines (Lawrence and Lawrence, 1959; Heusser, 1960, 1964; Pisano, 1978; Rabassa et al., 1981; Veblen et al., 1989; Dollenz, 1991; Armesto et al., 1992) and the influence of some specific biological interactions on plant colonization (Henríquez, 2004; Henríquez and Lusk, 2005). These studies have mostly focused on *Nothofagus* establishment and have rarely considered other components of the vegetation, such as algae, lichens, and bryophytes. Moreover, the study of the primary succession in general has received little attention within *Nothofagus* forests in the southern hemisphere (Orwin, 1972; Archer et al., 1973; Ashton and Moore, 1978; Wardle, 1980; Sommerville et al., 1982).

The aim of this study is to document and assist understanding of the primary succession process on rock outcrops following glacier retreat. Glaciar Frías, in the north Patagonian Andes, offers an excellent opportunity for the study of vegetation succession on rocky environments. Five rock outcrops have been exposed by glacier recession from its Neoglacial maximum and provide a chronosequence of sites covering the last 140 years (Fig. 1). In this paper we describe the pattern of primary succession of vegetation on the Glaciar Frías outcrops and infer the possible mechanisms driving vegetation development in these stressful environments. The study integrates all vegetation groups (algae, lichens, ferns, mosses, and vascular plants) allowing a thorough analysis of the community changes over time. In addition, successional pattern on rock outcrops at Glaciar Frías were compared with vegetation establishment on unconsolidated glacial deposits reported in the literature for the Patagonian Andes. By conducting this analysis we explore possible future vegetation landscape changes in mountain regions as glaciers continue retreating along bedrock slopes.

Methods

STUDY AREA

Glaciar Frías is the northernmost ice body of Mount Tronador (41°10'S, 71°50'W), one of the highest mountains (3554 m) in the northern Patagonian Andes (Fig. 1). The climate of the zone is temperate and wet. Available climatic data from the Mascardi weather station, 20 km east from Glaciar Frías, indicate a mean annual temperature of 7.6 °C (January mean 12.9 °C, July mean 2.4 °C). Total annual precipitation in the Frías valley is ca. 4300 mm (Barros et al., 1983; Villalba et al., 1990). The vegetation in the Frías valley corresponds to the Valdivian temperate rain forest, which is a multistratified forest dominated by the evergreen *Nothofagus dombeyi* and the conifer *Fitzroya cupressoides* (Ezcurra and Brion, 2005).

The chronology of Glaciar Frías recession is well known from the study of historical drawings, written records, terrestrial and aerial photographs, direct measurements of ice front, and the dendrochronological dating of moraines. The maximum Neoglacial advance of Glaciar Frías was reached ca. AD 1660 (Rabassa et al., 1978; Villalba et al., 1990), and the glacier has subsequently retreated more than 1500 m along the Frías valley (Fig. 1). A sharp forest trim-line defines the boundary between the glacier foreland and the mature forest not affected by the last major Neoglacial advance. A well-preserved sequence of frontal moraines remains in the bottom of the valley as evidence of seven minor readvances of Glaciar Frías since the last Neoglacial maximum (Villalba et al., 1990; Masiokas, 2008). The most recent readvance occurred during the years 1976–1977 (Rabassa et al., 1979).

At the Glaciar Frías foreland the bottom and slopes of the valley show contrasting environments. Along the bottom of the Frías valley wet meadows cover the glaciofluvial deposits and vascular plants grow on the frontal moraines. In contrast, along the valley slopes, bedrock outcrops without glacial sediments on top are mostly covered by lichens and bryophytes with sparsely distributed vascular plants. Inclination of the valley slopes is around 30°.

The study sites are located on the southern valley side, where five granodiorite rock outcrops have been successively exposed by the retreat of Glaciar Frías (Fig. 1). Exposure dates for the rock outcrops were estimated from the dates of the moraines (Villalba




FIGURE 2. Selected historical photographs and drawings of Glaciar Frías used to constrain the exposure dates of the rock outcrops in the study sites. Historical sources: (a) Fonck (1896), (b) Steffen (1909), (c) De Agostini (1945), (d) anonymous (IANIGLA archive), (e and f) R. Villalba.

et al., 1990; Masiokas, 2008), direct measurements of ice front positions (Rabassa et al., 1978), and the analysis of old terrestrial and aerial photographs (Fig. 2). Study sites were located between points of known exposure times, so the mean of these dates was used as an estimation of the sampling site age. The five rock outcrops studied are a chronosequence covering the past 140 years (Table 1). As the glacier forefield represents a spatial chronosequence, the analysis of the vegetation growing on sites exposed at different ages was used to infer the process of primary succession (Pickett, 1989; Foster and Tilman, 2000; Walker et al., 2010).

SAMPLING DESIGN

Rock outcrops are characterized by high compositional heterogeneity in vegetation due to the presence of a large variety of microhabitats modulated by topography (Wiser et al., 1996; Matthes-Sears and Larson, 2006; Opazo Medina et al., 2006). The rock outcrops exhibit meso- and micro-scale topographic variability due to the presence of drainage channels between outcrops and small cracks on the rock surface, respectively. Inspection of the area evidences striking differences in vegetation between rock surfaces and drainage channels. However, drainage channels constitute a minor fraction of the Glaciar Frías rock outcrop landscape. In consequence, our study focused in the analysis of the vegetation growing on the rock surfaces, the major landscape features on the deglaciated valley slopes.

We randomly located 10 sampling plots in each study site. Plots of 5×10 m were used for analyzing ferns and vascular

plants, and subplots of 1×1 m for analyzing the cryptogamic flora (algae, lichens, ferns, and bryophytes). Within each plot and subplot the species were recorded and their coverage visually estimated. Cover estimates for lichens and bryophytes were recorded to the species level whenever possible or by genera or morphological groups when field identification was not feasible. In addition, a floristic inspection throughout the study area was conducted to detect the presence of species not occurring in the plots. Lichens and mosses not identified in the field were collected for later taxonomical determination in the laboratory. Voucher specimens are deposited at the Argentinean Institute of Snow, Ice and Environmental Sciences (IANIGLA). Nomenclature for vascular plants follows Zuloaga and Morrone (1999a, 1999b), and for lichens and bryophytes Brummitt and Powell (1992).

Altitude, aspect, slope, surface topography, canopy coverage, percentage of the substrate covered by pebbles, and percentage of bare rock outcrop and bare soil were recorded at each plot. Surface topography was categorized as elevated, depressed, and flat with respect to the surrounding area. Inclination of the sampling plots depends on microtopography, and the slope at each sampling plot was estimated using clinometers. Percent canopy coverage was estimated using a spherical densitometer.

DATA ANALYSES

Patterns of variation in plant diversity along the successional sequence were assessed using indices of species diversity and plots of the species-abundance distribution (Lambshead et al., 1983;

TABLE 1

Exposure dates for sampling sites estimated from historical information and dendrochronological dating of frontal moraines. Historical information includes: TP (terrestrial photographs), AP (aerial photographs), D (drawings), and FP (direct measurement of glacier front positions). The historical drawing by Hess in 1856 and the terrestrial photos are shown in Figure 2. Dates of moraines from Villalba et al. (1990).

Sampling site	Estimate date of site deglaciation	Estimated site age (years)	Dates of moraines limiting site date	Evidences of site exposition dates from historical information
S1	1984	24	none	The AP from 1981 and the TP from 1983 (Anonymous) show Glacier Frías covering S1 at these dates. The TP from 1986 (IANIGLA) shows S1 uncovered by ice.
S2	1960	48	AD 1942 (M7) to 1977 (M8)	The TP taken between 1935 and 1938 by De Agostini (1945) and the AP from 1944 show S2 covered by ice. The FP by Rabassa et al. (1978) shows that S2 was uncovered by ice in 1977.
S3	1924	84	AD 1914 (M6) to 1942 (M7)	The TP taken between 1935 and 1938 by De Agostini (1945) shows S3 uncovered by ice.
S4	1903	109	AD 1881 (M5) to 1914 (M6)	The TP from 1893 (Steffen, 1909) shows a >50-m-tall ice front. This photograph relocation suggests that S4 was covered by ice in 1893.
S5	1868	140	AD 1839 (M4) to 1881 (M5)	The etching (D) by Hess (Fonck, 1896) shows the whole study area covered by Glacier Frías in 1856.

Magurran, 1988). Three diversity indices which provide complementary information on community structure were selected: the Shannon index (H') combines species richness and evenness; the reciprocal of Simpson's index ($1/D$) measures species dominance depending on the proportional abundance of all species; and the Berger-Parker index gives a value of dominance of the most abundant species, dividing the coverage of the dominant species by the total coverage of the species.

The vegetation data were analyzed by Detrended Correspondence Analysis (DCA), which ordines sampling plots according to their floristic composition and species coverage, allowing assessment of overall patterns in vegetation changes (Jongman et al., 1995). The ordination matrix contained 71 species in 50 sampling plots. Detrending was performed by segments, and rare species were not down-weighted. Vegetation data were also analyzed using a Two-Way Indicator Species Analysis (TWINSPAN) in order to determine vegetation groups (VG) characteristic of the different successional stages (Leps and Šmilauer, 1999). This analysis works with qualitative data, so quantitative data of species coverage was transformed to qualitative variables called pseudospecies, which are defined by cut-levels of species coverage (Jongman et al., 1995). In our analysis pseudospecies cut levels were set at 0, 2, 5, 20 and 50%, representing the whole range of species coverage. The minimum group size for division was 7, and a maximum of 4 levels of division was used.

The influence of different variables on vegetation changes were indirectly assessed correlating the DCA ordination axes with the dates of site exposure and environmental variables. In addition, a Canonical Correspondence Analysis (CCA) was applied to directly assess the main patterns of variation in the vegetation community accounted for by the explanatory variables (Jongman et al., 1995). The variables included were site exposure date, slope, surface topography, canopy coverage, and percentage of bare rock outcrop, bare soil, and pebbles. A Monte Carlo permutation test was used to test the significance of the first ordination axis (Leps and Šmilauer, 1999). In addition, the statistical significance of the partial effect of each explanatory variable (variability explained by a given variable after accounting for the effects of the other variables under analysis) was estimated by a Monte Carlo permutation test as the respective variable was step-wise added to the model.

Results

GENERAL PATTERNS IN SPECIES RICHNESS, COVER, AND DIVERSITY

A total of 97 species were identified in the floristic surveys performed on the rock outcrops studied at Glacier Frías foreland. These include species from 10 different life forms, trees; shrubs; herbs; graminoid herbs; ferns; mosses; foliose, fruticose, and crustose lichens; and algae (the full list of species is available from the corresponding author upon request). Fourteen additional species (9 graminoid herbs, 2 mosses, and 3 crustose lichens) were not identified due to the absence of reproductive structures at the time of sampling.

A general pattern of increasing species numbers and total plant cover occurred along the spatial chronosequence (Table 2). From the youngest to the oldest site (sites 1 to 5) the number of species increased from 31 to 44 and total plant cover increased from 44 to 154%, respectively. In the earliest exposed sites, total plant cover exceeded 100% due to the development of a multistratified community. Crustose lichens dominate the recently exposed sites (sites 1 to 3), but decline in the oldest sites 4 and 5. Cover of vascular plants, mosses, and foliose and fruticose lichens gradually increase with age. Ferns and algae are poorly represented at the study outcrops.

The species-abundance distributions at each study site are shown in Figure 3. The curves for all five sample sites indicate the presence of one or two dominant species (>10% cover), a variable number of species with intermediate abundance (between 1 and 10% cover), and a large number of rare species (<1% cover). Dominance increases over time, as indicated by the steep initial portions of the species rank-abundance curves, but there is also a general trend of increasing diversity through the successional sequence, with higher species richness and more even distribution of abundance among the species of intermediate abundance (Fig. 3). Interpretation of the species diversity indices is problematic because the rank-abundance curves from sites 2, 3, and 4 intersect each other, indicating that these communities are not comparable in terms of intrinsic diversity (Lambhead et al., 1983). Comparison of the species diversity indices for the oldest site 5 and the earlier site 1 (their rank-abundance curves do not intersect; Fig. 3) agree with the interpretation of the rank-

TABLE 2

Community structure in the five study sites along the Glaciar Frías forefield: mean \pm standard deviation cover of each vegetation group and diversity indices. Estimated exposure dates of sites are given in Table 1; site 1 is the youngest and site 5 the oldest.

	Site 1	Site 2	Site 3	Site 4	Site 5
Species number	31	33	37	33	44
Total vegetation cover (%)	44.6 \pm 21.0	81.1 \pm 34.7	116.8 \pm 33.9	82.0 \pm 33.2	153.9 \pm 46.9
Vascular plant cover (%)	8.7 \pm 4.9	15.7 \pm 9.6	17.5 \pm 16.1	10.4 \pm 8.2	47.2 \pm 46.7
Fern cover (%)	0.1 \pm 0.3	0.1 \pm 0.3	0.5 \pm 1.6	0.1 \pm 0.3	0
Moss cover (%)	11.8 \pm 8.0	23.5 \pm 18.3	41.3 \pm 24.0	45.4 \pm 29.0	72.7 \pm 22.5
Crustose lichen cover (%)	17.1 \pm 22.8	27.4 \pm 14.0	34.0 \pm 19.1	12.6 \pm 8.9	5.3 \pm 4.9
Foliose and fruticose lichen cover (%)	6.7 \pm 8.8	14.4 \pm 10.0	23.5 \pm 16.2	13.5 \pm 9.1	28.4 \pm 15.5
Algae cover (%)	0.2 \pm 0.4	0	0	0	0.3 \pm 0.9
Shannon index (H')	2.34	2.48	2.45	2.21	2.45
Simpson index (1/D)	1.18	1.14	1.15	1.23	1.23
Berger-Parker index (d)	0.33	0.24	0.28	0.39	0.41

abundance curves suggesting a trend of increasing species diversity in the vegetation community and of dominance of the most abundant species as succession progresses (Table 2).

VEGETATION ASSEMBLAGES

The species classification by TWINSPLAN differentiated six major vegetation assemblages. Based on the time of entering in the successional sequence and the period in which a species achieves the maximum coverage, the vegetation assemblages were differentiated as corresponding to the pioneer, mid-, or late-successional stage (Table 3). The pioneer species are those that colonized the recently exposed sites but disappeared in older sites, the mid-successional species prevailed in the middle-aged sites, whereas the late-successional species appear late in the succession.

The pioneer species dominate the community for about 50 years (sites 1 and 2) and decrease in coverage later in the succession (Table 3). The dominant species in the pioneer-successional stage are the crustose lichen *Placopsis perrugosa*, forming a dense cover on the rock outcrops, and the moss *Andreaea* sp., growing in small cracks on the rock surfaces. Besides these, the lichens *Stereocaulon speciosum* and *Placopsis stenophylla*, the moss *Racomitrium lanuginosum*, and the small shrubs *Senecio argyreus*, *Baccharis racemosa*, and *Gaultheria pumila* are also present with relatively high coverage in the early stage (Table 3). A large diversity of vascular plants (classified in the VG1) colonizes early the fine material accumulated between rock outcrops (Table 3). Most of these vascular species are only sporadically present in the younger and not in the older sites, thus we assumed that they are present by chance and not actually part of the successional sequence on the rock outcrops. Indeed, ordination analysis of the composition and coverage data shows that these species are clearly separated in the right side of the ordination graph, outside the center of the diagram where the sample sites lie (Fig. 4), indicating a minor influence of VG1 plants on the sites ordination.

The mid-successional stage is dominated by the moss *R. lanuginosum*. This species became prominent about 80 years after site exposure (site 3), and in combination with the fruticose lichens *Stereocaulon* spp., *Cladonia lepidophora*, and *C. subchordalis* forms a dense lichen-moss carpet on the rock surfaces (Table 3). After about 140 years the lichen-moss mat covered more than 90% of the rock surface on site 5. Other characteristic species of the mid-successional stage were the shrubby vascular plants *Gaultheria pumila* and *G. caespitosa*. In addition, species typical of the pioneer and late-successional stages, such as *Senecio argyreus* and

Escallonia alpina, respectively, were present with relatively high coverage in middle-aged sites (Table 3).

The late-successional stage is characterized by the invasion of a large diversity of vascular plant species, which on average contribute 47.2% of the total vegetation cover (Tables 2 and 3). Local variability in vegetation development is high, as indicated by the large standard deviation of the mean cover values for the vegetation groups (Table 2). The most relevant vascular plants, in order of their coverage are *Empetrum rubrum*, *Berberis buxifolia*, *Quinchamalium chilense*, *Discaria nana*, *Escallonia alpina*, *Baccharis racemosa*, and some graminoid herbs.

VEGETATION PATTERNS AND EXPLANATORY VARIABLES

Patterns in vegetation composition and species coverage along the spatial chronosequence were explored with DCA and CCA. Both analyses showed the same general pattern, thus only CCA ordination diagrams are shown (Fig. 4). Samples from each study site form groups relatively distinct in the ordination space, although the high within-site variability determines zones of overlap between groups (Fig. 4a). Sites and vegetation assemblages arranged successively along the first ordination axis (Fig. 4), indicating a trend of progressive vegetation change along the chronosequence.

The eigenvalues for the DCA (0.482 and 0.152 for DCA 1 and DCA 2, respectively) were similar to those recorded for the CCA (0.364 and 0.162 for CCA 1 and CCA 2, respectively), suggesting




FIGURE 3. Species-abundance distribution plots for each study site. Site 1 is the youngest and site 5 the oldest.

TABLE 3

Mean percent cover (%) of the species in the five study sites at Glaciar Frías foreland. Cover values higher than 1.5% are indicated in bold. Vegetation groups (VG) as defined by TWINSpan. Estimated exposure date of sites is given in Table 1.

VG	Species	Abbreviated name	Species life form	Site 1	Site 2	Site 3	Site 4	Site 5
Pioneer species								
1	<i>Cerastium arvense</i>	Carve	Herb	0.1				
	<i>Draba gilliesii</i>	Dgill	Herb	0.2				
	<i>Fuchsia magellanica</i>	Fmage	Shrub		0.1			
	<i>Gamochoaeta nivalis</i>	Gniva	Herb	0.1				
	<i>Luzula racemosa</i>	Lrace	Herb	0.1				
	<i>Leucheria papillosa</i>	Lpapi	Herb	0.3				
	<i>Rhytidosperra picta</i>	Rpict	Graminae	0.9	0.9	1.9	0.3	
	<i>Ribes magellanicum</i>	Rmage	Shrub		0.5			
	<i>Senecio fistulosus</i>	Sfist	Herb	0.3				
	Graminoid herb 2	Gram	Graminae	0.3				
	<i>Placopsis cribellans</i>	Perib	Lichen		0.5			
2	<i>Baccharis racemosa</i>	Brace	Shrub	1.2	1.2	0.1		1.8
	Graminoid herb 1	Gram	Graminae	0.7	0.6	0.4	0.1	0.1
	<i>Placopsis stenophylla</i>	Psten	Lichen	1.7	2.7		2.4	0.1
	<i>Stereocaulon speciosum</i>	Sspec	Lichen	3.1	0.7		0.1	0.1
3	<i>Gunnera tinctoria</i>	Gtinc	Herb	0.6				
	<i>Senecio argyreus</i>	Sargy	Herb	1.1	3.2	3.9	1.5	
	Graminoid herb 3	Gram	Graminae	0.5	0.2	0.2	0.2	0.3
	<i>Andreaea</i> sp.	Andre	Moss	9.5	17.5	9.1	13.7	8.5
	<i>Placopsis perrugosa</i>	Pperru	Lichen	14.8	19.8	22.0	5.1	2.5
	<i>Pseudophebe pubescens</i>	Poube	Lichen		0.2			0.1
	<i>Trentepohlia</i> sp.	Trent	Lichen	0.2				0.3
Mid-successional species								
4	<i>Placopsis bicolor</i>	Pbico	Lichen		2.8	6.4	1.0	
	<i>Rhizocarpon geminatum</i>	Rgemi	Lichen		0.6	2.3		0.4
	<i>Calceolaria biflora</i>	Cbifl	Herb	0.1		0.1		
	<i>Rhizocarpon superficiale</i>	Rsupe	Lichen	0.2				
	<i>Acaena pinnatifida</i>	Apinn	Herb		0.8	0.5		
	<i>Lathyrus magellanicus</i>	Lmage	Herb			0.2		
	Graminoid herb 4	Gram	Graminae		0.1		0.1	
	<i>Acarospora sinoptica</i>	Asino	Lichen			0.4		
	<i>Peltigera rufescens</i>	Prufe	Lichen		0.1			
	<i>Xanthoparmelia cordillerana</i>	Xcord	Lichen			0.5	0.1	
5	<i>Gaultheria pumila</i>	Gpumi	Shrub	1.3	4.0	2.9	0.2	6.5
	<i>Rumohra adiantiformis</i>	Radia	Fern	0.1	0.1	0.5	0.1	
	Graminoid herb 5	Gram	Graminae	0.2	0.1	0.4	0.2	0.6
	Graminoid herb 6	Gram	Graminae	0.6	1.1	1.4	0.7	2.3
	Unidentified crustose lichen	Lich	Lichen	0.1	0.1	0.3	0.1	0.1
	<i>Cladonia lepidophora</i>	Clepi	Lichen		5.4	9.5	2.6	8.7
	<i>Rhizocarpon geographicum</i>	Rgeog	Lichen		0.7	2.6	0.8	1.0
	<i>Stereocaulon</i> spp.	Ster	Lichen	2.3	8.0	11.2	8.9	13.5
	<i>Stereocaulon botryosum</i>	Sbotr	Lichen	1.3		2.0	1.1	
	<i>Racomitrium lanuginosum</i>	Rlanu	Moss	2.3	6.0	32.2	31.7	63.2
	<i>Gaultheria caespitosa</i>	Gcaes	Shrub		1.1	1.2	1.9	3.4
	<i>Muhlenbergia</i> sp.	Muhl	Graminae				0.4	
	<i>Silene chilensis</i>	Schil	Herb			0.3		0.7
	<i>Lecidea auriculata</i>	Lauri	Lichen	0.2				0.2
	<i>Rinodina</i> sp.	Rino	Lichen	0.1	0.2		3.3	1.1
Late-successional species								
6	<i>Berberis buxifolia</i>	Bbuxi	Shrub		0.8	1.3	0.1	5.4
	<i>Chloraea alpina</i>	Calpi	Herb		0.2	0.1	0.1	0.8
	<i>Fragaria chiloensis</i>	Fchil	Herb			0.6	0.3	0.2
	Graminoid herb 7	Gram	Graminae	0.1		0.1	0.2	1.1
	<i>Cladonia subchordalis</i>	Csubc	Lichen			0.3	0.7	5.2
	<i>Neofuscelia plana</i>	Nplan	Lichen				0.1	0.1
	<i>Quinchamalium chilense</i>	Qchil	Herb		0.7	0.7	1.1	4.3
	<i>Escallonia alpina</i>	Ealpi	Shrub		0.1	0.2	2.0	3.0
	<i>Nothofagus dombeyi</i>	Ndomb	Tree			0.5		1.0
	<i>Pseudopanax laetevirens</i>	Plaet	Shrub					1.0
	<i>Buellia</i> sp.	Buel	Lichen					0.1

TABLE 3
Continued.

VG	Species	Abbreviated		Site 1	Site 2	Site 3	Site 4	Site 5
		name	Species life form					
	<i>Baccharis nivalis</i>	Bniva	Herb					0.5
	<i>Discaria nana</i>	Dnana	Shrub				1.2	3.2
	<i>Empetrum rubrum</i>	Erubr	Shrub			0.2		7.5
	<i>Sisyrinchium arenarium</i>	Saren	Herb					0.2
	Graminoid herb 8	Gram	Graminae					0.2
	Graminoid herb 9	Gram	Graminae					0.5
	<i>Trisetum</i> sp.	Tris	Graminae			0.2		2.5
	Unidentified moss	Moss	Moss					1.0
	<i>Cladonia pocillum</i>	Cpoci	Lichen					0.1
	<i>Pseudocyphellaria encoensis</i>	Penco	Lichen					0.7
	<i>Carex</i> sp.	Carex	Graminae			0.1		0.1

that the explanatory variables included in the canonical analysis are adequate for explaining the variation in species composition and cover. In addition, the species-environmental correlations were relatively high for both analyses (Table 4), revealing a strong relationship between changes in the vegetation and the explanatory variables available. The first DCA and CCA axes are significantly and strongly correlated to site exposure dates (Fig. 4a, Table 4), indicating that the dominant pattern in community structure is the successional change associated with increasing time since site deglaciation. The second ordination axes are significantly correlated to surface topography, showing positive values for depressed surfaces and negative values for elevated surfaces (Fig. 4a, Table 4). Thus, microtopographic heterogeneity, such as elevated, depressed, and flat surface topography, can partially explain the within-site variability in vegetation structure. The canonical axes are also significantly correlated with percentage of bare rock surface, percentage of bare soil, and canopy coverage, reflecting the progressive occupation of the bare terrain by the vegetation as succession progresses.

Discussion

Primary succession on the bedrock outcrops of Glaciar Frías foreland follows a model of directional replacement of species (*sensu* Svoboda and Henry, 1987). An initial stage dominated by a crustose lichen is followed by a mid-successional stage characterized by a lichen-moss mat, whereas vascular plants diversified and increased in coverage during the late-successional stage (Fig. 4, Table 3).

The pioneer crustose lichen *Placopsis perrugosa* is a successful colonizer, with high growth rate and dispersal ability, and is frequently found dominating recently deglaciated terrains in Chile, New Zealand, and Antarctica (Orwin, 1970; Lindsay, 1978; Galloway, 1992). This lichen formed pure stands on the study rock outcrops during the first 50 years after deglaciation (Table 3). With increasing terrain age, *P. perrugosa* centers disintegrate probably due to limitations in nutrient transport from the periphery to the center of the thallus (Nash, 1996). The declining of *P. perrugosa* on the older study surfaces seems to be exclusively related to the species life-cycle, as no different lichens or mosses overgrow or develop in close contact to *P. perrugosa* competing for space.

Lichens are considered to be initiators of succession on rock surfaces because they significantly enhance rock weathering, derive inorganic nutrients from the rocks, and provide organic materials (Adamo and Violante, 2000). In addition, the presence of external cephalodia with cyanobacteria capable of fixing

nitrogen in *P. perrugosa* significantly increases the nitrogen stock in the new substrates, where nitrogen is a limiting nutrient for plant colonization (Vitousek, 1994). However, we noted that rock surfaces exposed by *P. perrugosa* as the lichen centers degraded show little evidence of disaggregation and fragmentation, suggesting that the colonization of this lichen does not contribute much to rock weathering. In addition, *P. perrugosa* disappears from the rock surfaces before the widespread colonization by mid-successional species, indicating that there is not a direct interaction between this lichen and later colonizers. Therefore, the early colonization by *P. perrugosa* seems to contribute little to the establishment of later colonizers on rock outcrops in the study area. Our observations are consistent with studies questioning the role of pioneer crustose lichens in the primary succession process. For example, Longton (1992) and Kurina and Vitousek (2001) considered that crustose lichens are able to colonize bare surfaces early because of their ruderal life-cycle traits, but do not have relevant positive effects on later colonizers.

The development of a lichen-moss mat between 50 and 80 years after deglaciation marks the transition from the pioneer to the mid-successional stage. The species that dominated the mat (*Racomitrium lanuginosum*, *Cladonia* spp., and *Stereocaulon* spp.) are common pioneer colonizers during primary succession on glacier forelands, lava flows, and high mountain environments (Veblen and Ashton, 1979; Veblen et al., 1989; Vetaas, 1994; Hodkinson et al., 2003; Cutler et al., 2008). They are highly tolerant to stressful environmental conditions and have the ability to rapidly spread over intact rock surfaces as their rhizoids are able to penetrate into the superficial layers of rocks (Longton, 1992; Adamo and Violante, 2000). After about 110 years the lichen-moss mat forms an almost continuous, thick layer (ca. 15 cm) of organic material on rock surfaces at Glaciar Frías.

The expansion of the lichen-moss mat is followed by the high recruitment and increase in coverage of vascular plants (Tables 2 and 3). It is well known that cryptogamic mats contribute to soil formation by entrapping particulate material and retaining remnants of dead vegetation (Longton, 1992). The formation of a soil layer is highly relevant in rocky environments where growth of large-sized vascular plants is controlled by the presence of sites with adequate soil volume for root deployment (Burbank and Phillips, 1983; Matthes-Sears and Larson, 1999). Cryptogams can also benefit vascular plants by ameliorating the physical and chemical environment, contributing nitrogen to the environment and entrapping plant seeds (Belnap et al., 2001; Breen and Lévesque, 2006). Therefore, the colonization of vascular plants observed during mid- and late-succession in the study area (Tables 2 and 3) likely has been facilitated by the earlier

TABLE 4

Correlation values (r) between vegetation ordination axes and explanatory variables. DCA (Detrended Correspondence Analysis); CCA (Canonical Correspondence Analysis). Species-environment correlations on DCA 1 and DCA2: $r = 0.86$ and $r = 0.37$, respectively; and on CCA 1 and CCA 2: $r = 0.89$ and $r = 0.83$, respectively. The statistical significance level of each explanatory variable was estimated in base of their partial effects (variability explained by the variable after accounting for the effects of other variables) as each variable is added to the model. All three categories of Topography were tested in conjunction for significance of their effect. Significance levels: $**P < 0.01$, $*P < 0.05$.

Variables	DCA correlations		CCA intra-set correlations		
	DCA 1	DCA 2	CCA 1	CCA 2	Partial F -value
Site age	-0.80	-0.05	-0.94	0.02	6.16**
Slope	0.24	0.08	0.26	-0.29	1.06
% bare rock outcrop	0.71	-0.04	0.81	-0.36	1.66*
% bare soil	0.26	0.15	0.33	0.75	2.89*
% pebble	0.30	-0.01	0.34	0.01	1.25
Canopy coverage	-0.19	-0.03	-0.21	-0.01	1.97*
Topography depressed	-0.16	0.02	-0.14	0.49	
Topography flat	0.02	0.22	0.01	-0.09	2.61**
Topography elevated	0.12	-0.17	0.12	-0.37	

progresses, herbs, erect shrubs, and small *Nothofagus dombeyi* trees (up to 2 m high) colonize the rock outcrops, forming a community with many overlapping strata (Table 3). On deglaciated terrains and alpine treelines in the southern Patagonian Andes, the emergence, growth, and survival of *N. antarctica* and *N. pumilio* seedlings increase under the canopy of shrubs (e.g. *Empetrum rubrum*) or adult trees (Cuevas, 2000; Henríquez and Lusk, 2005). Veblen and Ashton (1979) also reported that the establishment of *Nothofagus* spp. on volcanic ashes needs the protection from strong winds given by prostrate shrubs. Although individuals of *Nothofagus* are sometimes initial colonizers on recently deglaciated surfaces (Lawrence and Lawrence, 1959; Villalba et al., 1990; Masiokas, 2008), in general they are isolated trees associated with safe sites, such as cracks protected from wind (Veblen et al., 1989). For example, Villalba et al. (1990) reported a solitary 29-year-old *N. pumilio* growing on moraine 7 at Glaciar Frías, indicating that no other specimen succeeded in establishing on this moraine during the subsequent three decades. Therefore, the protection provided by other plants can be of outmost importance for the regeneration of a *Nothofagus* stand under the severe environmental conditions prevailing in the Patagonian Andes.

Our study provides a detailed example of vegetation development on bedrock slopes following deglaciation, which can be compared to successional trajectories reported in the literature for unconsolidated glacial deposits in the Patagonian Andes. Unfortunately, few studies on unconsolidated deposits have included cryptogams, making difficult the comparison of the earlier stages of vegetation development on the different surfaces. Lichenological studies nearby Glaciar San Rafael, a wet-maritime area in the Patagonian Andes, showed that *Placopsis* spp. not only dominate boulders and rock surfaces but also consolidated gravel (Galloway, 1992). The dominance of *Placopsis* spp. has also been reported in other Patagonian glaciers (Winchester and Harrison, 2000). At Glaciar Casa Pangué, approximately 4 km to the west of Glaciar Frías (Fig. 1), cryptogams account for 57% of the total coverage on moraines 40 years after deglaciation (Veblen et al., 1989). The lichen *Stereocaulon* and the moss *Racomitrium* are common constituents of the community in the earlier stages of vegetation development in Glaciar Casa Pangué (Veblen et al., 1989). Therefore, there is a striking similarity in the vegetation communities developing on rock faces and glacial deposits during the pioneer and mid-stages of the primary succession.

The woody carpet of prostrate shrubs observed on Glaciar Frías outcrops is a common stage of vegetation succession on areas exposed by glacier retreat or affected by volcanism in the

Patagonian Andes (Heusser, 1964; Veblen and Ashton, 1979; Veblen et al., 1989; and personal observations). On unconsolidated glacial deposits, the shrubby vegetation is usually replaced by *Nothofagus*-dominated forests within 75 years since glacier retreat (Heusser, 1964; Villalba et al., 1990). In contrast, we recorded a sparse coverage of *N. dombeyi* saplings in the rock outcrops exposed for more than 140 years (Table 3). This indicates that vegetation development is greatly delayed on rock outcrops compared to unconsolidated deposits. It is known that the slower community establishment on bare rock outcrops is related to the long time required for the formation of a ground vegetation cover and an organic matter layer (Asselin et al., 2006). These observations are consistent with our results indicating that once the lichen-moss mat develops, a large diversity of vascular plants rapidly colonized the rock surfaces. Indeed, the coverage of the vascular plants increased almost fivefold between sites 4 to 5, i.e. in a period of approximately 35 years (Fig. 3, Table 2). In summary, vegetation development requires at least 100 years longer on bedrock outcrops than on unconsolidated glacial deposits and is closely related to the time required for the formation of a cryptogamic carpet.

Acknowledgments

This work was funded by the Argentinean Agency for the Promotion of Science (grant PICTR02-186 and PICT32003), the Argentinean Council of Research and Technology (CONICET), and by the Inter-American Institute for Global Change Research (IAI) CRN II # 2047, which is supported by the U.S. National Science Foundation (grant GEO-0452325). We thank T. Ahti (Helsinki University, Finland) and G. Calabrese (University of Río Negro, Argentina) for help identifying lichen and moss species, respectively. The authors thank two anonymous reviewers for constructive criticism of the original manuscript.

References Cited

- Adamo, P., and Violante, P., 2000: Weathering of rocks and neogenesis of minerals associated with lichen activity. *Applied Clay Sciences*, 16: 229–256.
- Archer, A. C., Simpson, M. J. A., and MacMillan, B. H., 1973: Soils and vegetation of the lateral moraine at Malte Brun, Mount Cook region, New Zealand. *New Zealand Journal of Botany*, 11: 23–48.
- Armesto, J. J., Casassa, I., and Dollenz, O., 1992: Age structure and dynamics of Patagonian beech forests in Torres del Paine National Park, Chile. *Vegetatio*, 98: 13–22.

- Ashton, D. H., and Moore, G. M., 1978: Vegetation of Pleistocene block streams and block fields in Victoria: a successional interpretation. *Australian Journal of Ecology*, 3: 43–56.
- Asselin, H., Belleau, A., and Bergeron, Y., 2006: Factors responsible for the co-occurrence of forested and unforest rock outcrops in the boreal forest. *Landscape Ecology*, 21: 271–280.
- Barros, V., Cordon, V., Moyano, C., Mendez, R., Forquera, J., and Pizzio, O., 1983: *Cartas de Precipitación de la Zona Oeste de las Provincias de Río Negro y Neuquén*. Cinco Saltos, Argentina: Facultad de Ciencias Agrarias, Universidad Nacional del Comahue, 27 pp.
- Belnap, J., Prasse, R., and Harper, K. T., 2001: Influence of biological soil crusts on soil environments and vascular plants. In Belnap, J., and Lange, O. L. (eds.), *Biological Soil Crusts: Structure, Function, and Management*. Berlin-Heidelberg: Springer-Verlag, 281–300.
- Breen, K., and Lévesque, E., 2006: Proglacial succession of biological soil crusts and vascular plants: biotic interactions in the High Arctic. *Canadian Journal of Botany*, 84: 1714–1731.
- Brummitt, R. K., and Powell, C. E., 1992: *Authors of Plants Names*. Kew: Royal Botanical Garden, 732 pp.
- Burbanck, M. P., and Phillips, D. L., 1983: Evidence of plant succession on granite outcrops of the Georgia piedmont. *The American Midland Naturalist*, 109: 94–104.
- Cuevas, J. G., 2000: Tree recruitment at the *Nothofagus pumilio* alpine timberline in Tierra del Fuego, Chile. *Journal of Ecology*, 88: 840–855.
- Cutler, N. A., Belyea, L. R., and Dugmore, A. J., 2008: The spatiotemporal dynamics of a primary succession. *Journal of Ecology*, 96: 231–246.
- Chapin, F. S., III, Walker, L. R., Fastie, C. L., and Sharman, L. C., 1994: Mechanisms of primary succession following deglaciation at Glacier Bay, Alaska. *Ecological Monographs*, 64: 149–175.
- De Agostini, A., 1945: *Andes Patagónicas*. Buenos Aires: Guillermo Kraft Ltda., 437 pp.
- del Moral, R., Titus, J. H., and Cook, A. M., 1995: Early primary succession on Mount St. Helens, Washington, USA. *Journal of Vegetation Science*, 6: 107–120.
- Dollenz, O., 1991: Sucesión vegetal en el sistema morrenico del Glaciar Dickson, Magallanes, Chile. *Anales del Instituto de la Patagonia Serie Ciencias Naturales*, 20: 49–60.
- Ezcurra, C., and Brion, C., 2005: *Plantas del Nahuel Huapi: Catálogo de la Flora Vasculare del Parque Nacional Nahuel Huapi, Argentina*. San Carlos de Bariloche, Argentina: Universidad Nacional del Comahue, 70 pp.
- Fonck, F., 1896: *Viajes del Fray Francisco Menéndez a la Cordillera*. Valparaiso, Chile: Niemeyer, 528 pp.
- Foster, B. L., and Tilman, D., 2000: Dynamic and static views of succession: testing the descriptive power of the chronosequence approach. *Plant Ecology*, 146: 1–10.
- Galloway, D. J., 1992: Lichens of Laguna San Rafael, Parque Nacional “Laguna San Rafael”, southern Chile: indicators of environmental change. *Global Ecology and Biogeography Letters*, 2: 37–45.
- Henríquez, J. M., 2004: Influencia de los defecaderos de camélidos sobre el desarrollo vegetal y riqueza de especies en morenas glaciales, Tierra del Fuego. *Revista Chilena de Historia Natural*, 77: 501–508.
- Henríquez, J. M., and Lusk, C. H., 2005: Facilitation of *Nothofagus antarctica* (Fagaceae) seedlings by the prostrate shrub *Empetrum rubrum* (Empetraceae) on glacial moraines in Patagonia. *Austral Ecology*, 30: 877–882.
- Heusser, C. J., 1960: Late-Pleistocene environments of the Laguna de San Rafael area, Chile. *Geographical Review*, 50: 555–577.
- Heusser, C. J., 1964: Some pollen profiles from Laguna San Rafael area, Chile. In Cranwell, L. M. (ed.), *Ancient Pacific Floras*. Honolulu, U.S.A.: University of Hawaii Press, 95–114.
- Hodkinson, I. D., Coulson, S. J., and Webb, N. R., 2003: Community assembly along proglacial chronosequences in the High Arctic: vegetation and soil development in north-west Svalbard. *Journal of Ecology*, 91: 651–663.
- Jongman, R. H. G., ter Braak, C. J. F., and Van Tongeren, O. F. R., 1995: *Data Analysis in Community and Landscape Ecology*. Cambridge, U.K.: Cambridge University Press, 299 pp.
- Körner, C., 2005: The green cover of mountains in a changing environment. In Huber, U. M., Bugmann, H. K. M., and Reasoner, M. A. (eds.), *Global Change and Mountain Regions, an Overview of Current Knowledge*. Dordrecht, The Netherlands: Springer, 367–375.
- Kurina, L. M., and Vitousek, P. M., 2001: Nitrogen fixation rates of *Stereocaulon vulcani* on young Hawaiian lava flows. *Biogeochemistry*, 55: 179–194.
- Lambhead, P. J. D., Platt, H. M., and Shaw, K. M., 1983: The detection of differences among assemblages of marine benthic species based on an assessment of dominance and diversity. *Journal of Natural History*, 17: 859–874.
- Lawrence, D. B., and Lawrence, E. G., 1959: *Recent Glacier Variations in Southern South America*. New York: American Geographical Society, 39 pp.
- Leps, J., and Šmilauer, P., 1999: *Multivariate Analysis of Ecological Data*. Ceske Budejovice: Faculty of Biological Sciences, University of South Bohemia, 110 pp.
- Lindsay, D. C., 1978: The role of lichens in Antarctic ecosystems. *The Bryologist*, 81: 268–276.
- Longton, R. E., 1992: The role of bryophytes and lichens in terrestrial ecosystems. In Bates, J. W., and Farmer, A. M. (eds.), *Bryophytes and Lichens in a Changing Environment*. Oxford: Oxford University Press, 32–75.
- Luckman, B. H., and Villalba, R., 2001: Assessing the synchronicity of glacier fluctuations in the western Cordillera of the Americas during the last millennium. In Makgraf, V. (ed.), *Interhemispheric Climate Linkages*. San Diego: Academic Press, 119–140.
- Magurran, A. E., 1988: *Ecological Diversity and its Measurements*. Princeton: Princeton University Press, 179 pp.
- Masiokas, M., 2008: Climate and glacier variability during past centuries in the north and south Patagonian Andes of Argentina. Ph.D. thesis. Faculty of Graduate Studies, The University of Western Ontario, London, Ontario, Canada, 268 pp.
- Masiokas, M., Luckman, B. H., Villalba, R., Ripalta, A., and Rabassa, J., 2010: Little Ice Age fluctuations of Glaciar Río Manso in the north Patagonian Andes of Argentina. *Quaternary Research*, 73: 96–106.
- Matthes-Sears, U., and Larson, D. W., 1999: Limitations to seedling growth and survival by the quantity and quality of rooting space: implications for the establishment of *Thuja occidentalis* on cliff faces. *International Journal of Plant Sciences*, 160: 122–128.
- Matthes-Sears, U., and Larson, D. W., 2006: Microsite and climatic controls of tree population dynamics: an 18-year study on cliffs. *Journal of Ecology*, 94: 402–414.
- Matthews, J. A., 1992: *The Ecology of Recently Deglaciated Terrain. A Geological Approach to Glacier Forelands and Primary Succession*. Cambridge, U.K.: Cambridge University Press, 386 pp.
- Matthews, J. A., 1999: Disturbance regimes and ecosystem response on recently-deglaciated substrates. In Walker, L. R. (ed.), *Ecosystems of Disturbed Ground*. Amsterdam: Elsevier, 17–37.
- Matthews, J. A., and Whittaker, R. J., 1987: Vegetation succession on the Storbreen Glacier Foreland, Jotunheimen, Norway: a review. *Arctic and Alpine Research*, 19: 385–395.
- Nash, T. H., III, 1996: Photosynthesis, respiration, productivity and growth. In Nash, T. H., III (ed.), *Lichen Biology*. Cambridge, U.K.: Cambridge University Press, 88–120.

- Opazo Medina, B. M., Torres Ribeiro, K., and Rubio Scarano, F., 2006: Plant-plant and plant-topography interactions on a rock outcrop at high altitude in southeastern Brazil. *Biotropica*, 38: 27–34.
- Orwin, J. F., 1970: Lichen succession on recently deposited rock surfaces. *New Zealand Journal of Botany*, 8: 452–477.
- Orwin, J. F., 1972: The effect of environment on assemblages of lichens growing on rock surfaces. *New Zealand Journal of Botany*, 10: 37–47.
- Paul, F., Kääb, A., and Haeberli, W., 2007: Recent glacier changes in the Alps observed by satellite: consequences for future monitoring strategies. *Global and Planetary Change*, 56: 111–122.
- Pelto, M. S., 2009: Forecasting temperate alpine glacier survival from accumulation zone observations. *The Cryosphere Discussions*, 3: 323–350.
- Pickett, S. T. A., 1989: Space-for-time substitution as an alternative to long-term studies. In Likens, G. E. (ed.), *Long-Term Studies in Ecology*. New York: Springer, 110–135.
- Pisano, V. E., 1978: Establecimientos de *Nothofagus betuloides* (Mirb.) Blume (Coigue de Magallanes) en un valle en proceso de deglaciación. *Anales del Instituto de la Patagonia, Serie Ciencias Naturales*, 9: 107–128.
- Rabassa, J., Rubulis, S., and Suarez, J., 1978: Los glaciares del Monte Tronador, Parque Nacional Nahuel Huapi, Río Negro, Argentina. *Anales de Parques Nacionales*, 14: 259–295.
- Rabassa, J., Rubulis, S., and Suarez, J., 1979: Rate of formation and sedimentology of (1976–1978) push-moraines, Frías Glacier, Mount Tronador (41° 10'S; 71° 53'W), Argentina. In Schlucher, C. H. (ed.), *Moraines and Varves*. Rotterdam: Balkema, 65–80.
- Rabassa, J., Rubulis, S., and Suarez, J., 1981: Moraine in-transit as parent material for soil development and the growth of Valdivian Rain Forest on moving ice: Casa Pangué Glacier, Mount Tronador (Lat. 41°10'S), Chile. *Annals of Glaciology*, 2: 97–102.
- Rivera, A., and Casassa, G., 2004: Ice elevation, areal, and frontal changes of glaciers from National Park Torres del Paine, southern Patagonia Icefield. *Arctic, Antarctic, and Alpine Research*, 36: 379–389.
- Sarthou, C., Kounda-Kiki, C., Vaculik, A., Mora, P., and Ponge, J.-F., 2009: Successional patterns on tropical inselbergs: a case study on the Nouragues inselberg (French Guiana). *Flora*, 204: 396–407.
- Shure, D. J., and Ragsdale, H. L., 1977: Patterns of primary succession on granite outcrop surfaces. *Ecology*, 58: 993–1006.
- Sommerville, P., Mark, A. F., and Wilson, J. B., 1982: Plant succession on moraines of the upper Dart Valley, southern South Island, New Zealand. *New Zealand Journal of Botany*, 20: 227–244.
- Steffen, H., 1909: *Viajes de Exploracion i Estudio en la Patagonia Occidental: 1892–1902*. Santiago de Chile: Imprenta Cervantes, 233 pp.
- Svoboda, J., and Henry, G. H. R., 1987: Succession in marginal Arctic environments. *Arctic and Alpine Research*, 19: 373–384.
- Uno, G. E., and Collins, S. L., 1987: Primary succession on granite outcrops in southwestern Oklahoma. *Bulletin of the Torrey Botanical Club*, 114: 387–392.
- Veblen, A. T., and Ashton, D. H., 1979: Successional pattern above timberline in south-central Chile. *Vegetatio*, 40: 39–47.
- Veblen, T. T., Ashton, D. H., Rubulis, S., Lorenz, D. C., and Cortes, M., 1989: *Nothofagus* stand development on in-transit moraines, Casa Pangué Glacier, Chile. *Arctic and Alpine Research*, 21: 144–155.
- Vetaas, O. R., 1994: Primary succession of plant assemblages on a glacier foreland—Bodalsbreen, southern Norway. *Journal of Biogeography*, 21: 297–308.
- Villalba, R., Leiva, J. C., Rubulis, S., Suarez, J., and Lenzano, L., 1990: Climate, tree-ring, and glacial fluctuations in the Río Frías Valley, Río Negro, Argentina. *Arctic, Antarctic, and Alpine Research*, 22: 215–232.
- Vitousek, P. M., 1994: Potential nitrogen fixation during primary succession in Hawai'i Volcanoes National Park. *Biotropica*, 26: 234–240.
- Walker, L. R., and Chapin, F. S., III, 1987: Interactions among processes controlling successional change. *Oikos*, 50: 131–135.
- Walker, L. R., Wardle, D. A., Bardgett, R. D., and Clarkson, B. D., 2010: The use of chronosequences in studies of ecological succession and soil development. *Journal of Ecology*. doi:10.1111/j.1365-2745.2010.01664.x.
- Wardle, P., 1980: Primary succession in Westland National Park and its vicinity, New Zealand. *New Zealand Journal of Botany*, 18: 221–232.
- Winchester, V., and Harrison, S., 2000: Dendrochronology and lichenometry: colonization, growth rates and dating of geomorphological events on the east side of the North Patagonian Icefield, Chile. *Geomorphology*, 34: 181–194.
- Wiser, S. K., Peet, R. K., and White, P. S., 1996: High-elevation rock outcrop vegetation of the southern Appalachian Mountains. *Journal of Vegetation Science*, 7: 703–722.
- Zuloaga, I. O., and Morrone, O., 1999a: *Catálogo de las Plantas Vasculares de Argentina I*. St. Louis, Missouri: Missouri Botanical Garden, 323 pp.
- Zuloaga, I. O., and Morrone, O., 1999b: *Catálogo de las Plantas Vasculares de Argentina II*. St. Louis, Missouri: Missouri Botanical Garden, 1267 pp.

MS accepted July 2010