



Yellow-crowned Elaenia *Myiopagis flavivertex*, new to Colombia

Authors: Flórez, Pablo , and Kirwan, Guy M.

Source: Bulletin of the British Ornithologists' Club, 137(2) : 150-151

Published By: British Ornithologists' Club

URL: <https://doi.org/10.25226/bboc.v137i2.2017.a4>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Yellow-crowned Elaenia *Myiopagis flavivertex*, new to Colombia

by Pablo Flórez & Guy M. Kirwan

Received 3 January 2017; revised 10 May 2017; published 18 June 2017

<http://zoobank.org/urn:lsid:zoobank.org:pub:49E0212D-2D55-40EE-AFE1-55E545D935B6>

An inhabitant of *várzea* and flooded, swampy forest, as well as more locally in flooded forest in sandy-soil areas, Yellow-crowned Elaenia *Myiopagis flavivertex* is comparatively widely distributed across much of northern and western Amazonia. It is known from north-east and southern Venezuela (including Monagas south to Delta Amacuro, and Amazonas), the Guianas, north-east Ecuador (Sucumbíos, Napo), north-east Peru (in eastern Loreto, Ucayali) and Brazil across the central Amazon Basin, but also south along the Madeira River to northern Rondônia (Fitzpatrick 2004, Ridgely & Tudor 2009). There are no published records in Colombia (Salaman *et al.* 2010, Donegan *et al.* 2016), although Hilty & Brown (1986) suggested that it might be found near San Fernando de Atabapo, which is just inside neighbouring Venezuela, and McMullan & Donegan (2014) illustrated the species given that its occurrence was expected. Recently, we have recorded the species three times in the extreme east of the country, in the environs of Puerto Inírida, eastern Guainía, thereby confirming its presence, precisely in the region predicted by Hilty & Brown (1986).

On 7 October 2014, in the environs of Caño Matraca, just upstream of Puerto Inírida on the left (west) bank of the Inírida River, GMK heard and briefly observed at least one *M. flavivertex* inside the canopy of lower-stature flooded *várzea*, together with N. Bostock *et al.* Unfortunately, at the time, they and the other observers present, including PF, were all focused on photographing and sound-recording a pair of Orinoco Softtails *Thripophaga cherriei*, and GMK did not capture any vocalisations in the background of his recordings. Subsequently, the observation was more or less forgotten about for almost two years—especially as GMK was unfamiliar with the species' (non-)status in Colombia but also because its presence in the extensive *várzea* forest around Puerto Inírida appeared so unsurprising. In mid-September 2016, PF sent GMK photographs and a sound-recording (XC345120; www.xeno-canto.org) from the same region, which GMK immediately recognised as being of *M. flavivertex*. It was only then that GMK noticed that the species was not previously known to occur in Colombia and a little later came across his notes from 2014 again.

PF's observations were made on 12 September 2016, when a single was seen with a mixed-species flock (which this species does not routinely follow), in *várzea* forest beside the Inírida River. In addition to the recording, his photographs show the majority of the salient features associated with a *Myiopagis* elaenia and specifically *M. flavivertex*, other than the wingbars, namely: the small bill with pinkish base to mandible, dark eyes, rather flat crown with no crest, greyish-olive breast and throat merging into more yellowish belly, lack of an obvious pale superciliary except on the supraloral region, slightly long-tailed appearance, and yellow-fringed tertials and flight feathers. However, although the underparts pattern differs from that of the otherwise very similar Forest Elaenia *M. gaimardii* (which has a greyish-white throat and upper breast, with a hint of olive streaking), the latter species is best identified by its voice. The only Tyrannidae that might be confused with *M. flavivertex* by voice is Euler's Flycatcher *Lathrotriccus euleri*, but the latter is very different in plumage. While *M. flavivertex* is a specialist of *várzea* forest, whereas Forest Elaenia is typically

found in dry-land forest habitats, although it also occurs in inundated habitats, at least in Suriname and Venezuela (P. Boesman *in litt.* 2017).

Subsequently, T. M. Donegan (*in litt.* 2016, 2017) alerted us to the fact that other observers have also reported *M. flavivertex* in the same area of Colombia, among them T. Friedel, who photographed one individual on 30 January 2016 (http://www.birdphotos.com/photos/index.php?q=gallery&g2_itemId=228164&g2_imageViewsIndex=1). Furthermore, during another visit to the area, on 16 February 2017, PF, GMK, G. R. R. Brito and J. Lobel heard, and GMK sound-recorded, the species in the, at this season, dry *várzea* forest along the trail to Caño Matraca (XC365007)

A recent ornithological inventory of the Puerto Inírida region recorded only Forest Elaenia there, listing this species as occurring in both *terra firme* and *várzea* forests (Stiles & Beckers 2016), which suggests that *M. flavivertex* might have been to some degree overlooked by previous workers, perhaps because it is easily missed even when vocalising. The species' discovery in easternmost Colombia is plainly unsurprising, given both the extensive availability of suitable habitat in this part of Guainía and the presence of *M. flavivertex* at several localities in immediately adjacent Venezuela (south-west Amazonas; Hilty 2003). Nevertheless, that it should have escaped previous notice, despite considerable recent survey effort around Puerto Inírida (Stiles & Beckers 2016), surely is somewhat remarkable and hints at the possibility of still-further ornithological discoveries in this, and many other regions of the country. Indeed, while the present manuscript was in production, Ramírez *et al.* (submitted) reported several records of *M. flavivertex* from the departments of Guaviare and Meta, indicating that its range in eastern Colombia is considerably broader than believed. Given records very close to the Colombian border inside Peru (e.g. XC16599; <http://www.hbw.com/ibc/photo/yellow-crowned-elaenia-myiopagis-flavivertex/yellow-crowned-elaenia-perched>), the species can also be expected to occur around Leticia, in the extreme south of dpto. Amazonas.

Acknowledgements

We are grateful to the following who made useful comments on the manuscript or related issues: Peter Boesman, Diego Calderón, Thomas Donegan, Jeremy Minns and Chris Sharpe. We are grateful to Guilherme Brito and John Lobel for their company in February 2017.

References:

- Donegan, T., Verhelst, J. C., Ellery, T., Cortés-Herrera, O. & Salaman, P. 2016. Revision of the status of bird species occurring or reported in Colombia 2016 and assessment of BirdLife International's new parrot taxonomy. *Conserv. Colombiana* 24: 12–36.
- Fitzpatrick, J. 2004. Yellow-crowned Elaenia (*Myiopagis flavivertex*). P. 266 in del Hoyo, J., Elliott, A. & Christie, D. A. (eds.) *Handbook of the birds of the world*, vol. 9. Lynx Edicions, Barcelona.
- Hilty, S. L. 2003. *Birds of Venezuela*. Princeton Univ. Press.
- McMullan, M. & Donegan, T. 2014. *Field guide to the birds of Colombia*. Second edn. Fundación ProAves, Bogotá.
- Ramírez, W. A., Arredondo, C., López, R. C. & Chaparro-Herrera, S. submitted. Range extensions for Yellow-crowned Elaenia *Myiopagis flavivertex* and Dugand's Antwren *Herpsilochmus dugandi* in eastern Colombia. *Bull. Brit. Orn. Cl.*
- Ridgely R. S. & Tudor, G. 2009. *Birds of South America: passerines*. Christopher Helm, London.
- Salaman, P., Donegan, T. M. & Caro, D. 2010. *Checklist of the birds of Colombia*. Fundación ProAves, Bogotá.
- Stiles, F. G. & Beckers, J. 2016. Un inventario de las aves de la región de Inírida, Guainía, Colombia. *Orn. Colombiana* 15: 21–52.

Addresses: Pablo Flórez, Calle 51A, No. 81 B22 Apto. 503, Medellín, 05001000, Colombia, e-mail: pablo@multicolorbirdingcolombia.com. Guy M. Kirwan, Field Museum of Natural History, 1400 South Lakeshore Drive, Chicago, IL 60605, USA, and Museu Nacional, Departamento de Vertebrados, Setor de Ornitologia, Quinta da Boa Vista, CEP 20940-040, Rio de Janeiro, RJ, Brazil, e-mail: GMKirwan@aol.com