



## **An Acoustic Trap to Survey and Capture Two Neoscapteriscus Species**

Authors: Rohde, Barukh B., Allen, Pablo E., Benda, Nicole, Brun, Abe, Mankin, Richard W., et al.

Source: Florida Entomologist, 102(3) : 654-657

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.102.0316>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# An acoustic trap to survey and capture two *Neoscapteriscus* species

Barukh B. Rohde<sup>1</sup>, Pablo E. Allen<sup>2</sup>, Nicole Benda<sup>2</sup>, Abe Brun<sup>3</sup>, Richard W. Mankin<sup>4</sup>, and Adam G. Dale<sup>2,\*</sup>

Invasive *Neoscapteriscus* (Orthoptera: Grilloalpidae) mole crickets damage turfgrasses and pastures throughout the southeastern USA (Hayslip 1943; Walker & Nickle 1981; Liskey 2000). The southern mole cricket, *Neoscapteriscus borellii* (Giglio-Tos) (Orthoptera: Grilloalpidae), and tawny mole cricket, *Neoscapteriscus vicinus* (Scudder) (Orthoptera: Grilloalpidae), were introduced inadvertently into southeastern Georgia before 1900, and thereafter spread rapidly throughout the southeastern US (Walker & Nickle 1981). These insects feed on plant roots and tunnel through the soil, uprooting and killing plants (Xu et al. 2012), which reduces forage for cattle, increases turfgrass sod production time, and reduces turfgrass quality and playability on golf courses (Frank & Walker 2006). Because mole crickets remain underground for much of their life cycle, the cause of initially observed damage may remain unknown until populations rise to destructive levels, as with many other subterranean insects (e.g., Mankin et al. 2007).

*Neoscapteriscus borellii* and *N. vicinus* attract mates using acoustic communication, and were among the earliest targets of insect acoustic trapping technology (Ulagaraj & Walker 1973; Walker 1996). During spring and fall mating seasons (Hayslip 1943; Walker & Nation 1982), males build carefully constructed, horn-shaped burrows in which they stridulate for several h after sunset (Ulagaraj 1976). The stridulations are amplified by resonances within the burrows (Bennet-Clark 1987, 1999), generating loud calls that attract conspecific, virgin, and mated flying females (Ulagaraj 1976; Forrest 1986). The females land, search for the burrow entrance, and mate with the calling male (Ulagaraj & Walker 1973; Ulagaraj 1976). Also, other males are attracted to the area around the burrow, possibly because the presence of calling males indicates suitable habitat for colonization (Ulagaraj 1975). By replicating the mating call and broadcasting it from a trapping device (Ulagaraj & Walker 1973; Ulagaraj 1975; Forrest 1980, 1983; Walker 1982), researchers can take advantage of the phonotactic behavior of both sexes to monitor local populations, alert turf managers to *Neoscapteriscus* presence, and test efficacy of control strategies (e.g., Parkman et al. 1993; Frank & Walker 2006; Kerr et al. 2014; Mhina et al. 2016; Aryal et al. 2019).

The first mole cricket traps were constructed in the 1970s with a battery-operated tape recorder as the sound source and a 1.2-m-diam sheet-metal funnel attached to a jar as the catching device (Ulagaraj 1975, 1976; Ulagaraj & Walker 1975). Walker (1982) and colleagues placed a synthesizer, amplifier, and power source with a switch to toggle between mole cricket species' mating calls into a battery-pow-

ered box that became the first-ever semi-automatic mole cricket trap. Walker (1996) and colleagues further advanced these traps by building the first microprocessor-based mole cricket trap, powered by a 12-volt battery, and furnished with a built-in photocell to autonomously time broadcasts. Development of such traps initially required electrical and audio engineering skills, and most mole cricket researchers contracted out their broadcast systems (e.g., Thompson & Brandenburg 2004).

Advances in audio and computer technology enabled Dillman et al. (2014) to construct an acoustic trap to survey *N. borellii* by combining a commercially available Arduino microcontroller platform (Mankin et al. 2012, 2016; Johnson et al. 2018) with an open-source 'wave shield' (Adafruit Inc., New York, New York, USA) that allows a nontechnical user to play recorded sounds from an SD memory card into audio output. Dillman et al. (2014) connected the audio output to an amplifier and a pair of motorcycle speakers (Pyle Audio, Brooklyn, New York, USA), powered with a 12-volt battery. Their initial collection device was an assembly of two 1.5-m diam wading pools constructed as in Thompson & Brandenburg (2004). One wading pool was filled to about 10 cm depth pure sand mixed with perlite to harbor captured mole crickets and limit desiccation, and the second pool was empty with 6 holes cut into it, and placed over top of the sand-filled pool, minimizing escape by captured mole crickets and predation from vertebrate predators (Thompson & Brandenburg 2004). This wading pool trap design enables retrieval of live specimens for use in experiments.

To conduct studies described here, we constructed a modified version of the Dillman et al. (2014) acoustic wading-pool trap, hereafter termed as the Borelli Vicinus Acoustic Pool trap, with 2 modifications. One modification was the use of an SD memory card that broadcast 4 s of separate recordings from both *N. borellii* and *N. vicinus* played in a continuous loop (Fig. 1), and the second was the addition of a 12-volt solar panel (model #41007; RDK Products, Buford, Georgia, USA) to recharge the 12-volt battery during daylight, eliminating the need for routine field recharges (Fig. 2A). We obtained recordings of each species' mating call from the Singing Insects of North America database (<http://entnemdept.ufl.edu/Walker/buzz/index.htm>). As in Dillman et al. (2014), the sound pressure level was set to 106 decibels at 15 cm from the speaker.

In 2017, we deployed 1 Borelli Vicinus Acoustic Pool trap at each of 7 pasture sites throughout north-central Florida (Fig. 2B). We collected mole crickets for 10 wk from 23 Apr to 6 Jul 2017. In 2018, we redeployed the same traps at 5 pasture locations (Fig. 2B). Mole crickets were collected for


<sup>1</sup>University of Florida, Department of Electrical and Computer Engineering, Gainesville, Florida 32611, USA; E-mail: barukh94-work@yahoo.com (B. B. R.)

<sup>2</sup>University of Florida, Entomology and Nematology Department, Gainesville, Florida 32611, USA; E-mails: pabloallen@ufl.edu (P. E. A.), nbenda@ufl.edu (N. B.), agdale@ufl.edu (A. G. D.)

<sup>3</sup>Custom Engineered Solutions, West Hempstead, New York 11552, USA; E-mail: anbk1@yahoo.com (A. B.)

<sup>4</sup>USDA ARS, Center for Medical, Agricultural, and Veterinary Entomology, Gainesville, Florida 32608, USA; E-mail: richard.mankin@ars.usda.gov (R. W. M.)


\*Corresponding author; E-mail: agdale@ufl.edu



**Fig. 1.** (A) Oscillogram and (B) spectrogram of *Neoscapteriscus borellii* and *N. vicinus* species-specific mating calls broadcast from speakers in a continuous loop. Darker shading in spectrogram indicates greater energy at specified frequency and time.

8 wk from 7 Feb to 5 Apr 2018. Sites were at least 48 km apart in 2017 and 16 km apart in 2018, and spanned approximately 9,000 km<sup>2</sup>. Each trap was placed in the middle of a bahiagrass (*Paspalum notatum* Flügge; Poaceae) pasture away from any acoustic (e.g., anthropogenic noise sources) or structural (e.g., trees, buildings) interference. Mole crickets were collected from the traps once per wk, and immediately placed individually into clear plastic vials, labeled by site and date, and taken to the laboratory for identification. All collected mole crickets were identified to species.

In 2017 and 2018, the traps caught varying numbers of *N. borellii* and *N. vicinus* at each location (Fig. 2A, B). In sum, we collected 675 *N. borellii* and 13 *N. vicinus* in 2017, and 71 *N. borellii* and 438 *N. vicinus* in 2018. The bias towards *N. borellii* in 2017 and *N. vicinus* in 2018 was expected because the 2017 surveys were conducted after peak *N. vicinus* flight activity had occurred, and the 2018 surveys before *N. borellii* peak flight activity. By connecting a solar panel to each trap's battery, no batteries needed a recharge through the dura-



**Fig. 2.** (A) The Borelli Vicinus Acoustic Pool trap used to attract and capture *N. borellii* and *N. vicinus* mole crickets throughout the study. (B) Total *Neoscapteriscus* mole crickets captured in Borelli Vicinus Acoustic Pool traps from late spring (23 Apr to 6 Jul) 2017 and early spring (7 Feb to 5 Apr) 2018 in northern Florida, USA. Dark circles represent *N. borellii* and light circles represent *N. vicinus*. The size of each circle indicates the total individuals captured per site from 0 to 250, as illustrated in the figure legend. The same traps with the exact same settings were used in both yr.

tion of this study. As expected based on previous studies (Ulagaraj & Walker 1973), traps captured predominantly female mole crickets (> 85% of captured individuals). Nevertheless, several hundred individuals of both species were captured by Borelli Vicinus Acoustic Pool traps each year, indicating that our Borelli Vicinus Acoustic Pool traps, playing a continuous loop of 4-s intervals of the 2 species-specific mating calls (Fig. 1), successfully attracted and captured both invasive mole cricket species. Our current Borelli Vicinus Acoustic Pool trap design costs approximately \$400 per trap for materials, a one-time \$500 program license fee for use on multiple traps, trap assembly labor costs, and technical training or support, which is prohibitive for many. Therefore, future work should develop trap designs that cost a fraction of this price and require little technical knowledge or training, enabling land managers and researchers to employ these tools to monitor invasive pests (not as a pest control tool) on their respective properties.

This research was supported by USDA-NIFA award number 2016-38503-25899 to AGD, the University of Florida Agricultural Experiment Station to AGD, and the National Science Foundation Graduate Research Fellowship DGE-1315138 and DGE-1842473 to BBR. We thank Custom Engineered Solutions, LLC (West Hempstead, New York, USA) for acoustic trap development and troubleshooting. We also thank Rebecca Perry and Alexander LoCastro for assistance with field work and data collection. Mention of a trademark or proprietary product is solely for the purpose of providing specific information and does not constitute a guarantee or warranty of the product by the USDA, and does not imply its approval to the exclusion of other products that also may be suitable. The USDA is an equal opportunity employer.

## Summary

There is an extensive history of mole cricket integrated pest management (IPM) research in Florida, USA (Kerr et al. 2014; Mhina et al. 2016), much of which has incorporated acoustic trapping as a monitoring tool. The acoustic trap design described in this report provides a method for surveying 2 *Neoscapteriscus* mole cricket species relatively autonomously at low cost, which can facilitate future efforts to study the biology, ecology, and distribution of invasive mole crickets (e.g., Walker 1988). In a broader context, however, there remains considerable need to reduce the costs and simplify the technology of these and other traps based on inexpensive microcontroller platforms, not only for *Neoscapteriscus* species, but also for other pests that mate based on either acoustic or vibrational communication (Mankin 2012).

Key Words: mole cricket; forage; turfgrasses; survey

## Sumario

Existe una extensa historia de investigaciones sobre el manejo integrado de plagas (MIP) del grillo topo en la Florida, EE. UU. (Kerr et al. 2014; Mhina et al. 2016), una gran parte de las cuales ha incorporado el atrapamiento acústico como herramienta de monitoreo. El diseño de la trampa acústica descrito en este informe proporciona un método para estudiar 2 especies de grillo topos del género *Neoscapteriscus* de forma relativamente autónoma a bajo costo, lo que puede facilitar los esfuerzos futuros para estudiar la biología, ecología y distribución de los grillo topos invasivos (por ejemplo, Walker 1988). Sin embargo, en un contexto más amplio, sigue existiendo una necesidad considerable de reducir los costos y simplificar la tecnología de estas y otras

trampas basadas en plataformas económicas de microcontroladores, no solo para las especies de *Neoscapteriscus*, sino también para otras plagas que se aparean en base a la comunicación acústica o vibracional (Mankin 2012).

Palabras Clave: grillo topo; forraje; céspedes; sondeo

## References Cited

- Aryal SK, Lu D, Le K, Allison L, Gerke C, Dillman AR. 2019. Sand crickets (*Gryllus firmus*) have low susceptibility to entomopathogenic nematodes and their pathogenic bacteria. *Journal of Invertebrate Pathology* 160: 54–60.
- Bennet-Clark HC. 1987. The tuned singing burrow of mole crickets. *Journal of Experimental Biology* 128: 383–409.
- Bennet-Clark HC. 1999. Resonators in insect sound production: how insects produce loud pure-tone songs. *Journal of Experimental Biology* 202: 3347–3357.
- Dillman AR, Cronin CJ, Tang J, Gray DA, Sternberg PW. 2014. A modified mole cricket lure and description of *Scapteriscus borellii* (Orthoptera: Gryllotalpidae) range expansion and calling song in California. *Environmental Entomology* 43: 146–156.
- Forrest TG. 1980. Phonotaxis in mole crickets: its reproductive significance. *Florida Entomologist* 63: 45–53.
- Forrest TG. 1983. Calling songs and mate choice in mole crickets, pp. 185–204. In Gwynne DT, Morris GK [eds.], *Orthopteran Mating Systems: Sexual Competition in a Diverse Group of Insects*. Westview, Boulder, Colorado, USA.
- Forrest TG. 1986. Oviposition and maternal investment in mole crickets (Orthoptera: Gryllotalpidae): effects of season, size, and senescence. *Annals of the Entomological Society of America* 79: 918–924.
- Frank JH, Walker TJ. 2006. Permanent control of pest mole crickets (Orthoptera: Gryllotalpidae: *Scapteriscus*) in Florida. *American Entomologist* 52: 138–144.
- Hayslip NC. 1943. Notes on biological studies of mole crickets at Plant City, Florida. *Florida Entomologist* 26: 33–46.
- Johnson BJ, Rohde BB, Zeak N, Staunton KM, Prachar T, Ritchie SA. 2018. A low-cost, battery-powered acoustic trap for surveilling male *Aedes aegypti* during rear-and-release operations. *PLoS ONE* 13: e0201709. <https://doi.org/10.1371/journal.pone.0201709> (last accessed 22 May 2019).
- Kerr CR, Leppla NC, Buss EA, Frank JH. 2014. Mole cricket IPM guide for Florida. UF/IFAS EDIS IPM 206. University of Florida, Gainesville, Florida, USA.
- Liskey E. 2000. Turf's most (un) wanted pests. *Grounds Maintenance* 35: 18–23.
- Mankin RW. 2012. Applications of acoustics in insect pest management. *Commonwealth Agricultural Bureaux International Reviews* 7: 001. doi: 10.1079/PAVSNNR20127001 (last accessed 22 May 2019).
- Mankin RW, Hubbard JL, Flanders KL. 2007. Acoustic indicators for mapping infestation probabilities of soil invertebrates. *Journal of Economic Entomology* 100: 790–800.
- Mankin RW, Rohde B, McNeill S. 2016. Vibrational duetting mimics to trap and disrupt mating of the devastating Asian citrus psyllid insect pest. *Proceedings of Meetings on Acoustics* 25: 010006. <https://www.ars.usda.gov/ARSUserFiles/3559/publications/poma25-2016.pdf> (last accessed 22 May 2019).
- Mhina GJ, Leppla NC, Thomas MH, Solis D. 2016. Cost effectiveness of biological control of invasive mole crickets in Florida pastures. *Biological Control* 100: 108–115.
- Parkman JP, Frank JH, Nguyen KB, Smart GC. 1993. Dispersal of *Steinernema scapterisci* (Rhabditida: Steinernematidae) after inoculative applications for mole cricket (Orthoptera: Gryllotalpidae) control in pastures. *Biological Control* 3: 226–232.
- Thompson SR, Brandenburg RL. 2004. A modified pool design for collecting adult mole crickets (Orthoptera: Gryllotalpidae). *Florida Entomologist* 87: 582–584.
- Ulagaraj SM. 1975. Mole crickets: ecology, behavior, and dispersal flight (Orthoptera: Gryllotalpidae: *Scapteriscus*). *Environmental Entomology* 4: 265–273.
- Ulagaraj SM. 1976. Sound production in mole crickets (Orthoptera: Gryllotalpidae: *Scapteriscus*). *Annals of the Entomological Society of America* 69: 299–306.
- Ulagaraj SM, Walker TJ. 1973. Phonotaxis of crickets in flight: attraction of male and female crickets to male calling songs. *Science* 182: 1278–1279.
- Ulagaraj SM, Walker TJ. 1975. Response of flying mole crickets to three parameters of synthetic songs broadcast outdoors. *Nature* 253: 530–532.
- Walker TJ. 1982. Sound traps for sampling mole cricket flights (Orthoptera: Gryllotalpidae: *Scapteriscus*). *Florida Entomologist* 65: 105–110.

- Walker TJ. 1988. Acoustic traps for agriculturally important insects. *Florida Entomologist* 71: 484–492.
- Walker TJ. 1996. Acoustic methods of monitoring and manipulating insect pests and their natural enemies, pp. 245–257 *In* Rosen D, Capinera JL, Bennett FD [eds.], *Pest Management in the Subtropics - Integrated Pest Management - A Florida Perspective*. Intercept, Andover, United Kingdom.
- Walker TJ, Nation JL. 1982. Sperm storage in mole crickets: fall matings fertilize spring eggs in *Scapteriscus acletus*. *Florida Entomologist* 65: 283–285.
- Walker TJ, Nickle DA. 1981. Introduction and spread of pest mole crickets: *Scapteriscus vicinus* and *S. acletus* reexamined. *Annals of the Entomological Society of America* 74: 158–163.
- Xu Y, Held DW, Hu XP. 2012. Potential negative effects of earthworm prey on damage to turfgrass by omnivorous mole crickets (Orthoptera: Gryllotalpidae). *Environmental Entomology* 41: 1139–1144.