



**First Record of *Erythmelus klopomor* (Hymenoptera: Mymaridae) as a Parasitoid of the Avocado Lace Bug, *Pseudacysta perseae* Heteroptera: Tingidae)**

Authors: Peña, J. E., Triapitsyn, S. V., Long, D., Evans, G. A., and Roltsch, W.

Source: Florida Entomologist, 92(2) : 394-395

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.092.0232>

---

BioOne Complete ([complete.BioOne.org](http://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

FIRST RECORD OF *ERYTHMELUS KLOPOMOR* (HYMENOPTERA: MYMARIDAE) AS A PARASITOID OF THE AVOCADO LACE BUG, *PSEUDACYSTA PERSEAE* (HETEROPTERA: TINGIDAE)

J. E. PEÑA<sup>1</sup>, S. V. TRIAPITSYN<sup>2</sup>, D. LONG<sup>1</sup>, G. A. EVANS<sup>3</sup> AND W. ROLTSCH<sup>4</sup>

<sup>1</sup>University of Florida, Entomology and Nematology Department, Tropical Research and Education Center, 18905 SW 280<sup>th</sup> Street, Homestead, FL 33031

<sup>2</sup>Department of Entomology, University of California, Riverside, CA 92521

<sup>3</sup>Systematic Entomology Laboratory, USDA/ARS/BARC-West, Building 005, Rm 09A, 10300 Baltimore Avenue, Beltsville, MD 20705

<sup>4</sup>Biological Control Program, Integrated Pest Control Branch, California Department of Food and Agriculture, 3288 Meadowview Road, Sacramento, CA 95832

The avocado lace bug, *Pseudacysta perseae* (Heidemann) (Hemiptera: Tingidae), is a secondary pest of avocado and other Lauraceae in Florida and other states within the continental USA (Peña et al. 1998; Wysoki et al. 2002; Hoddle et al. 2005). Due to a resurgence of *P. perseae* in Florida in the 1990s, a survey for natural enemies was conducted from 1995 to 1997 (Peña et al. 1998). During this survey, an unidentified mymarid (Hymenoptera: Mymaridae) was reared from parasitized eggs of *P. perseae* together with *Oligosita* sp. (Hymenoptera: Trichogrammatidae) (Identified by M. Schauff, USDA, Beltsville, MD). That study, however, did not include the extent of parasitization caused by both natural enemies. Eggs of *P. perseae* are deposited as clusters in the adaxial surface of avocado leaves, and they hatch within 9 to 19 d, depending on the temperature (Morales, 2000). At temperatures above 30°C, which are common in much of Florida, especially in the avocado growing season in summer. Therefore, any egg parasitoid must discover the *P. perseae* egg mass within the shorter part of the range. According to the literature, no other egg parasitoids have been reported from *P. perseae* (Aboud-Antun 1991; Medina-Gaud et al. 1991; Morales 2000; Sandoval 2004; Streito & Morival 2005).

An additional survey for parasitoids of *P. perseae* was initiated from Jan 2006 through Jun 2007. The survey consisted of monthly collection of 20 avocado leaves infested with *P. perseae*, from 3 sites within Miami-Dade County (USDA, Miami Research Station, latitude 25.64194 and longitude -80.29038; Princeton, FL, latitude 25°30' 49.99N and longitude 80°26' 18.71W, and the University of Florida, Tropical Research and Education Center, Homestead, FL, latitude 25°30' 37.64N and longitude 80°30' 11.02W). All sites were avocado groves of different the West Indian cultivars Dunedin, Simmonds, Gottfried, and Booth, and all were insecticide free throughout the study. Leaves were placed in individual plastic bags in an ice chest and transferred to the laboratory. Eggs were counted, and the section of the

leaf containing them was placed inside individual test tubes (12 × 75 mm) and sealed with a Kim-wipes tissue. These were held for 30 d at 25°C and 75% RH and 12:12 L:D and then inspected for the presence of any parasitoid. Emerged parasitoids were placed in 5-mL vials with 75% ETOH.

*Pseudacysta perseae* egg densities increased from 48.5 ± 5.0 eggs per leaf during Jan 2006 to 95.3 ± 7.0 eggs per leaf during Apr and decreased thereafter. Egg densities during 2007 increased from 29.0 ± 0.1 in Jan to 37.7 ± 3.2 during Mar 2007. Two species of parasitoids emerged from *P. perseae* eggs: the mymarid, *Erythmelus (Erythmelus) klopomor* S. Triapitsyn (Triapitsyn et al. 2007) was the more common one, while probably a new species of Trichogrammatidae (J. D. Pinto, pers. comm.) was the less common (only 1 specimen emerging from 1 egg throughout the survey). This is the first record of *E. klopomor* parasitizing *P. perseae*. Previously, *E. klopomor* was known as a parasitoid of *Corythucha* spp. (Tingidae) eggs in the eastern USA (Triapitsyn et al. 2007). Evidence of parasitism of *P. perseae* was considered low because of our collecting technique. For instance, the highest number of *E. klopomor* parasitoids emerging from an egg mass was 0.36 ± 0.2, which translates as 1.71% parasitism per cluster of 20 eggs. These numbers however, can be misleading. For instance, since the parasitoid emerges through a circular hole (Fig. 1) in the operculum in the same manner as *P. perseae*, those collected eggs with a circular hole through the operculum could have been either parasitized or not. The importance of egg parasitism or other egg mortality factor needs to be further studied. For instance, during an evaluation of egg mortality factors in Homestead, during 2007, emergence of the tingid nymphs from eggs exposed to biotic and abiotic mortality factors was reduced 67% compared to a 20% reduction of nymphal emergence from eggs where these mortality factors were excluded (Peña et al. unpublished). Only these 2 parasitoids and the predator, *Paracarniella cubana* (Peña et al. 1998) have been observed as mortality factors for *P. perseae* eggs in Florida.



Fig. 1. *Erythmelus klopomor* emerging from eggs of *P. perseae*. Credits. D. Long.

#### SUMMARY

The mymarid, *Erythmelus* (*Erythmelus*) *klopomor* S. Triapitsyn was found as an egg parasitoid of the avocado lace bug, *Pseudacysta perseae* (Heteroptera: Tingidae) for the first time in Miami-Dade County, Florida, USA.

#### REFERENCES CITED

- ABUD-ANTUN, A. 1991. Presence of avocado lacebug *Pseudacysta perseae* (Heidemann) (Hemiptera: Tingidae) in Dominican Republic. Primera Jornada de Protección Vegetal, Universidad de Santo Domingo, República Dominicana (Abstract), p. 4.
- HODDLE, M., MORSE, J., STOUTHAMER, R., TRIAPITSYN, S., HUMERES, E., BENDER, G., KELLUM, D., DOWELL, R., ROLTSCH, W., PHILLIPS, P., AND WHITNEY, G. 2005. Biology and management of avocado lacebug in California. Semiannual Report, California Avocado Growers. Downloaded as: [www.avocado.org/growers/pdf/2005\\_midyear/HoddeLaceBugProgress05.pdf](http://www.avocado.org/growers/pdf/2005_midyear/HoddeLaceBugProgress05.pdf).
- MEAD, F., PEÑA, J. E. 1991. Avocado Lace Bug, *Pseudacysta perseae* (Hemiptera: Tingidae). Florida Dept. Agric. Cons. Serv., Div. Plant Ind., Entomol. Circ. 346.
- MEDINA-GAUD, S., SEGARRA, A., AND FRANQUI, R. 1991. The avocado lace bug, *Pseudacysta perseae* (Heidemann) (Hemiptera: Tingidae). J. Agric. Univ. Puerto Rico 75: 185-188.
- MORALES, L. 2000. Estudios bioecológicos de *Pseudacysta perseae* (Heid.) (Heteroptera: Tingidae). Tesis Presentada para aspirar al título de Master en Agricultura Sostenible. Mención Sanidad Vegetal. Facultad de Ciencias Agropecuarias. Universidad Central Marta Abreu de las Villas. Santa Clara. Villa Clara, Cuba.
- PEÑA, J. E., SUNDHARI, S., HUNSBERGER, A., DUNCAN, R., AND SCHAFFER, B. 1998. Monitoring, damage, natural enemies and control of the avocado lace bug, *Pseudacysta perseae* (Hemiptera: Tingidae). Proc. Florida State Hort. Soc. 111: 330-334.
- SANDOVAL, M. F. 2004. Presencia de *Pseudacysta perseae* (Heid.) (Insecta: Hemiptera: Tingidae) en Venezuela. En: Resúmenes V Seminario Científico Internacional de Sanidad Vegetal. 24-28 Mayo. La Habana, Cuba.
- STREITO, J. C., AND MORIVAL, T. 2005. Première capture en Guyane Française de *Pseudacysta perseae* (Heidemann, 1908), un ravageur de l'avocatier (Heteroptera: Tingidae). Nouv. Revue Entomol. 22: 191-192.
- TRIAPITSYN, S. V., BEREZOVSKIY, V. V., HODDLE, M. S., AND MORSE, J. G. 2007. A review of the Nearctic species of *Erythmelus* (Hymenoptera: Mymaridae), with a key and new additions to the New World fauna. Zootaxa 1641: 1-64.