

**Larinus minutus (Coleoptera: Curculionidae), A
Biological Control Agent of Spotted Knapweed
(Centaurea stoebe ssp. Micranthos), Established in
Northern Arkansas**

Authors: Minter, C. R., Kring, T. J., Shen, J., and Wiedenmann, R. N.

Source: Florida Entomologist, 94(2) : 350-351

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.094.0233>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

***LARINUS MINUTUS* (COLEOPTERA: CURCULIONIDAE), A BIOLOGICAL CONTROL AGENT OF SPOTTED KNAPWEED (*CENTAUREA STOEBE* SSP. *MICRANTHOS*), ESTABLISHED IN NORTHERN ARKANSAS**

C. R. MINTEER, T. J. KRING, J. SHEN AND R. N. WIEDENMANN

Department of Entomology, University of Arkansas, Fayetteville, AR 72701 USA

Larinus minutus Gyllenhal (Coleoptera: Curculionidae) is a univoltine weevil that feeds on the seeds of spotted and diffuse knapweeds *Centaurea stoebe* ssp. *micranthos* (Gugler) Hayek and *C. diffusa* Lamarck. After emerging from overwintering sites in the leaf litter, adult weevils begin feeding on the vegetative portions of the plants. Adults, however, prefer to feed on flowers when they are available and development of beetle ovaries is dependent upon flower feeding (Groppe 1990). Females oviposit on newly opened flower heads (capitula). Two or 3 eggs can occur in each flower head, but only 1 larva usually develops in smaller capitula. Multiple larvae can survive in large spotted knapweed capitula (Groppe 1990). Under laboratory conditions (25°C), eggs hatch in 3-4 d (Groppe 1990). Larval development takes approximately 4 weeks and larvae go through 3 instars. Larvae feed on knapweed seeds and pupate in the capitula, making a cocoon out of the seed head material (Kashefi & Sobhian 1998). Larvae can destroy up to 100% of the seeds in a capitulum (Kashefi & Sobhian 1998). In the Western United States, adult weevils emerge in late Sep and feed on plants until winter, when the adults overwinter in leaf litter and emerge in the following Jun (Jordan 1995).

Larinus minutus was first released into the United States in 1991 with collections from Greece and Romania (Story 2002). Although 12 other natural enemy species were introduced into the Western United States and Canada to control spotted and diffuse knapweeds, only recently has adequate suppression of some populations been seen (Myers 2004; Smith 2004). Myers (2007) suggested that knapweed populations did not significantly decline until the establishment of *L. minutus*. Populations of *L. minutus* have been established in Washington, Wyoming, Oregon, Montana, Minnesota, Colorado, and Indiana (Lang et al. 1996; Story 2002).

No natural enemies of spotted knapweed have been released in Arkansas until the inception of this study. However, we found *Urophora quadrfasciata* (Meigen) (Diptera: Tephritidae) established throughout the range of spotted knapweed in the state in a survey in 2006 for knapweed natural enemies. This seedhead galling fly was introduced from Russia into Canada in 1980 and has since been re-distributed or spread on its own to several states in the northeastern and northwestern United States (Story 2002). Duguma (2008) found that *U. quadrfasciata* reduced the number of seeds produced by

spotted knapweed by 44% late in the season (Aug), at a time when plants are more environmentally stressed. However, the fly did not significantly reduce the number of seeds produced earlier in the season, a time when knapweed is most robust (Duguma 2008). Thus, it is likely that *U. quadrfasciata* alone will not significantly suppress knapweed populations in Arkansas, or stop its spread further into the southern United States.

To increase the level of control beyond that provided by *U. quadrfasciata*, we have introduced *L. minutus* in Arkansas. Collections of *L. minutus* were made from diffuse knapweed infestations in Colorado Springs, CO, for the Arkansas releases. A total of 8 releases were made during 2008 and 2009 (Fig. 1). Two releases were made in 2008 with 400 weevils released at the University of Arkansas Agriculture Experiment Station Farm, and 300 weevils released approximately 5 km south of the station. After the releases, sites were visually surveyed for surviving weevils each week. Weevils were observed for the remainder of the growing season in 2008. Six additional releases were made in 2009 in Washington County, Arkansas, with 600 to 700 weevils per release. Release sites in 2009 were of sufficient distance away from the release sites in 2008 that we believe the weevils had not spread to the new release sites. After releases in 2009, all sites were visually surveyed for adult weevils for the remainder of the growing season in 2009. *Larinus minutus* were found at all release sites (2008 and 2009) except one—the 2008- site just south of the University of Arkansas Agriculture Experiment Station.




Fig. 1. Locations of releases of *Larinus minutus* in Washington County, Arkansas during 2008 and 2009.

Beginning in mid-Apr 2010, emerging adult weevils were collected weekly by sweep-netting at all Arkansas release sites, once plants had bolted, but before flowers were present. *Larinus minutus* were first found on May 5, 2010. Weekly sweeps yielded *L. minutus* at each of the 2008- and 2009-release sites, including the 2008-site at which no weevils were found during 2009. We believe that low population levels the first year after release prohibited us from collecting weevils at the south Fayetteville site. Finding weevils at all sites demonstrated establishment at the 2008-release sites and successful overwintering of adults at the 2009-release sites. Thus, *L. minutus* joins *U. quadrifasciata* as an established natural enemy of spotted knapweed in northern Arkansas.

Additional releases in northern Arkansas of ~17,000 *L. minutus* were made in 2010 with weevils collected from Colorado Springs, CO.

We thank Jerry Michels and his crew from Texas A&M AgriLife for help in locating and collecting weevils in Colorado.

SUMMARY

Larinus minutus, a biological control agent of spotted knapweed, has been established at 2 sites in Washington County, Arkansas. The weevils were collected in Colorado and introduced into Arkansas at 2 sites in 2008 and 6 additional site in 2009. No *L. minutus* weevil was recorded in Arkansas prior to these releases.

REFERENCES CITED

- DUGUMA, D. 2008. Seasonal Dynamics and Impact of *Urophora quadrifasciata* (Meigen) (Tephritidae: Diptera) on Spotted Knapweed in the Arkansas Ozarks., M.S. University of Arkansas, Fayetteville, AR.
- GROPPE, K. 1990. *Larinus minutus* Gyll. (Coleoptera: Curculionidae), A Suitable Candidate for the Biological Control of Diffuse and Spotted Knapweed in North America. CABI Institute of Biological Control Report, European Station, Delémont, Switzerland, 30 p.
- JORDAN, K. 1995. Host specificity of *Larinus minutus* Gyll. (Col., Curculionidae), an agent introduced for the biological control of diffuse and spotted knapweed in North America. *J. Appl. Entomol.* 119: 689-693.
- KASHEFI, J. M., AND SOBHIAN, R. 1998. Notes on the biology of *Larinus minutus* Gyllenhal (Col., Curculionidae), an agent for biological control of diffuse and spotted knapweeds. *J. Appl. Entomol.* 122: 547-549.
- LANG, R. F., STORY, J. M., AND PIPER, G. L. 1996. Establishment of *Larinus minutus* Gyllenhal (Coleoptera: Curculionidae) for biological control of diffuse and spotted knapweed in the western United States. *Pan-Pacific Entomol.* 72: 209-212.
- MYERS, J. H. 2004. A silver bullet in the biological control of diffuse knapweed. Entomological Society of America 2004 Annual meeting abstract. <http://abstracts.co.allenpress.com/pweb/esa2004/document/36421>.
- MYERS, J. H. 2007. How many and what kind of agents for the biological control of weeds: a case study with diffuse knapweed, pp. 70-79 *In* C. Vincent, M. S. Goettel, and G. Lazarovitis [eds.], *Biological Control: A Global Perspective*. CAB International. Trowbridge, UK, 79 pp.
- SMITH, L. 2004. Impact of biological control agents on *Centaurea diffusa* (diffuse knapweed) in central Montana, pp. 589-593 *In* J. M. Cullen, D. T. Brieske, D. J. Kriticos, W. M. Lonsdale, L. Morin and J. K. Scott [eds.], *Proc. XI Intl. Symp. on Biol. Control of Weeds*, April 27-May 2 2003, Canberra, Australia.
- STORY, J. M. 2002. Spotted knapweed, pp. 169-180 *In* R. G. Van Driesche, B. Blossey, M. Hoddle, S. Lyon and R. Reardon [eds.], *Biological Control of Invasive Plants in the Eastern United States*. USDA Forest Service, Morgantown, West Virginia.