

## **Testes and Spermatozoa as Characters for Distinguishing Two Ant Species of the Genus *Neoponera* (Hymenoptera: Formicidae)**

Authors: Barcellos, Marcelo S., Martins, Luiza C. B., Cossolin, Jamile F. S., Serrão, José Eduardo, Delabie, Jacques H. C., et al.

Source: Florida Entomologist, 98(4) : 1254-1256

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.098.0441>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# Testes and spermatozoa as characters for distinguishing two ant species of the genus *Neoponera* (Hymenoptera: Formicidae)

Marcelo S. Barcellos<sup>1</sup>, Luiza C. B. Martins<sup>2</sup>, Jamile F. S. Cossolin<sup>1</sup>, José Eduardo Serrão<sup>1</sup>, Jacques H. C. Delabie<sup>3</sup>, and José Lino-Neto<sup>1,\*</sup>

The correct diagnosis of a species is often challenging, especially in the case of cryptic species. The misidentification can invalidate results already published, contribute negatively to the knowledge of the organism in question, and waste financial resources associated with research and control programs. Therefore, it is important to find easily visible traits that allow the diagnosis of the species under study. The external morphologies of the ant species *Neoponera inversa* (Smith) and *Neoponera villosa* (F.) (Hymenoptera: Formicidae) are very similar, and as a result, they were previously considered as the single species *N. villosa*. However, studies comparing petiole morphology and isozyme patterns (Lucas et al. 2002), chromosome number (Mariano et al. 2007), and external morphologies (Fernandes et al. 2014) revealed that they form, in fact, a complex of sympatric cryptic species. These characteristics make the distinction difficult between these species. Thus, to identify them requires a specialist and, in general, methods that are time consuming and expensive.

Morphological characteristics of the male reproductive system have been used in the systematics of insects particularly in Hymenoptera (Wheeler & Krutzsch 1992; Dias et al. 2013). In addition, spermatozoa have been used for phylogenetic analyses to determine taxonomic variations in insects (Jamieson et al. 1999; Dallai 2014). In Hymenoptera, the sperm may be free in the seminal vesicle, as reported for parasitic wasps (Lino-Neto et al. 1999) and most Aculeata (Moreira et al. 2004), or arranged in bundles as described for ants such as *Crematogaster victima* Smith (Oliveira et al. 2014) and *Lasius pallitarsis* (Provancher) (Burnett & Heinze 2014) and for sawflies (Schiff et al. 2001).

Morphological characteristics of testes and sperm may be useful in the differentiation of these sympatric, cryptic species of ants. In this study, we describe the numbers of follicles per testis and the morphometries of the spermatozoa of *N. inversa* and *N. villosa* to establish additional characters by which they can easily be distinguished.

Five colonies of *N. inversa* and 5 colonies of *N. villosa* were collected in Ilhéus, state of Bahia, Brazil, and reared for 3 mo within artificial nests in the laboratory. These nests were kept at  $25 \pm 4$  °C with about 50 to 80% relative humidity. All colonies were fed on a mixture of honey, apple, and *Tenebrio molitor* L. (Coleoptera: Tenebrionidae) larvae ad libitum. Our studies used in total 50 males consisting of 5 males from each colony.

To quantify the number of follicles per testis, 15 males of *N. inversa* and 15 of *N. villosa* (3 specimens per colony) were dissected, and the

reproductive system was photographed with a stereoscopic microscope (Zeiss, Stemi 2000-C). For the morphometric analysis of sperm, the seminal vesicles of 2 males from each colony (totaling 10 specimens per species) were transferred to histology slides, dissected in pH 7.2 phosphate buffered saline (PBS), fixed in 4% paraformaldehyde in 0.1 M phosphate buffer for 20 min, and washed in distilled water for 20 min. The spermatozoa were observed and photographed under a light microscope (Olympus, BX-60). To visualize and measure the nuclei, some slides were stained for 15 min with 4,6-diamidin-2-phenylindole (DAPI) at a concentration of 0.2 mg/mL in PBS and observed under an epifluorescence microscope (Olympus, BX-60) equipped with a BP 360–370 nm filter. For the measurements, 50 nuclei and 50 flagella of each species (10 cells per colony of both species) were randomly measured using the program Image-Pro Plus Version 4.5. The mean values were estimated and compared using the Mann–Whitney test at 5% significance.

Results showed that in the 2 species *N. inversa* and *N. villosa*, the reproductive system consisted of a pair of testes surrounded by a thin peritoneal sheath that can be broken easily (Fig. 1A, B). Each testis opened into a long and thin vas deferens, forming the seminal vesicle, and both seminal vesicles and paired accessory glands terminated into a short ejaculatory duct (Fig. 1A, B). When the peritoneal sheath was broken, 3 follicles per testis were observed in *N. inversa* (Fig. 1C), whereas 4 follicles per testis were observed in *N. villosa* (Fig. 1D).

In the seminal vesicle, the spermatozoa were individualized, long, and threadlike. The total lengths of the sperm of *N. inversa* and *N. villosa* were  $155 \pm 10.1$  µm and  $175 \pm 5.4$  µm, respectively (Fig. 2A, B). Lengths of flagella did not differ between the 2 species ( $P = 0.3734$ ) and averaged 125 µm. However, the nucleus of *N. inversa* sperm was smaller than the nucleus of *N. villosa* ( $P < 0.001$ ), with a length of  $30 \pm 1.6$  µm and  $50 \pm 2.3$  µm, respectively (Fig. 2C, E).


The different numbers of follicles per testis in *N. inversa* (3) and *N. villosa* (4) support the proposition that they are distinct species within the genus *Neoponera* (Lucas et al. 2002) and facilitate greatly the distinction between them. The numbers of testicular follicles vary in different groups within Hymenoptera. Three follicles per testis have been observed in Vespidae (Brito et al. 2005) and Andrenidae, Halictidae, and some species of Megachilidae (Ferreira et al. 2004). Four follicles per testis have been observed in Mellitidae, other species of Megachilidae, and Apidae sensu stricto (Roig-Alsina & Michener 1993;

<sup>1</sup>Departamento de Biologia Geral, Universidade Federal de Viçosa, 36570-900, Viçosa, Minas Gerais, Brazil

<sup>2</sup>Departamento de Química e Biologia, Universidade Estadual do Maranhão—UEMA, 65604-375, Caxias, Maranhão, Brazil

<sup>3</sup>Laboratório de Mirmecologia, Convênio CEPLAC/UESC, CEPEC - 45600-000, Itabuna, Bahia, Brazil

\*Corresponding author; E-mail: lino-neto@ufv.br




**Fig. 1.** Male reproductive system of (A) *Neoponera inversa* and (B) *Neoponera villosa*, showing the testes (t) united by peritoneal sheath. Seminal vesicle (sv); vas deferens (vd); accessory glands (ag). Testes of (C) *N. inversa* and (D) *N. villosa* without peritoneal sheath, showing 3 and 4 follicles (arrows), respectively.

Ferreira et al. 2004). Interestingly, the number of testicular follicles can be up to 250 in *Apis mellifera* L. (Hymenoptera: Apidae) (Snodgrass 1984), whereas in ants, the number varies from 1 to 25 among different species (Wheeler & Krutzsch 1992).

In *N. inversa* and *N. villosa*, the spermatozoa were long, threadlike, linear, and free from each other in the seminal vesicle as also observed in the ant *Acromyrmex subterraneus* (Forel) (Hymenoptera: Formicidae) (Moreira et al. 2004). However, in *C. victima* (Oliveira et al. 2014) and *L. pallitarsis* (Burnett & Heinze 2014), these cells are maintained in bundles in the seminal vesicle in the mature male as reported for sawfly wasps (Schiff et al. 2001).

In Hymenoptera, the difference in lengths of the sperm can be used to differentiate cryptic species (Pereira et al. 2008). Total lengths of sperm are variable among species, for example, 13 to 577  $\mu\text{m}$  in Vespidae (Quicke et al. 1992) and 80 to 1,500  $\mu\text{m}$  in Apoidea (Quicke et al. 1992; Zama et al. 2004). In Formicidae, the mean length of the sperm in *Solenopsis invicta* Buren is about 70  $\mu\text{m}$  (Lino-Neto & Dolder 2002), whereas in *Pseudomyrmex*, it ranges from 53 to 70  $\mu\text{m}$  (Moya et al. 2007). Thus, our results showed that sperm lengths in *N. inversa* (155  $\mu\text{m}$ ) and *N. villosa* (175  $\mu\text{m}$ ) are the longest recorded to date in ants. Furthermore, we showed that the difference in the size of the sperm between *N. inversa* and *N. villosa* is due to a difference in the lengths of nuclei, with the nucleus being 40% longer in *N. villosa*. This



**Fig. 2.** Sperm of (A) *Neoponera inversa* and (B) *Neoponera villosa* in phase contrast, showing the transition (arrows) between the nucleus (n) and the flagellum (f). Nuclei stained with DAPI of (C and D) *N. inversa* and (E and F) *N. villosa* sperm.

result is consistent with a cytogenetic study on differences in chromosome number between these 2 species, with *N. inversa* showing fewer chromosomes ( $2n = 30$ ) than *N. villosa* ( $2n = 34$ ) (Mariano et al. 2007). Considering the importance of these reproductive cells, and that the amount of DNA in the same species is generally constant, the difference in nuclear size of the sperm of these 2 ant taxa supports that they are different species.

Although all colonies have been collected in the same region (Ilhéus, Bahia State, Brazil), the way the data were sampled and their uniformity, as well as the sample size, provide strong evidence that the number of follicles per testis and the nuclear length of sperm are species specific. Therefore, the characters cited above can be used to distinguish easily these 2 cryptic species of ants.

We are grateful to the Brazilian research agencies FAPEMIG, CNPq, PRONEX SECTI-FAPESB/CNPq - PNX 0011/2009, to José Adade, José Raimundo Maia, and José Crispin (Laboratory of Myrmecology at the Cocoa Research Center, Ilhéus, Bahia, Brazil) and to the 2 reviewers who greatly improved the manuscript with their suggestions.

## Summary

We examined the morphologies of the testes and spermatozoa of the ant species *Neoponera inversa* (Smith) and *Neoponera villosa* (F.) (Hymenoptera: Formicidae), collected in Ilhéus, Bahia State, Brazil, looking for information to distinguish easily one species from another. In *N. inversa*, 3 follicles per testis occurred, and the nucleus of sperm measured  $30 \pm 1.6 \mu\text{m}$  in length. In *N. villosa*, 4 follicles per testis occurred, and the nucleus of sperm measured  $50 \pm 2.3 \mu\text{m}$  in length. The morphologies of the testes and the spermatozoa offer an easy and fast method to distinguish between the 2 sympatric, cryptic species *N. inversa* and *N. villosa*.

**Key Words:** *Neoponera inversa*; *Neoponera villosa*; testicular follicles; sperm morphometry

## Sumário

---

Nós analisamos a morfologia dos testículos e dos espermatozoides das formigas *Neoponera inversa* (Smith) and *Neoponera villosa* (F.) (Hymenoptera: Formicidae), coletadas em Ilhéus, Bahia, Brasil, buscando por informações que possam facilmente distinguir uma espécie da outra. Em *N. inversa* ocorre 3 folículos por testículo e o núcleo dos espermatozoides medem  $30 \pm 1,6 \mu\text{m}$  de comprimento, já em *N. villosa* são 4 folículos e o núcleo dos espermatozoides medem  $50 \pm 2,3 \mu\text{m}$ . Portanto, a morfologia dos testículos e dos espermatozoides oferece um método fácil e rápido para distinguir *N. inversa* de *N. villosa*, duas espécies crípticas simpátricas.

Palavras Chave: *Neoponera inversa*; *Neoponera villosa*; folículos testiculares; morfometria espermática

## References Cited

---

- Brito P, Moreira J, Lino-Neto J. 2005. Morphology of male reproductive system and sperm of *Mischocyttarus* sp. (Hymenoptera: Vespidae: Polistinae). Brazilian Journal of Morphological Sciences 22: 175-176.
- Burnett WE, Heinze J. 2014. Sperm bundles in the seminal vesicles of sexually mature *Lasius* ant males. PLoS One 9: e93383.
- Dallai R. 2014. Overview on spermatogenesis and sperm structure of Hexapoda. Arthropod Structure and Development 43: 257-290.
- Dias G, Oliveira CM, Lino-Neto J. 2013. Testicular and spermatogenic characteristics of *Lagria villosa* (Tenebrionidae: Lagriinae) with taxonomic inferences. Tissue and Cell 45: 227-230.
- Fernandes IO, Oliveira ML, Delabie JHC. 2014. Description of two new species in the Neotropical *Pachycondyla foetida* complex (Hymenoptera: Formicidae: Ponerinae) and taxonomic notes on the genus. Myrmecological News 19: 133-163.
- Ferreira A, Abdalla FC, Kerr WE, Cruz-Landim C. 2004. Comparative anatomy of the male reproductive internal organs of 51 species of bees. Neotropical Entomology 33: 569-576.
- Jamieson BGM, Dallai R, Afzelius BA. 1999. Insects: Their Spermatozoa and Phylogeny, 1st edition. Science Publishers Inc., Enfield, New Hampshire, USA.
- Lino-Neto J, Dolder H. 2002. Sperm structure and ultrastructure of the fire ant *Solenopsis invicta* (Buren) (Hymenoptera, Formicidae). Tissue and Cell 34: 124-128.
- Lino-Neto J, Bão SN, Dolder H. 1999. Structure and ultrastructure of the spermatozoa of *Bephratelloides pomorum* (Fabricius) (Hymenoptera: Eurytomidae). International Journal of Insect Morphology and Embryology 28: 253-259.
- Lucas C, Fresneau D, Kolmer K, Heinze J, Delabie JHC, Pho DB. 2002. A multidisciplinary approach to discriminating different taxa in the species complex *Pachycondyla villosa* (Formicidae). Biological Journal of the Linnean Society 75: 249-259.
- Mariano CSF, Delabie JHC, Santos JRM, Pompolo SG. 2007. Evolução cariotípica em *Pachycondyla* spp. (Ponerinae) neotropicais. Biológico 69: 409-412.
- Moreira J, Zama U, Lino-Neto J. 2004. Release, behavior and phylogenetic significance of spermatozoa in bundles in the seminal vesicle during sexual maturation in Aculeata (Hymenoptera). Brazilian Journal of Morphological Sciences 21: 185-189.
- Moya J, Mancini K, Lino-Neto J, Delabie J, Dolder H. 2007. Sperm ultrastructure of five species of the Neotropical ant genus *Pseudomyrmex* (Hymenoptera: Formicidae). Acta Zoologica 88: 181-187.
- Oliveira CM, Moreira J, Gomes LF, Camargo-Mathias MI, Lino-Neto J. 2014. Sperm bundles in the seminal vesicle of the *Crematogaster victima* (Smith) adult males (Hymenoptera: Formicidae). Neotropical Entomology 43: 201-208.
- Pereira RAS, Lino-Neto J, Prado AP. 2008. Sperm morphology: a novel way to associate female-males of highly sexual dimorphic fig wasp species. Symbiosis 45: 143-147.
- Quicke DLJ, Ingram SN, Baillie HS, Gaitens PV. 1992. Sperm structure and ultrastructure in the Hymenoptera (Insecta). Zoologica Scripta 21: 381-402.
- Roig-Alsina A, Michener CD. 1993. Studies of the phylogeny and classification of long-tong bees (Hymenoptera: Apoidea). University of Kansas Science Bulletin 55: 123-162.
- Schiff N, Flemmenc AJ, Quicke DLJ. 2001. Spermatodesmata of the sawflies (Hymenoptera: Symphyta): evidence for multiple increases in sperm bundle size. Journal of Hymenoptera Research 10: 119-125.
- Snodgrass RE. 1984. Anatomy of the Honey Bee, 1st edition. Cornell University Press, Ithaca, New York, USA.
- Wheeler DE, Krutzsch PH. 1992. Internal reproductive system in adult males of the genus *Camponotus* (Hymenoptera: Formicidae: Formicinae). Journal of Morphology 211: 307-317.
- Zama U, Lino-Neto J, Dolder H. 2004. Structure and ultrastructure of spermatozoa in Meliponini (stingless bees) (Hymenoptera: Apidae). Tissue and Cell 36: 29-41.