



TRICHOMONIASIS IN A BONELLI'S EAGLE POPULATION IN SPAIN

Authors: Real, Joan, Mañosa, Santi, and Muñoz, Elena

Source: Journal of Wildlife Diseases, 36(1) : 64-70

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-36.1.64>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

TRICHOMONIASIS IN A BONELLI'S EAGLE POPULATION IN SPAIN

Joan Real,^{1,3} Santi Mañosa,¹ and Elena Muñoz²

¹ Departament de Biologia Animal (Vertebrats), Facultat de Biologia, Universitat de Barcelona, Diagonal, 645. 08028 Barcelona, Catalonia, Spain

² Departament de Patologia i Producció Animals, Facultat de Veterinària, Universitat Autònoma de Barcelona, 08193 Bellaterra, Catalonia, Spain

³ Corresponding author (e-mail: jreal@porthos.bio.ub.es)

ABSTRACT: During 1980–97, trichomoniasis was detected in nestlings of Bonelli's eagle *Hieraetus fasciatus* in Catalonia (Spain). In 1993 *Trichomonas gallinae* was isolated in 36% of nestlings ($n = 39$) and affected 41% of broods ($n = 22$). Overall, trichomoniasis was one of the most important single nestling mortality factor, accounting for 22% of total chick mortality, and causing the death of 2% of chicks. Trichomoniasis deaths took place during the second half of the nestling period. The median age at death was 45.5 days. Although the presence of the parasite was not related to the composition of the diet or parental age, pairs that developed the disease ate more pigeons and included more often non-adult birds. At present trichomoniasis apparently has little demographic impact on the Bonelli's eagle population in Catalonia, but the eventual spread of this disease in chicks and its unknown effects on adults might be of concern.

Key words: Birds of prey, Bonelli's eagle, *Hieraetus fasciatus*, nestlings, survey, *Trichomonas gallinae*, trichomoniasis.

INTRODUCTION

Feral pigeons (*Columba livia*) and other columbiform birds are the main primary hosts of *Trichomonas gallinae*, a sarcomastigophoran parasite that causes trichomoniasis in birds (Stabler, 1951; Locke and James, 1962). Most of the strains of the parasite are found in the upper digestive tract of their hosts and are considered to be non-pathogenic (Stabler, 1969; Honigberg, 1970). However, others are highly virulent and are responsible for proliferative lesions in the oropharynx that cause the death of a non-immune bird by starvation or secondary bacterial infections (Stabler, 1954; Mesa et al., 1961; Cooper and Petty, 1988; Samour et al., 1995).

Trichomoniasis has been reported frequently in birds of prey from several areas (Stabler, 1969; Keymer, 1972; Beecham and Kochert, 1975; Rettig, 1978; Tangredi, 1978; Stone and Nye, 1981; Redig, 1987; Pepler and Oettlé, 1992; Samour et al., 1995; Boal et al., 1998), but there is little information on the prevalence of infection or disease incidence of this parasite in wild populations (Beecham and Kochert, 1975; Cooper and Petty, 1988; Kietzmann, 1988; Boal et al., 1998). The occurrence of trichomoniasis in this group of birds is re-

lated to the ingestion of infected pigeons (Stabler and Shelanski, 1936; Stabler, 1969; Boal et al., 1998) or other infected birds (Halliwell, 1979). In some species, there is increasing concern that the expansion of the disease in the population may impair or reduce population growth rate, or even put the population at risk (Cooper and Petty, 1988; Boal et al., 1998).

The Bonelli's eagle (*Hieraetus fasciatus*) is an endangered bird of prey whose populations have undergone a marked decline in Europe in recent years (Rocamora, 1994; Real and Mañosa, 1997). During the last decades, in some areas the eagles have increased the consumption of feral pigeons as a consequence of the decline on wild prey and the increase of feral pigeon populations (Real, 1991; Del Hoyo et al., 1994). This change increases the risk of exposure to *T. gallinae* and nestling mortality, which has been shown to entail the decline of some populations (Cugnasse, 1989; Real 1991; Fernández et al., 1998). The objectives of this paper are to describe some cases of trichomoniasis in wild Bonelli's eagle nestlings, to quantify the prevalence of infection of *T. gallinae* among nestlings, and to analyze the influence of diet and eagle age on the presence

TABLE 1. Cases of trichomoniasis detected in Bonelli's eagle chicks in Catalonia (Spain) during 1980–97.

Case	Year	Sex	Hatching order	Weight (g)	Age (days)	Status at finding ^a	Fate ^b	Number of siblings and fate
1	1987	F	1	1,150 ^d	58	D	D	1 (dead at postnestling ^c)
2	1992	M	2	1,010 ^d	47	A	D	1 (fledged)
3	1992	M	—	1,350 ^d	48	A	D	0
4	1993	F	—	—	28	D	D	1 (case 5)
5	1993	M	—	1,480	43	A	F	1 (case 4)
6	1995	F	1	—	44	D	D	1 (dead at 32 days ^c)
7	1997	M	—	1,690	43	D	D	0

^a D = dead, A = alive.

^b D = dead, F = fledged.

^c Unknown cause of death.

^d Underweight.

of the parasite and on the development of the disease. The potential impact of the disease in the Bonelli's eagle breeding success and conservation also is discussed.

MATERIALS AND METHODS

Between 1980–97 the breeding success of the Bonelli's eagle population was monitored in Catalonia (northeastern Spain), where about 70 pairs still remain (Real et al., 1997). A preliminary monitoring scheme was conducted during 1980–85 involving only six pairs to determine nestling mortality and breeding success. From 1986–93, an intensive study was undertaken on a large sample. During these years, the nests of 41 breeding sites were regularly checked for hatching success and nestling survival. Nests were visited 20, 35, 55, and 65 days after hatching to determine the levels and causes of nestling mortality during the nestling period. When possible, the age and sex of the nestlings were determined following Mañosa et al. (1995).

Clinical trichomoniasis in nestlings (living or dead) was verified by visual inspection of the buccal cavity, where characteristic *T. gallinae* lesions can be observed (Cooper, 1978; Ward, 1986; Cooper and Petty, 1988).

The prevalence of infection by *T. gallinae* was assessed in nestlings aged 32 to 46 days from 22 nests in 1993. Culture specimens were obtained by swabbing the surface areas of the upper crop, pharynx, palatin region, and mouth with a cotton-tipped swab previously soaked in culture medium. The samples were immediately transferred to a 5 ml of CPLM (Cysteine, Peptone, Liver extract and Maltose) *Trichomonas* broth with CAF (Chloramphenicol) (Bioline, le Monza, Milano, Italy) medium, (supplemented with 0.1 g mycostatin, 0.061 g penicillin, 0.1 g streptomycin and 5 ml of inactivated rabbit serum/100 ml). The cultures

were stored at room temperature and taken to the laboratory the same day, where they were incubated at 36 C. Each sample was examined under an inverted microscope (100×) after 24 to 48 hrs of incubation and checked again after 120 to 144 hrs. If no trichomonads were detected at this time, the sample was considered free of *T. gallinae*.

The diet of nestlings was assessed by collecting prey remains and pellets in the nest, which were analysed and quantified following Real (1996). Breeding birds were classified as adults (those with full adult plumage, >4-yr-old) or non-adults (≤4-yr-old) based on plumage characteristics (Parellada, 1984; J. Real, pers. data). We used chi-square and Fisher's exact tests for comparisons (Zar, 1984). The level of α was established at $P \leq 0.05$.

RESULTS

Between 1980–97, seven cases of trichomoniasis were recorded (Table 1, 2). Of those cases, three involved females and four involved males. Of these nestlings, six died before leaving the nest, and the remaining fledged.

The nestling period of Bonelli's eagle lasts for 65 to 70 days. Most deaths (five of six) associated with trichomoniasis occurred in the second half of this period, when nestlings were between 28-days and 58-days-old, usually after 40 days of life. The median age at death was 45.5 days ($n = 6$). Three of five nestlings were underweight (Mañosa et al., 1995) (Table 1, 2). All the nestlings had buccal cavity nodular lesions of different size, usually more than 2 cm long (Table 2). These nodules caused

TABLE 2. Characteristics of Bonelli's eagle chicks affected by trichomoniasis during 1980–97 in Catalonia (Spain).

Case	Nodule size	Nodule position in buccal cavity	Esophagus obstruction	External swelling	Weight loss	Rotted food into the mouth	Deteriorated plumage
1	>3 cm	Lower jaw and laterally	Total	Yes	Yes	Yes	No
2	3 cm	Lower jaw and laterally	Important	Yes	Yes	No	No
3	>3 cm	Superior jaw and posterior	Important	No	Yes	No	Yes
4	>3 cm	Lower jaw	Total	Yes	?	—	No
5	2 cm	Lower jaw and posterior	Little	No	No	—	No
6	2 cm 3 nod.	Lower jaw and posterior	Important	Yes	?	No	No
7	2 cm	Lower jaw near glottis	No	No	No	No	No

TABLE 3. Causes of nestling Bonelli's eagle mortality in relation to age during 1986–93.

Causes	1–20 days	21–40 days	41–60 days	Total chicks
Trichomoniasis	0	1	3	4
Starvation	1	0	0	1
Hatching asynchrony	4	0	0	4
Predation	1	0	0	1
Accident	0	0	2	2
Other diseases	0	1	0	1
Unknown	2	1	2	5
Total	8	3	7	18

moderate to severe occlusion of the esophagus and prevented the passage of food. One nestling may have died by asphyxia caused by a large nodule on the inferior surface of the tongue, which occluded the glottis. This chick was in good physical condition and the crop and stomach were full of recently ingested food. One of the underweight nestling showed deficient feather growth (remiges, tail feathers and coverts), consisting in bands without cornial deposits as a result of malnutrition (Grubb, 1995). The other underweight nestlings did not have deteriorated plumage but one had delayed development of the feathers.

The presence of *T. gallinae* was checked in 22 broods in 1993; one had 3 chicks, 15 had 2, and 6 consisted of one, involving 21 males and 18 females. The parasite was isolated in 13 live nestlings and diagnosed in one dead nestling (36%, 7 males and 7 females), corresponding to 9 broods (41%) (two single broods and seven double broods). Only two (5%) of these 14 infected chicks had lesions clearly related to

the disease and only one (3%) died. For the 105 broods checked in the 1986–93 intensive study period, involving 179 nestlings, trichomoniasis caused the death of 2% of chicks, accounting for 22% of total nestling mortality (Table 3). Mortality affected 0% of chicks and broods in 1986 ($n = 8$ chicks, and $n = 5$ broods), 1988 ($n = 18$, and $n = 10$), 1989 ($n = 14$, and $n = 8$), 1990 ($n = 14$, $n = 9$), and 1991 ($n = 13$, $n = 8$) respectively, 2% of chicks ($n = 54$) and 3% of broods ($n = 30$) in 1993, 5% of chicks ($n = 41$) and 8% of broods ($n = 25$) in 1992, and 6% of chicks ($n = 17$) and 10% of broods ($n = 10$) in 1987.

There were no differences in diet composition between broods that were positive or negative for *T. gallinae* in 1993 ($\chi^2 = 9.3955$, $df = 6$, $P = 0.1525$) (Table 4). However, the development of the disease was related to the consumption of pigeons. Taking into account only the four pairs that developed the disease at least once between 1980 to 1997, and where dietary information could be collected, differences in the consumption of *C. livia* were ob-

TABLE 4. Diet of pairs of Bonelli's eagle monitored in 1993 related to the presence of *Trichomonas gallinae*.

	Infected		Uninfected	
	<i>n</i>	%	<i>n</i>	%
European rabbit (<i>Oryctolagus cuniculus</i>)	39	28	57	25
Other mammals	26	18	23	10
Domestic pigeon (<i>Columba livia</i>)	15	11	22	9
Woodpigeon (<i>Columba palumbus</i>)	17	12	34	15
Phasianidae	17	12	32	14
Other birds	16	11	45	19
Ocellated lizard (<i>Lacerta lepida</i>)	12	8	19	8

served between three broods developing the disease (31%, $n = 26$) and four broods without the disease (11%, $n = 73$) ($\chi^2 = 4.187$, $df = 1$, $P = 0.04$). No differences were detected in the consumption of *C. palumbus*, 4% and 10% respectively ($\chi^2 = 0.254$, $df = 1$, $P = 0.61$).

In 1993 no difference of infection was found between adult or non-adult pairs (formed at least by one non-adult individual) (40%, $n = 15$ and 43%, $n = 7$, respectively, Fisher Exact Test, $P > 0.05$). However, when we take into account only the five pairs in which trichomoniasis developed at any time from 1980–97, we found that the disease developed more often when the breeding attempt involved a non-adult bird (83% $n = 6$), than when both parents were adult (5%, $n = 20$, Fisher Exact Test, $P = 0.0005$).

DISCUSSION

Despite the high prevalence of *T. gallinae* in nestling Bonelli's eagles, only a small proportion developed clinical trichomoniasis and died during the nestling period. This may indicate that most strains of *T. gallinae* are non-pathogenic or that nestlings of Bonelli's eagle have certain immunity, as happens in pigeons (Stabler, 1969; Honigberg, 1970) or other raptors (Samour et al. 1995; Boal et al. 1998). In spite of that, trichomoniasis appears as one of the most important causes of chick death and might produce up to 6% nestling mortality in particular years.

Although the presence of the parasite in a brood seems to be independent of the proportion of domestic pigeons or woodpigeons in the diet or parental age, the development of the disease in infected chicks is more frequent in broods consuming more domestic pigeons or where non-adult breeders are involved. This suggests that although most of the population is exposed to the protozoan (since most pairs consume pigeons to some extent), pairs consuming large amounts of domestic pigeons have a higher probability of being exposed to a virulent strain (Boal et al.,

1998). Also, the disease would mainly develop in broods raised by non-adult birds, which might be related to lower experience (Newton, 1979), resulting in poor chick condition, or to non-adult birds consuming more feral pigeons than adult birds.

Although the monitoring of Bonelli's eagle in Catalonia began in 1980, trichomoniasis was not detected until 1987, which might be related to the changes in feeding habits of Bonelli's eagles which took place in late 1980s (Real, 1991). Whatever the causes of these changes are (decline in wild prey populations, increased feral pigeon availability, increased proportion of non-experienced parents), increased consumption of feral pigeons may have led to an increased prevalence of the disease and chick mortality, since the prevalence of the parasite in rural pigeons *C. livia* from the area (90 to 97%) is far higher than that exhibited by woodpigeons *C. palumbus* (50%) (Muñoz, 1995). Increased trichomoniasis prevalences in raptor populations consuming large proportions of columbiform birds have repeatedly been reported (Cooper and Petty, 1988; Boal et al., 1998).

Although chick mortality associated with *T. gallinae* in Catalonia is low at present and may have low demographic effects (Real and Mañosa, 1997), it may be of future concern if natural prey continue to be replaced by infected prey such as domestic pigeons in eagle's diet. In Portugal, where Bonelli's eagle consume large amounts of domestic pigeons (Palma et al. 1984), nestling mortality as high as 14% associated with the disease has been reported (L. Palma, pers. comm.).

Future monitoring and research is needed to understand the susceptibility of adult birds to trichomoniasis, as well as the ecological factors and kind of prey related to the spread of the disease. In the mean time, the most sensible measure to be undertaken must be the restoration of wild prey populations (rabbits and red-legged partridges), in order to ameliorate the

availability of food to the eagles and to reduce the risk of exposure to the parasite. Further studies should indicate the eventual need of feral pigeon control and prophylaxis in some areas where prevalence of the disease is high.

ACKNOWLEDGMENTS

We are grateful to J. Codina, D. Molina and R. del Amo for helping in the field. Four anonymous reviewers gave very useful comments that improved the manuscript. Field work was carried out with the financial support of Fundació Miquel Torres of Vilafranca del Penedès (Catalonia, Spain).

LITERATURE CITED

- BEECHAM, J. J., AND M. N. KOCHERT. 1975. Breeding biology of the golden eagle in southwestern Idaho. *The Wilson Bulletin* 87: 506–513.
- BOAL, C. W., R. W. MANNAN, AND K. S. HUDELSON. 1998. Trichomoniasis in Cooper's Hawk from Arizona. *Journal of Wildlife Diseases* 34: 590–593.
- COOPER, J. E. 1978. Parasites. *In* Veterinary aspects of captive birds of prey, J. E. Cooper (ed.). The Standfast Press, Cherington, UK, pp. 82–96.
- , AND PETTY, S. J. 1988. Trichomoniasis in free living goshawks (*Accipiter gentilis*) from Great Britain. *Journal of Wildlife Diseases* 24: 80–87.
- CUGNASSE, J. M. 1989. Schutzstrategien für den Habichtsadler (*Hieraetus fasciatus*) im Französischen Languedoc-Roussillon. *Laufener Seminarbeiträge*, 1: 65–66.
- DEL HOYO, J., A. ELLIOTT, AND J. SARGATAL. 1994. Handbook of the birds of the world, Vol. 2. Lynx Edicions, Barcelona, Spain, 638 pp.
- FERNANDEZ, A., J. ROMAN, J. A. DE LA TORRE, L. M. ANSOLA, J. SANTAMARIA, R. VENTOSA, F. ROMAN, AND C. PALMA. 1998. Demografía y conservación de una población de águilas perdicera *Hieraetus fasciatus* en declive. *In* Holarctic birds of prey, Proceedings of an International Conference, B. U. Meyburg, R. D. Chancellor and J. J. Ferrero (eds.). Adenex-WWGBP, Badajoz, Spain, pp. 305–321.
- GRUBB, T. C. 1995. Ptilochronology. A review and prospectus. *In* Current ornithology, Vol. 12. D. M. Power (ed.). Plenum Press, New York, New York, pp. 89–114.
- HALLIWELL, W. H. 1979. Diseases of birds of prey. *Veterinary Clinics of North America* 9: 541–568.
- HONIGBERG, B. M. 1970. Trichomonads. *In* Immunity to parasitic animals, Vol. 2. G. J. Jackson, R. Herman and I. Singer (eds.). Appleton-Century-Crofts, New York, New York, pp. 469–550.
- KEYMER, I. F. 1972. Diseases of birds of prey. *Veterinary Record* 90: 579–594.
- KIETZMANN, G. E. 1988. Host parasite interactions of *Trichomonas gallinae* (Rivolta, 1878). Ph.D. Thesis, Iowa State University, Ames, Iowa, 144 pp.
- LOCKE, L. N., AND P. JAMES. 1962. Trichomonad cancer in the Inca dove, *Scardafella inca* (Lesson). *The Journal of Parasitology* 48: 497.
- MAÑOSA, S., J. REAL, AND J. CODINA. 1995. Age estimation and growth patterns in nestling Bonelli's eagles. *Journal of Raptor Research* 29: 273–275.
- MESA, C. P., R. M. STABLER, AND M. BERTHRONG. 1961. Histopathological changes in the domestic pigeon infected with *Trichomonas gallinae* (Jones-Barn strain). *Avian Diseases* 5: 48–60.
- MUÑOZ, E. 1995. Estudio de la prevalencia y susceptibilidad a la infección por *Trichomonas gallinae* en aves domésticas y silvestres. Valoración de la sensibilidad del protozoo a diferentes derivados nitroimidazólicos. Ph.D. Thesis. Universitat Autònoma de Barcelona, Cerdanyola, Catalonia, Spain, 276 pp.
- NEWTON, I. 1979. Population ecology of raptors. T & A D Poyser, Berkhamsted, UK, 399 pp.
- PALMA, L., CANCELA, L., AND L. OLIVEIRA. 1984. L'alimentation de l'aigle de Bonelli (*Hieraetus fasciatus*) dans la côte portugaise. *In* Rapinyaires Mediterranis, Vol. 2. Centre de Recerca i Protecció dels Rapinyaires (eds.). Barcelona, Spain, pp. 87–96.
- PARELLADA, X. 1984. Variació dels plomatges i identificació de l'àliga cuabarrada *Hieraetus fasciatus*. *In* Rapinyaires Mediterranis, Vol. 2. Centre de Recerca i Protecció dels Rapinyaires (eds.). Barcelona, Spain, pp. 70–79.
- PEPLER, D., AND E. E. OETTLÉ. 1992. *Trichomonas gallinae* in wild raptors on the Cape peninsula. *South African Journal of Wildlife Research* 22: 87–88.
- REAL, J. 1991. L'àliga perdiguera *Hieraetus fasciatus* a Catalunya: status, ecologia tròfica, biologia reproductora i demografia. Ph.D. Thesis, University of Barcelona, Catalonia, Spain, 241 pp.
- . 1996. Biases in diet study methods in the Bonelli's eagle. *The Journal of Wildlife Management* 60: 632–638.
- , AND S. MANOSA. 1997. Demography and conservation of western european Bonelli's eagle *Hieraetus fasciatus* populations. *Biological Conservation* 79: 59–66.
- , ———, AND CODINA, J. 1997. El àguila perdicera en Catalunya. Evolució demogràfica y situació actual. Technical Report. University of Barcelona, Barcelona, Catalonia, Spain, 11 pp.
- REDIG, P. 1987. Medical management of birds of prey. The Raptor Center. Technical Report. University of Minnesota, Saint Paul, Minnesota, 159 pp.
- RETTIG, T. 1978. Trichomoniasis in a bald eagle (*Haliaeetus leucocephalus*) diagnosis and successful treatment with dimetridazol. *Journal of Zoo Animal Medicine* 9: 98–100.
- ROCAMORA, G. 1994. Bonelli's eagle *Hieraetus fas-*

- ciatus*. In Birds of Europe. Their conservation status, G. M. Tucker and M. F. Heath (eds.). Birdlife International, Cambridge, UK, 600 pp.
- SAMOUR, J. H., T. A. BAILEY, AND J. E. COOPER. 1995. Trichomoniasis in birds of prey (order falconiformes) in Bahrain. *Veterinary Record* 136: 358–362.
- STABLER, R. M. 1951. A survey of Colorado band-tailed pigeons, mourning doves, and wild common pigeons for *Trichomonas gallinae*. *The Journal of Parasitology* 37: 471–472.
- . 1954. *Trichomonas gallinae*: A review. *Experimental parasitology* 3: 368–402.
- . 1969. *Trichomonas gallinae* as a factor in the decline of the peregrine falcon. In *Peregrine falcon population. Their biology and decline*. University of Wisconsin, Madison, Wisconsin, pp. 435–437.
- , AND H. A. SHELANSKI. 1936. *Trichomonas columbae* as a cause of death in the hawk. *The Journal of Parasitology* 22: 539–540.
- STONE, W. B., AND P. E. NYE. 1981. Trichomoniasis in bald eagles. *Wilson Bulletin* 93: 109.
- TANGREDI, B. P. 1978. Occurrence of trichomoniasis on Long Island. *New York Fish and Game Journal* 25: 89–90.
- WARD, F. P. 1986. Parasites and their treatment in birds of prey. In *Zoo and wild animal medicine*, M. E. Fowler (ed.). Saunders, Philadelphia, Pennsylvania, pp. 425–430.
- ZAR, J. H. 1984. *Biostatistical analysis*. Prentice-Hall International, Inc. Englewood Cliffs, New Jersey, 718 pp.

Received for publication 11 January 1999.