

Antibodies to *Toxoplasma gondii* in Eurasian Badgers

Authors: Anwar, Ali, Knaggs, Joanna, Service, Katrina M., McLaren, Graeme W., Riordan, Philip, et al.

Source: Journal of Wildlife Diseases, 42(1) : 179-181

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-42.1.179>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Antibodies to *Toxoplasma gondii* in Eurasian Badgers

Ali Anwar,¹ Joanna Knaggs,¹ Katrina M. Service,^{1,4} Graeme W. McLaren,^{1,3,5} Philip Riordan,¹ Chris Newman,¹ Richard J. Delahay,² Chris Cheesman,² and David W. Macdonald¹ ¹ Wildlife Conservation Research Unit, University of Oxford, Tubney House, Abingdon Road, Tubney, Abingdon, OX13 5QL, UK; ² Central Science Laboratory, Sand Hutton, York, YO4 1LZ, UK; ³ Environment Agency, North East Area Office, Apollo Court, 2 Bishops Square Business Park, St Albans Road West, Hatfield, Hertfordshire, AL10 9EX, UK; ⁴ Present address: Department of Health and Biosciences, University of East London, Stratford Campus, Romford Road, Stratford, London, E15 4LZ, UK; ⁵ Corresponding author (email: gwmcl@hotmail.com)

ABSTRACT: Antibodies to *Toxoplasma gondii* were detected in samples collected from 90 live-trapped adult Eurasian badgers (*Meles meles*) sampled at three sites (two agricultural and one woodland) in southern England. Serum was tested using a qualitative latex agglutination test procedure and 63 of 90 (70%) badgers tested positive for *T. gondii* antibodies. Antibody prevalence varied between the sites; 67% and 77% of badgers from agricultural sites and 39% from a nonagricultural site tested positive.

Key words: *Toxoplasma*, Eurasian badger, *Meles*, parasite, protozoan.

Toxoplasma gondii is an intracellular zoonotic protozoan parasite and is possibly the most widespread parasite of warm-blooded animals (Kreier and Baker, 1987), and it is maintained in an indirect life cycle with felids representing the only known definitive hosts. In infected cats, oocysts are produced and are shed in the feces. Oocysts require a few days of adequate temperature and moisture to sporulate before they become infective to birds and mammals, which serve as intermediate hosts (Kreier and Baker 1987; Gajadhar et al., 2004). In intermediate hosts *T. gondii* does not produce oocysts but instead forms cysts in muscular or neural tissues, causing latent toxoplasmosis. The life cycle is completed when a felid ingests infected tissue. Intermediate hosts can also acquire infection through this route and by transplacental transmission (Kreier and Baker, 1987). Prevalence levels of *T. gondii* infection can be high, for example, 62% in outdoor-living domestic cats (Dubey et al., 2002), 35% in wild rats (*Rattus norvegicus*; Webster, 1994), and 12–90% in humans (Hegab and Al-Mutawa, 2003).

In intermediate hosts, latent toxoplasmosis can have behavioral consequences (Webster et al., 1994; Berdoy et al., 1995a, b, 2000). *T. gondii* can manipulate a rat's perception of predation risk, in some cases completely overturning their innate aversion to cats (Berdoy et al., 2000). This change in rat behavior increases the likelihood of cat predation, therefore enhancing the chances of *T. gondii* completing its life cycle. The development of immunity rarely results in the complete destruction of the parasite (Kreier and Baker, 1987) and the presence of antibodies to *T. gondii* is therefore used as an indicator of persistent infection.

In this report we provide the first evidence of *T. gondii* antibodies in Eurasian badgers (*Meles meles*) from the UK. Badgers were sampled from three study areas in the south of England, one in Wiltshire (51°28'N, 2°03'W), one site on the South Gloucestershire/Somerset border (51°27'N, 2°25'W), and one site in Wytham woods (51°46'N, 1°18'W), Oxfordshire. Badgers from the first two sites were largely captured on agricultural land (mainly dairy and beef farms), while the Oxfordshire site is an area of ancient woodland. Animals were live-trapped between 2002 and 2004 (under Home Office and English Nature licenses) as part of a continuing study of their ecology and behavior, following protocols as described by Macdonald and Newman (2002) for Wytham, and Cheesman and Mallinson (1979) for the other sites. Individuals were anesthetized by intramuscular administration of 20 mg/kg ketamine (Vetalar; Pharmacia & Upjohn, Crawley, UK) or a com-

ination of ketamine hydrochloride (100 mg/ml), medetomidine hydrochloride (1 mg/ml, Domitor[®], Pfizer, Sandwich, UK), and butorphanol tartrate (10 mg/ml, Torbugesic[®], Fort Dodge Animal Health Ltd, Southampton, UK) by intramuscular (IM) injection at a ratio of 2:1:2 by volume, respectively, and a dose rate of 0.2 ml/kg⁻¹. Blood was collected by jugular venepuncture, and serum was stored at -20 C. All animals were released at their site of capture.

Serum samples were tested for antibodies to *T. gondii* by using a commercial qualitative latex agglutination test on undiluted serum as described by the manufacturer (Biokit, Barcelona, Spain; Gray et al., 1990). In a previous study, the performance of the latex agglutination test, when compared with that of the Dye test, showed a sensitivity of 98.7%, the specificity was 95.8%, the positive predictive value was 99.4%, and the negative predictive value was 92.0% (Dunford and Johnson, 1991).

Sixty-three of 90 (70%) badgers tested positive for *T. gondii* antibodies; none of the animals sampled showed any clinical signs of toxoplasmosis. There was apparent variation in antibody prevalence between sites, with the two sites in the southwest having 67% ($n=15$) and 77% ($n=60$) testing seropositive, while the Wytham population had a lower prevalence (39% seropositive, $n=15$).

Although this is the first record of *T. gondii* antibodies in badgers, given the widespread occurrence of this parasite, our finding is perhaps not surprising. In the Mustelidae, infection has been reported in black-footed ferrets (*Mustela nigripes*; Burns et al., 2003), mink (*Mustela vison*; Frank, 2001), river otters (*Lontra canadensis*; Tociłowski et al., 1997) and southern sea otters (*Enhydra lutris nereis*; Cole et al., 2000). In clinical situations, where badgers display symptoms such as lethargy, anorexia, and abnormal behavior, a quantitative test for toxoplasmosis should be considered.

LITERATURE CITED

- BERDOY, M., J. P. WEBSTER, AND D. W. MACDONALD. 1995a. The manipulation of rat behaviour by *Toxoplasma gondii*. *Mammalia* 59: 605-613.
- , ———, AND ———. 1995b. Parasite-altered behavior—is the effect of *Toxoplasma gondii* on *Rattus norvegicus* specific. *Parasitology* 111: 403-409.
- , ———, AND ———. 2000. Fatal attraction in rats infected with *Toxoplasma gondii*. *Proceedings of the Royal Society of London Series B-Biological Sciences* 267: 1591-1594.
- BURNS, R., E. S. WILLIAMS, D. O'TOOLE, AND J. P. DUBEY. 2003. *Toxoplasma gondii* infections in captive black-footed ferrets *Mustela nigripes*, 1992-1998, clinical signs, serology, pathology, and prevention. *Journal of Wildlife Diseases* 39: 787-797.
- CHEESEMAN, C. L., AND P. J. MALLINSON. 1979. Radio tracking in the study of bovine tuberculosis in badgers. In *Handbook on biotelemetry and radio tracking*, C. J. Amlaner, Jr., and D. W. MACDONALD (eds.). Pergamon Press, Oxford, UK, pp. 649-656.
- COLE, R. A., D. S. LINDSAY, D. K. HOWE, C. L. RODERICK, J. P. DUBEY, N. J. THOMAS, AND L. A. BAETEN. 2000. Biological and molecular characterizations of *Toxoplasma gondii* strains obtained from southern sea otters *Enhydra lutris nereis*. *Journal of Parasitology* 86: 526-530.
- DUBEY, J. P., W. J. A. SAVILLE, J. F. STANEK, AND S. M. REED. 2002. Prevalence of *Toxoplasma gondii* antibodies in domestic cats from rural Ohio. *Journal of Parasitology* 88: 802-803.
- DUNFORD, P., AND J. JOHNSON. 1991. Detection of toxoplasma-specific immunoglobulin-G: assessment of a slide agglutination test. *Medical and Laboratory Science* 48: 137-141.
- FRANK, R. K. 2001. An outbreak of toxoplasmosis in farmed mink *Mustela vison*. *Journal of Veterinary Diagnostic Investigation* 13: 245-249.
- GAJADHAR, A. A., L. MEASURES, L. B. FORBES, C. KAPEL, AND J. P. DUBEY. 2004. Experimental *Toxoplasma gondii* infection in grey seals *Halichoerus grypus*. *Journal of Parasitology* 90: 255-259.
- GRAY, J. J., A. H. BALFOUR, AND T. G. WREGHITT. 1990. Evaluation of a commercial latex agglutination-test for detecting antibodies to *Toxoplasma gondii*. *Serodiagnosis and Immunotherapy in Infectious Disease* 4: 335-340.
- HEGAB, S. M., AND S. A. AL-MUTAWA. 2003. Immunopathogenesis of toxoplasmosis. *Clinical and Experimental Medicine* 3: 84-105.
- KREIER, J. P. K., AND J. R. BAKER. 1987. *Parasitic protozoa*. Allen & Unwin, Winchester, Massachusetts.
- MACDONALD, D. W., AND C. NEWMAN. 2002. Population dynamics of badgers *Meles meles* in

- Oxfordshire U.K.: Numbers density and cohort life histories and a possible role of climate change in population growth. *Journal of Zoology* 256: 121–138.
- TOCIDLOWSKI, M. E., M. R. LAPPIN, P. W. SUMNER, AND M. K. STOSKOPF. 1997. Serologic survey for toxoplasmosis in river otters. *Journal of Wildlife Diseases* 33: 649–652.
- WEBSTER, J. P. 1994. Prevalence and transmission of *Toxoplasma gondii* in wild brown-rats, *Rattus norvegicus*. *Parasitology* 108: 407–411.
- , C. F. A. BRUNTON, AND D. W. MACDONALD. 1994. Effect of *Toxoplasma gondii* upon neophobic behavior in wild brown-rats, *Rattus norvegicus*. *Parasitology* 109: 37–43.

Received for publication 20 October 2004.