



Occasional Field Observations of the Predation on Mice, Dove and Ants by Black-Tufted-Ear Marmosets (*Callithrix penicillata*)

Authors: Silva, Ita de O., Alvarenga, Adriano B. B., and Boere, Vanner

Source: Neotropical Primates, 15(2) : 59-62

Published By: Conservation International

URL: <https://doi.org/10.1896/044.015.0209>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

beges mais ou menos evidentes; (2) a coloração laranja, marrom ou marrom-alaranjada da cauda; (3) a região cervical branca, bege ou bege-esbranquiçada; (4) o corpo bege estriado de castanho ferrugíneo, bege e ocre, bege com estrias castanhas, cinzas e pretas e bege-brancacentas. Por outro lado, a coloração negra das mãos e pés foi observada em todos os indivíduos. Portanto, consideramos a coloração da pelagem pouco confiável como caráter taxonômico, embora ela tenha sido utilizada juntamente com a morfometria do crânio e a dentição na descrição dessa espécie por Kobayashi e Langguth (1999). Além disso, a coloração da pelagem foi utilizada para distinguir *C. coimbrai* e *C. barbarabrownae*, apesar da variação no padrão de coloração dentro de uma população poder ser maior que aquela observada entre essas duas espécies do grupo *personatus*. A carência de espécimes depositados em coleções científicas compromete a resolução deste desafio taxonômico.

Agradecimentos

À Dra Cecília Kierulff pelas críticas e sugestões ao manuscrito e à Andréa Margit, Anthony Rylands, Russell Mittermeier e Michael Tobias pelo apoio. À Fundação Margot Marsh, Universidade Tiradentes e às Empresas JP pelo apoio financeiro para a realização do estudo.

Marcelo Cardoso de Sousa, Universidade Tiradentes, Av. Murilo Dantas, 300- Aracaju-SE, CEP 49.032-490, mcsousa@infonet.com.br, **Sidnei Sampaio dos Santos**, Associação Baiana para a Conservação dos Recursos Naturais (ABCNRN), Salvador-BA, e **Mariana Coelho Marques Valente**, Universidade Católica de Salvador-BA

Referências Bibliográficas

- Brasil, MMA, 2003. *Lista oficial das espécies da fauna brasileira ameaçada de extinção*. Ministério do Meio Ambiente (MMA), Brasília. Diário oficial da união – Seção 1 101: 88–97.
- Brasil, MMA, 2007. *Relatório de Atualização de Áreas Prioritárias para a Conservação, Uso Sustentável e Repartição de Benefícios da Biodiversidade Brasileira*, Ministério do Meio Ambiente (MMA), Brasília. Portaria n.9 de 23 de janeiro de 2007.
- Coimbra-Filho, A. F. e Câmara, I. de G. 1996. *Os limites originais do Bioma Mata Atlântica na Região Nordeste do Brasil*. Fundação Brasileira para a Conservação da Natureza, Rio de Janeiro, Brasil.
- Jerusalinsky, L., Oliveira, M. M., Pereira, R. F., Santana, V., Bastos, P. C. R. and Ferrari, S. F. 2006. Preliminary evaluation of the conservation status of *Callicebus coimbrai* Kobayashi & Langguth, 1999 in the Brazilian State of Sergipe. *Primate Conservation*. 21: 25–32.
- IUCN, 2007. 2007 IUCN Red List of Threatened Species. Website: <http://www.iucnredlist.org>. Downloaded on the 23rd of March 2007 (“Acessada em 23 de março de 2007”).
- Kinzev, W. G. 1981. The titi monkey, genus *Callicebus*. In: *Ecology and Behavior of Tropical Primates*, Vol. 1, A. F. Coimbra-Filho and R. A. Mittermeier (eds.), pp.241–277. Academia Brasileira de Ciências, Rio de Janeiro, Brazil.
- Kobayashi, S. and Langguth, A. L. 1999. A new species of titi monkey, *Callicebus* Thomas, from north-eastern Brazil (Primates, Cebidae). *Rev. Bras. de Zool.* 16: 531–551.
- Printes, R. C. 2005. Novos registros sobre a distribuição do guigó da caatinga *Callicebus barbarabrownae* (Hershkovitz, 1990) e novo limite sul de *Callicebus coimbrai* (Kobayashi e Langguth, 1999). Em: *Programa e Livro de Resumos: XI Congresso Brasileiro de Primatologia*, p. 154. Sociedade Brasileira de Primatologia, Porto Alegre, Brasil.
- Sistema Nacional de Unidades de Conservação (SNUC). Lei N. 9985 de 18 de julho de 2000. disponível em: <http://www.universia.com.br/materia/img/ilustra/2006/abr/artigos/snuc.pdf> Acessada em: 15 de maio de 2007.
- Sousa, M. C. 2000. New localities for Coimbra-Filho's titi monkey, *Callicebus coimbrai*, in North-east Brazil. *Neotrop. Primates* 8(4): 151.
- Sousa, M. C. 2003. Distribuição do guigó (*Callicebus coimbrai*) no Estado de Sergipe. *Neotrop. Primates* 11(2): 89–91.
- van Roosmalen, M. G. M., van Roosmalen, T. and Mittermeier, R. A. 2002. A taxonomic review of the titi monkeys, genus *Callicebus* Thomas, 1903, with description of two new species, *Callicebus bernhardi* and *Callicebus stephennashi*, from Brazilian Amazonia. *Neotrop. Primates* 10 (Suppl.): 1–52.

OCCASIONAL FIELD OBSERVATIONS OF THE PREDATION ON MICE, DOVE AND ANTS BY BLACK-TUFTED-EAR MARMOSETS (*CALLITHRIX PENICILLATA*)

Ita de O. Silva
Adriano B. B. Alvarenga
Vanner Boere

The black-tufted-ear marmoset (*Callithrix penicillata*) mainly feeds on plant exudates which can comprise up to 70% of the diet (Miranda and Faria, 2001). Other items included in the diet are insects, nectar and flowers, in different proportions. Stevenson and Rylands (1988) classify marmosets from the *Jacchus*-group as exudativore-insectivores. It is known that marmosets opportunistically feed on protein sources i.e. prey, and that their small size allows for a quite diverse diet (Rylands and Faria, 1993). Insects are the most important source of protein, but other protein-rich food currently described for this species are ants, spiders, lizards, snails, frogs, eggs and bird hatchlings (Stevenson and Rylands, 1988; Passamani and

Rylands, 2000). In this article, we report two new food sources, that may be included in the diet of wild marmosets, albeit probably very rarely: mouse and dove. We also report the ingestion of ants, confirming an animal food item in the diet of black-tufted-marmosets mentioned in the literature. Observations on predation by marmosets were made opportunistically when researchers observed the social behavior of two marmosets groups at two different study sites. Both sites, the Estação Ciência São Jose (ECSJ) and the Jardim Botânico de Brasília (JBB), include many vegetation types of the Cerrado biome. The observations were in the *cerradão*, a typical forest with medium to high semi-deciduous trees and xeromorphic vegetation (Ratter *et al.*, 1997).

Predation on a dove (Columbina talpacoti)

The dove predation was observed in the ECSJ, a field station of the Catholic University of Goiás (16° 44' 06" S, 49° 2' 48" W; Goiânia, Brazil), close to the suburb of the city, on 15 March 2001. Around 08:00, a group of black-tufted-ear marmoset moved toward the area around the field laboratory, staying at approximately 3 m height in small trees (*Grevillea robusta*). Suddenly, the marmoset group became very agitated. We succeeded in recording with a digital camera the moment when an adult male, located approximately 2 m above ground, captured a dove (*C. talpacoti*), immediately biting it into the head and starting to eat it (Figure 1). We did not observe the prey being shared with any other group member. The marmoset showed piloerection during the voracious consumption of the dove.

Predation on a mouse (Mus musculus)

The predation on a mouse happened in an area near the entrance of JBB (15° 55' 58" S, 47° 51' 02" W; Brasília, DF, Brazil). On 20 November 2006, at 08:30 h, we saw

the reproductive female of the group capturing a mouse while foraging in the forest understory, approximately 1.5 m above ground, and immediately biting of the head of the mouse (Figure 2). Although other group members, particularly infants, approached the female with characteristic submissive vocalizations, the female did not share the prey, dropping part of the carcass (mostly skin) on the ground.

Predation on ants (Labidus sp.)

The predation of the ants happened along a forest border of the JBB. On 23 August 2006, beginning at 11:15 we observed the marmosets descending to the forest understory above and ground close to a swarm of army ants, identified as *Labidus* (Ecitoninae, Formicidae) by Dr. C. R. F Brandão (Zoology Museum, São Paulo University). The marmosets caught and quickly ingested ants and did not seem to be intimidated by the ants' bites. This continued for approximately 3.5 hours. During this period, the whole group (15 animals) accompanied the swarm front, but among them, only two male adults and two juveniles fed on the ants.

Callithrix penicillata is widely distributed throughout the Cerrado (Stevenson and Rylands, 1988), one of the world's hot spots for biodiversity conservation (Myers *et al.*, 2000). This species, like *C. jacchus*, is found in urbanized areas and has been successfully introduced in several regions (Cunha *et al.*, 2006; Mendes Pontes and Soares, 2005; Miranda and Faria, 2001; Stevenson and Rylands, 1988; Vilela and Faria, 2004). It is presumed to have a flexible and opportunistic diet. Most data available on the diet of the marmosets and tamarins is focused on fruits and exudates, making it necessary to better describe and comprehend the role of prey in the behavioral ecology of Callitrichidae (Nickle & Heymann, 1996; Heymann *et al.*,



Figure 1. Adult male of *Callithrix penicillata* eating a dove (*Columbina talpacoti*) at the Estação Ciência São Jose.



Figure 2. Adult female of *Callithrix penicillata* biting the head of a *Mus musculus* individual previously captured in the forest understory at the Jardim Botânico de Brasília.

2000). The predation of bird nests, mostly for obtaining eggs and hatchlings is well described (Marini and Melo, 1998; Mendes Pontes and Soares, 2005), but according to Stevenson and Rylands (1988), marmosets rarely feed on birds and hatchlings when in their natural habitat. The relevant literature has few reports on the predation of adult birds (Cunha *et al.*, 2006), and the predation of this species of dove in particular has not been previously described. The bird preyed upon is commonly found in urban areas in Brazil (Sick, 1997). The contact from this bird with marmoset groups is presumably common in cities and their surrounding areas. The common mouse is an invasive species of the Brazilian fauna and is closely connected to human activity. The mouse predation reported here occurred in an area with pronounced human influence, and proximity to garbage cans. Newborn mice are used to complement the diet of marmosets kept in captivity (Coimbra-Filho, *et al.*, 1981). However, this is the first description of an adult mouse predation by a marmoset in a wild environment.

Our observations on predation of the ant genus *Labidus* by black-tufted-ear marmoset are in line with recent observations of Melo Jr and Zara (2007) in the Cerrados and Atlantic Forest. Rylands and collaborators (1989) and Melo Jr and Zara (2007) have already described marmosets as predators of ants and insects that are displaced by the raiding ant swarms. Mendes Pontes and Soares (2005) also mention the presence of ants in the marmoset's diet. Although there is a relatively high abundance of this ant species in the woods of the JBB, predation is not commonly observed. While foraging to attain the dove and the mouse was notably an individual behavior, the ants and insects flushed by the ants were eaten while the whole group foraged, as described previously by Passamani and Rylands (2000). The predation on the mouse and the dove, although interesting from the point of view of flexibility in feeding habits, also adds a potential epidemiological link between the several diseases that are present in the region (i.e. zoonotic hemorrhagic fevers, Figueiredo, 2006) and the managing of wild marmosets in preservation areas with strong human influence. Doves and particularly mice are important agents for the dissemination of several pathogens (Pereira *et al.*, 2001; Sick, 1997) that infect both human and non-human primates.

Acknowledgments

We are grateful to Rui Chaves Bozza Jr. for the help with the dove picture. We thank Dr. Carlos Roberto F. Brandão (Zoology Museum, São Paulo University) for the ant identification, Christian Hoffmann for kindly helping with the English translation, and Prof. Dr. Eckhard W. Heymann for editorial improvement of the manuscript.

Ita de O. Silva, Adriano B. B. Alvarenga, and Vanner Boere, Universidade de Brasília, Instituto de Biologia, Departamento de Ciências Fisiológicas, CEP 70910-900, Brasília, DF, Brazil. Corresponding author: Vanner Boere, e-mail: <vanner@unb.br>.

References

- Coimbra-Filho, A. F., Silva, R. R. and Pissinatti, A. 1981. Sobre a dieta de Callitrichidae em cativeiro. *Rev. Biot.* 1: 83–93.
- Cunha, A. A., Vieira, M. V. and Grelle, C. E. V. 2006. Preliminary observations on habitat, support use and diet in two non-native primates in an urban Atlantic forest fragment: The capuchin monkey (*Cebus* sp.) and the common marmoset (*Callithrix jacchus*) in the Tijuca forest, Rio de Janeiro. *Urban Ecosyst.* 9: 351–359.
- Figueiredo, L. T. M. 2006. Febres hemorrágicas por vírus no Brasil. *Rev. Soc. Bras. Med. Trop.* 39: 203–210.
- Heymann, E. W., Knogge, C. and Tirado Herrera, E. R. 2000. Vertebrate predation by sympatric tamarins, *Saguinus mystax* and *Saguinus fuscicollis*. *Am. J. Primatol.* 51: 153–158.
- Marini, M. A. and Melo, C. 1998. Predators of quail eggs, and the evidence of the remains: Implications for nest predation studies. *Condor* 100: 395–399.
- Melo Jr, T. A. and Zara, F. J. 2007. Black-tufted-ear marmoset *Callithrix penicillata* (Primates: Callitrichidae) following the army ant *Labidus praedator* (Formicidae: Ecitoninae) in the Cerrado and Atlantic Forest, Brazil. *Neotrop. Primates* 14: 32–33.
- Mendes Pontes, A. R. and Soares, M. L. 2005. Sleeping sites of common marmosets (*Callithrix jacchus*) in defaunated urban forest fragments: a strategy to maximize food intake. *J. Zool., Lond.* 266: 55–63.
- Miranda, G. H. B. and Faria, D. S. 2001. Ecological aspects of black-pinelled marmoset (*Callithrix penicillata*) in the *cerradão* and dense *cerradão* of the Brazilian Central Plateau. *Braz. J. Biol.* 61: 397–404.
- Myers, N., Mittermeier, R. A., Mittermeier, C. G., Fonseca, G. A. B. and Kent, J. 2000. Biodiversity hotspots for conservation priorities. *Nature* 403: 853–858.
- Nickle, D.A. & Heymann, E. W. 1996. Predation on Orthoptera and other orders of insects by tamarin monkeys, *Saguinus mystax mystax* and *Saguinus fuscicollis nigrifrons* (Primates: Callitrichidae), in north-eastern Peru. *J. Zool.* 239: 799–819.
- Passamani, M. and Rylands, A. B. 2000. Feeding behavior of Geoffroy's marmoset (*Callithrix geoffroyi*) in an Atlantic Forest fragment of south-eastern Brazil. *Primates* 41: 27–38.
- Pereira, L. E., Suzuki, A., Coimbra, T. L. M., Souza, R. P. and Chamelet, E. L. B. 2001. Ilheus arbovirus in wild birds (*Sporophila caerulea* and *Molothrus bonariensis*). *Rev. Sau. Pub.* 35: 119–123.
- Ratter, J. A., Ribeiro, J. F. and Bridgewater, S. 1997. The Brazilian *Cerrado* vegetation and threats to its biodiversity. *Ann. Bot.* 80: 223–230.
- Rylands, A. B., Monteiro Da Cruz, M. A. O. and Ferrari, S. F. 1989. An association between marmosets and army ants in Brazil. *J. Trop. Ecol.* 5:113–116.
- Rylands, A. B. and Faria, D. S. 1993. Habitats, feeding ecology and home range size in the genus *Callithrix*. In: *Marmosets and tamarins: systematics, behaviour and ecol-*

- ogy, A.B. Rylands (ed.), pp. 263–271. Oxford University Press, Oxford.
- Sick, H. 1997. Famílias e espécies: ordem Columbiformes. In: *Ornitologia Brasileira*, J. F. Pacheco (ed.), pp. 341–350. Nova Fronteira, Rio de Janeiro.
- Stevenson, M. F. and Rylands, A.B. 1988. The marmosets, genus *Callithrix*. In: *Ecology and behavior of Neotropical primates*, Mittermeier, R.A., Rylands, A.B., Coimbra-Filho, A. and Fonseca, G.A.B. (eds.), pp.131–222. WWF, Washington.
- Vilela, S. L. and Faria, D. S. 2004. Seasonality of the activity pattern of *Callithrix penicillata* (Primates, Callitrichidae) in the cerrado (scrub savanna vegetation). *Braz. J. Biol.* 64: 363–370.

OBSERVATION OF BLACK-CAPPED CAPUCHINS (*CEBUS APELLA*) FEEDING ON AN OWL MONKEY (*AOTUS BRUMBACKI*) IN THE COLOMBIAN LLANOS

Xyomara Carretero-Pinzón
Thomas R. Defler
Stephen F. Ferrari

Black-capped capuchins (*Cebus apella*) are known to feed on a wide variety of vertebrate prey, including lizards, frogs, birds, bats, marsupials, rodents, and squirrels (Izawa, 1978, 1990; Terborgh, 1983; Galetti, 1990; Rímoli, 2001; Resende *et al.*, 2003; Defler, 2004), although to date, there is only one report involving another primate, an infant titi, *Callicebus moloch* (Sampaio & Ferrari, 2005). Here we describe the behavior of a group of black-capped capuchins feeding on an adult female owl monkey (*Aotus brumbacki*) in a fragment of gallery forest in the Colombian Llanos. This appeared to be the scavenging of a carcass, rather than a predation event, but the observation nevertheless emphasizes the potential of the capuchins for the exploitation of prey this size.

The event was recorded during the long-term monitoring (September 2005 to January 2007) of a group of 43 squirrel monkeys, *Saimiri sciureus albigena* (Carretero-Pinzón, 2008) on the Arrayanes Farm (3°3'30"N, 73°35'40"W) near San Martín in the Colombian Llanos, department of Meta (Fig. 1). The group occupied a matrix of small fragments of gallery forest of up to 21 ha, including those on the neighboring Santa Rosa Farm, that are interconnected by fences. We collected quantitative behavioral data in scan samples (one minute scans at five minute intervals), and whenever an association was formed with the local group of *Cebus apella* [during 28.3% of monitoring time (1,113 hours of total observation time)]. We also collected data on the members of this group (1 adult male, 1 adult female, 2 juveniles, and 1 infant) using the same sampling schedule. During the event reported here we abandoned this schedule and the capuchins were monitored continuously for the first half hour and then at five-minute inter-

vals, until the carcass was abandoned.

On the morning of January the 12th, 2006, when the squirrel monkeys and capuchins were foraging together in a mixed troop at approximately 08:35, the adult male *C. apella* apparently found a dead female owl monkey (*Aotus brumbacki*) in a tree hole. The assumption that the capuchin found the owl monkey rather than captured it alive is based on the lack of visible or audible evidence of hunting activity or prey capture. On finding the carcass, the adult male became visibly excited, and emitted feeding vocalizations, that were answered in kind by the four other group members, who then approached to a distance of 3–5 m from the tree-hole, from where they observed the male. The male pulled the owl monkey's legs out of the hole and began biting, tearing and eating the flesh of one of the legs. The male was the only individual to feed on the carcass during the first eight minutes, but then he moved to a neighboring tree to rest and looked on as the other group members moved in to feed. The adult female and the infant fed on the second leg; then one of the juveniles pulled the carcass out of the hole as far as its neck, bit off the right radius and hand, and ran to the neighboring tree to consume these parts. The four individuals each spent some 8–10 minutes feeding on different parts of the carcass. Tolerance and sharing appears to be typical of vertebrate predation in the capuchins (e.g. Perry and Rose, 1994; Resende *et al.*, 2003), at least where relatively large-bodied prey is concerned (Izawa, 1978). After approximately 20 minutes, the adult male returned to feed on the carcass in close proximity with the infant, while vocalizing aggressively towards the female, who responded submissively and moved to a neighboring tree. The infant eventually joined the female to be groomed, and one juvenile rested nearby while the other foraged for arthropods. All five group members ingested parts of the owl monkey's limbs, tail or dorsal musculature, but, while one of the juveniles probed the abdomen with its hand, none of the capuchins appeared to feed on the intestines or internal organs. Curiously, Resende *et al.* (2003) recorded the opposite pattern in the predation of a rat by *C. apella*, where the soft parts were consumed and the musculature ignored. In the pres-



Figure 1. Location of the study area, showing the main fragments of gallery forest on the Arrayanes (16 ha) and Santa Rosa (21 ha) farms in the Colombian Llanos.