

Early Behavioral Development of a Free-Ranging Howler Monkey Infant (*Alouatta Guariba Clamitans*) in Southern Brazil

Authors: Podgaiski, Luciana Regina, and de Assis Jardim, Mércia Maria

Source: Neotropical Primates, 16(1) : 27-31

Published By: Conservation International

URL: <https://doi.org/10.1896/044.016.0106>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

- Grieser Johns, A. D. and Grieser Johns, B.G. 1995. Tropical forest primates and logging: long-term coexistence? *Oryx* 29: 205–211.
- Henderson, C. L. 2002. *Field Guide to the Wildlife of Costa Rica*. University of Texas Press, Austin.
- Isbell, I. A. 1994. Predation on primates: ecological patterns and evolutionary consequences. *Evol. Anthropol.* 3(2): 61–71.
- Isbell, L. A. and Young, T. P. 1993. Human presence reduces predation in a free-ranging vervet monkey population in Kenya. *Anim. Behav.* 45: 1233–1235.
- Julliot, C. 1994. Predation of a young spider monkey (*Ateles paniscus*) by a crested eagle (*Morphnus guianensis*). *Folia Primatol.* 63: 75–77.
- Ludwig, G., Aguiar, L. M., Miranda, J. M. D., Tixeira, G. M., Svoboda, W. K., Malanski, L. S., Shiozawa, M. M., Hilst, C. L. S., Navarro, I. T., and Passos, F. C. 2007. Cougar predation on black-and-gold howlers on Mutum Island, Southern Brazil. *Int. J. Primatol.* 28(1): 39–46.
- Miranda, J. M. D., Bernardi, I. P., Abreu, K. C., and Passos, F. C. 2005. Predation on *Aloaatta guariba clamitans* Cabrera (Primates, Atelidae) by *Leopardus pardalis* (Linnaeus) (Carnivora, Felidae). *Rev. Bras. Zool.* 22(3): 793–795.
- Miranda, J. M. D., Bernardi, I. P., Moro-Rios, R. F. and Passos, F. C. 2006. Antipredator behavior of brown howlers attacked by black hawk-eagle in Southern Brazil. *Int. J. Primatol.* 27(4): 1097–1101.
- Morrison, J. L. and Humphrey, S. R. 2001. Conservation value of private lands for crested caracaras in Florida. *Conserv. Biol.* 15(3): 675–684.
- Morrison, J. L. and Phillips, L. M. 2000. Nesting habitat and success of the Chimango caracara in southern Chile. *The Wilson Bulletin* 112(2): 225–232.
- Peetz, A., Norconk, M. A. and Kinzey, W. G. 1992. Predation by jaguar on howler monkeys (*Alouatta seniculus*) in Venezuela. *Am. J. Primatol.* 28: 223–228.
- Phillips, K. 1995. Differing responses to a predator (*Eira Barbara*) by *Alouatta* and *Cebus*. *Neotrop. Primates* 3(2): 45–46.
- Raguet-Schofield, M. 2008. The effects of human encroachment and seasonality on the risk of mantled howler monkey (*Alouatta palliata*) predation by dogs on Ometepe Island, Nicaragua. *Am. J. Phys. Anthropol.* Sup 46: 176.
- Richmond, A. R. 1976. Feeding of nestlings by the caracara in Costa Rica. *Wilson Bull.* 88(4): 667.
- Rodríguez-Estrella, R., Donazar, J. A. and Hiraldo, F. 1998. Raptors as indicators of environmental change in the scrub habitat of Baja California Sur, Mexico. *Conserv. Biol.* 12(4): 921–925.
- Sánchez-Zapata, J. A. and Calvo, J. F. 1999. Raptor distribution in relation to landscape composition in semi-arid Mediterranean habitats. *J. Appl. Ecol.* 36: 254–262.
- Seidensticker, J. 1983. Predation by *Panthera* cats and measures of human influence in habitats of south Asian monkeys. *Int. J. Primatol.* 4(3): 323–326.
- Sherman, P. T. 1991. Harpy eagle predation on a red howler monkey. *Folia Primatol.* 56(1): 53–56.
- Shultz, S., Noë, R., McGraw, W. S. and Dunbar, R. I. M. 2004. A community-level evaluation of the impact of prey behavioural and ecological characteristics on predator diet composition. *Proc. R. Soc. B.* 271: 725–732.
- Stanford, C. B. 2002. Avoiding predators: expectations and evidence in primate antipredator behavior. *Int. J. Primatol.* 23(4): 741–757.
- Travaini, A., Donazar, J. A., Ceballos, O. and Hiraldo, F. 2001. Food habits of the crested caracara (*Caracara planicus*) in the Andean Patagonia: the role of breeding constraints. *J. Arid Environ.* 48: 211–219.
- Treves, A. 2002. Predicting predation risk for foraging, arboreal primates. In: *Eat or be Eaten: Predator Sensitive Foraging Among Primates*. L. E. Miller (ed.), pp.222–241. Cambridge University Press, Cambridge.
- Van Schaick, C. P. 1983. Why are diurnal primates living in groups? *Behaviour* 87: 120–144.
- Vargas, R. J., Bó, M. S. and Favero, M. 2007. Diet of the southern caracara (*Caracara planicus*) in Mar Chiquita Reserve, Southern Argentina. *J. Raptor Res.* 41(2): 113–1.

EARLY BEHAVIORAL DEVELOPMENT OF A FREE-RANGING HOWLER MONKEY INFANT (*ALOUATTA GUARIBA CLAMITANS*) IN SOUTHERN BRAZIL

Luciana Regina Podgański
Márcia Maria de Assis Jardim

Introduction

In comparison with other mammals, the offspring of primates undergo a relatively long period of behavioral development and dependency upon their mothers. During this period, the mother is a secure base from which the infant is able to explore the environment and engage in social behavior (Vochteloo *et al.*, 1993), acquiring the ecological skills of the species and the social traditions of the family troop (Southwick and Siddiqi, 1974). Independence is the result of a long period of increasing self-sufficiency in activities such as locomotion and feeding, as well as growing sociability and distance from the mother (Altmann, 1980; Odalia-Rímoli, 1992). Mother-infant ties constitute one of the most intense types of social relationship in primates (Harlow and Zimmermann, 1958, Altmann, 1959), and have a positive effect on infant survival and development.

Howler monkeys (genus *Alouatta* Lacépède, 1799) are the most widely-distributed Neotropical primates (Chapman and Balcomb, 1998), and are also the best studied in the wild. They occupy a variety of habitat types, and are well-adapted to anthropogenic fragmentation (Crocket, 1998), although some species are declining rapidly in the wild and have been classified as threatened (IUCN, 2009). Howlers have been translocated and re-introduced at a number of

sites (e.g. Agoramoorthy, 1995), and a critical factor for such management procedures is the successful handling of females with nursing infants (Baker, 2002). Knowledge of the behavioral development of infants can obviously contribute to the efficiency of such procedures (International Primatological Society, 2007).

Studies of the behavioral development of primate infants have focused on their social behavior and spatial relationships with their mothers. In howlers, studies have been conducted on free-ranging *Alouatta guariba* (Kats and Otta, 1991), *Alouatta palliata* (Altmann, 1959; Clarke, 1990; Lyall, 1996), and *Alouatta seniculus* (Mack, 1979; Cabrera, 1997). Allomaternal care is well documented in female howlers (Calegario-Marques and Bicca-Marques, 1993), but is rare in males (Marques and Adis, 2000). In this study, the early behavioral development of a male infant *A. guariba clamitans* was monitored in a free-ranging group in southern Brazil.

Materials and methods

Study area and subjects

The study was undertaken in Itapuá State Park (30°23'S, 51°30'W), in the city of Viamão, Rio Grande do Sul, Brazil, from August 2003 to April 2004. The climate in the region is temperate, with hot summers and no clear dry season (Cfa type of Köppen's classification; Peel *et al.*, 2007). The home range of the study group was estimated to be approximately 8.71 ha of semi-deciduous forest on a granite hillside bordering a sandy beach, known as Pedreira (Marques, 2001; Jardim, 2005). In August 2003, the study group was composed of two adult males, one sub-adult male, one juvenile male, two adult females, and one male infant (the study subject—see below). Although the exact date of this infant's birth could not be determined, we estimated that it was between two and three months of age, based primarily on body size (Carpenter, 1934; Altmann, 1959; Kowalewski and Zunino, 2004) and the ontogeny of independent behaviors. In March 2004, another male infant was born in the troop, an adult female immigrated, and an adult male disappeared.

Observation methods

From August, 2003 to April, 2004, the behavior of the infant male subject was monitored for two days each month, except October (one day). Data were collected using focal-animal sampling with continuous recording (Altmann, 1974) from sunrise to sunset (around 9 hours of observation per day). Samples of three minutes duration were collected at ten-minute intervals, with a total of 810 samples collected over 155 hours of monitoring. During each sample, the infant's behavior and its position in relation to its mother were recorded according to the categories defined in Table 1. Other behaviors, such as drinking and rejection by the mother, were recorded in *ad libitum* fashion (Altmann, 1974). Nursing was not recorded here because of the difficulty of determining this behavior reliably.

Results

Infant-mother relationships

In the first month of observations, when the infant was two or three months old, he spent 86.8% of the day on the mother, mostly in the ventral position (Fig. 1), although he was carried on the dorsum during troop movements. In subsequent months, the infant spent increasingly less time on the mother. Ventral carrying during troop movements ceased in November (5–6 months), and dorsal carrying in December (6–7 months), although ventral and dorsal contact with the mother were recorded up to March and April, respectively. By the end of the study period, the infant was in contact with its mother less than 30 percent of the time.

Resting (Fig. 2a)

Initially, the infant spent most of its time at rest, and almost always rested while being carried by the mother until 4–5 months old. The infant was observed resting at a distance from its mother for the first time in November, but this only became common by the end of the study. By the age of 8–9 months, time spent resting stabilized at 55–60% of observation time.

Locomotion (Fig. 2b)

The earliest bouts of independent locomotion occurred in September (3–4 months old). This activity increased progressively until 6–7 months, but was subsequently irregular. Independent movement of the infant was monitored and stimulated by the mother, by moving very slowly through the trees and waiting for the infant to follow. Sometimes, the mother would move to a branch and emit a vocalization until the infant arrived. When crossing a wide gap in the trees, the mother used her own tail or body as a bridge or carried the infant on her dorsum.




Figure 1. Mean percentage observation time spent by the *A. guariba* infant in different contact categories (see Table 1) in each month during the study period.




Figure 2. Percentage of time spent by the howler infant in different behaviors and in different forms of contact with its mother during the study period. See Table 1 for behavior and contact categories.

Feeding (Fig. 2c)

The infant also started eating solid food at 3–4 months. Initially, the infant was usually either being carried by its mother or close to her during this behavior, which allowed it to observe the items ingested by the mother and then repeat her movements. Contact declined rapidly by the following month. The howlers usually drank water from bromeliads, and this behavior was first noticed in the infant at 8–9 months, the same period when the mother began rejecting the infant when it tried to nurse.

Grooming (Fig. 2d)

The infant was observed grooming on two occasions. On the first, he was 4–5 months old, and mimicked his mother when grooming an adult female. On the second occasion, at 6–7 months, the infant groomed its mother.

Play (Fig. 2e–f)

The infant already exhibited play behavior when observations began. The highest frequency occurred at 5–6 months, reaching around 50% of the daily observation period. After this peak, the behavior tended to decline towards the end of the study. Play almost always took place when the troop was at rest. During individual play, the infant explored its immediate environment. Social play initially involved the infant’s mother, and gradually involved a juvenile male. Play was also recorded with the dominant adult male.

Discussion

While the infant grew progressively independent, it maintained relatively close ties with its mother, invariably resting together, even at the end of the study. Overall, the behavioral development of the infant was similar to that recorded in previous studies of howlers (Altmann, 1959; Mack, 1979; Clarke, 1990; Kats and Otta, 1991; Lyall, 1996;

Cabrera, 1997). The transition from exclusive ventral carrying in the first few weeks of life to a predominance of the dorsal position is a typical pattern in monkeys, including howlers (Altmann, 1959; Mack, 1979; Shoemaker, 1979; Kats and Otta, 1991; Lyall, 1996; Miranda *et al.*, 2005). Increasing distance from the mother is also typical. Kats and Otta (1991) confirmed a progressive increase in distance from the mother at 3–4 month of age in *A. guariba*, while Miranda *et al.*, (2005) found that contact with the mother declined to 10% of activity time by the fifth month of age. In the present study, the infant was more than 1 m from its mother for 6% of the time at 2–3 months old, increasing to 70% only at 1 year. A possible regression in this trend was recorded at 7–8 months, however, which may represent a critical period of insecurity or a regression in suckling (Horwich, 1989).

According to Mack (1979), prior to six months of age, howler infants may bite and chew the same type of object that the adults are eating, but not necessarily consume it. Maybe, the high feeding percentages observed in the first months of this study refer to an exploratory feeding which occurred in contact or near the mother. The stabilization of rest at 55–60% by 7–8 months corresponds to the typical rate of adults in this study group (Marques, 2001). This stabilization occurred at the same time as the frequency of play behavior began to decline. Both play and grooming have an important social function (Southwick and Siddiqi, 1974). One possible factor determining rates of social play in the present study was the availability of potential partners, i.e. other immature individuals. Grooming appears to be relatively common in *A. guariba* in comparison with other howlers (Kinzey, 1997); this behavior was part of the behavioral development of the studied infant. While behavior patterns are best evaluated on the basis of a sample of different individuals, the present study provides some useful preliminary insights on

Table 1. Howler monkey infant behavioral categories based on proximity to the mother and activity.

Position	Description
Ventral	Carried by the mother in a ventral position
Dorsal	Carried by the mother in a dorsal position
In contact	In contact with the mother’s body, other than the ventrum or dorsum
< 1m	Less than a meter from the mother’s body
> 1m	More than a meter from the mother’s body
Activity	Description
Rest	Sitting or lying
Locomotion	Moving independently
Feed	Handling, processing or ingesting solid foods
Grooming	Sifting through another animal’s fur with the hands
Individual play	Handling objects, hanging from tail and jumping between branches
Social play	Interactions with other troop members involved mock fighting (holding, pulling, baring teeth, and biting) and chasing, often involving jumping
Drink	Ingesting water

the behavioral development of infant howlers of the species *Alouatta guariba*. The patterns observed appeared to be typical of the genus.

Acknowledgments

We thank Ana Alice B. de Marques for initial advice on the project; Paula C. Colombo for technical support; Aline F. Quadros, André F. B. Lima, and Verônica G. Sydow for comments on earlier versions of the manuscript.

Luciana Regina Podgaiski, Programa de Pós-Graduação em Ecologia, Instituto de Biociências, Universidade Federal do Rio Grande do Sul (UFRGS), Av. Bento Gonçalves, 9500; Prédio 43422, Postal box: 15007, Porto Alegre, Rio Grande do Sul, Brazil, <podgaiski@gmail.com> and **Márcia Maria de Assis Jardim**, Setor de Mastozoologia, Museu de Ciências Naturais (MCN), Fundação Zoobotânica do Rio Grande do Sul (FZBR), Porto Alegre, Rio Grande do Sul, Brazil <marcia.jardim@fzb.rs.gov.br>

References

- Agoramoorthy, G. 1995. Red howling monkey (*Alouatta seniculus*) reintroduction in a gallery forest of Hato Flores Moradas, Venezuela. *Neotrop. Primates* 3: 9–10.
- Altmann, S. A. 1959. Field observation on howling monkey society. *J. Mammal.* 40: 317–330.
- Altmann, J. 1974. Observational study of behavior: sampling methods. *Behaviour* 49: 227–267.
- Altmann, J. 1980. *Baboon mothers and infants*. Harvard University Press, Cambridge.
- Baker, L. R. 2002. Guidelines for nonhuman primate reintroductions. *Reintroduction News* 21: 29–57.
- Cabrera, J. A. 1997. Cambios en la actividad de juegos en infantes y jóvenes de mono aullador (*Alouatta seniculus*). *Neotrop. Primates* 5: 108–111.
- Calegari-Marques, C. and Bicca-Marques, J. C. 1993. Allomaternal care in the black howler monkey (*Alouatta caraya*). *Folia Primatol.* 61: 104–109.
- Carpenter, C. R. 1934. A field study of the behaviour and social relations of howling monkeys (*Alouatta palliata*). *Com. Psychol. Monogr.* 10: 1–168.
- Chapman, C. A. and Balcomb, S. R. 1998. Population characteristics of howlers: ecological conditions or group history. *Int. J. Primatol.* 19: 385–403.
- Clarke, M. R. 1990. Behavioral development and socialization of infants in a free-ranging group of howling monkeys (*Alouatta palliata*). *Folia Primatol.* 54: 1–15.
- Harlow, H. F. and Zimmermann, R. R. 1958. The development of affectional responses in infant monkeys. *Proc. Am. Phil. Soc.* 102: 501–509.
- Horwich, R. H. 1989. Cyclic development of contact behaviour in apes and humans. *Primates* 30: 269–279.
- International Primatological Society. 2007. IPS International guidelines for acquisition, care and breeding of nonhuman primates. Website: <http://www.internationalprimatologicalsociety.org/docs/IPS>
- International Guidelines for the Acquisition Care and Breeding of Nonhuman Primates Second Edition 2007. pdf. Accessed 18 August 2008.
- IUCN. 2009. IUCN Red List of Threatened Species. Version 2009.1. Website: www.iucnredlist.org Accessed 18 June 2009.
- Jardim, M. M. A. 2005. Ecologia populacional de bugios-ruivos (*Alouatta guariba*) nos municípios de Porto Alegre e Viamão, RS, Brasil. Doctoral thesis, Universidade Estadual de Campinas, Campinas, SP.
- Kats, B. and Ota, E. 1991. Comportamento lúdico do bugio (*Alouatta fusca clamitans*, Cabrera, 1940) (Primates: Cebidae: Alouattinae). *Biotemas* 4: 61–82.
- Kinzey, W. G. 1997. *Alouatta*. In: *New world primates: ecology, evolution and behaviour*, W. G. Kinzey (ed.), pp. 174–185. Aldine de Gruyter, New York.
- Kowalewski, M. and Zunino, G. E. 2004. Birth seasonality in *Alouatta caraya* in Northern Argentina. *Int. J. Primatol.* 25: 383–400.
- Lyall, Z. S. 1996. The early development of behavior and independence in howler monkeys *Alouatta palliata mexicana*. *Neotrop. Primates* 4: 4–8.
- Mack, D. 1979. Growth and development of infant red howling monkeys (*Alouatta palliata*) in a free-ranging population. In: *Vertebrate Ecology in the Northern Neotropics*, J. F. Eisenberg (ed.), pp. 127–136. Smithsonian Institution Press, Washington.
- Marques, A. A. B. 2001. Estratégias de uso do espaço por *Alouatta guariba clamitans* Cabrera, 1940 em habitats temperado e subtropical no sul do Brasil. Doctoral thesis, Universidade Federal de Minas Gerais, Belo Horizonte, MG.
- Marques, A. A. B. and Adis, C. 2000. Male care in a group of wild *Alouatta fusca clamitans* in Southern Brazil. *Folia Primatol.* 71: 409–412.
- Menegat, R., Mohr, F. V., Carraro, C. C. and Flores, R. 1998. Porto Alegre em dados. In: *Atlas Ambiental de Porto Alegre*, R. Menegat (ed.), pp. 203–205. Editora da Universidade, Porto Alegre.
- Miranda, J. M. D., Aguiar, L. M., Ludwig, G., Moro-Rios, R. F. and Passos, F. C. 2005. The first seven months of an infant of *Alouatta guariba* (Humboldt) (Primates, Atelidae): interactions and the development of behavioral patterns. *Rev. Bras. Zool.* 22: 1191–1195.
- Odalia-Rímoli, A. 1992. O filhote muriqui (*Brachyteles arachnoides*): um estudo do desenvolvimento da independência. Master dissertation. Universidade de São Paulo, São Paulo, SP.
- Shoemaker, A. H. 1979. Reproduction and development of the Black howler monkey *Alouatta caraya* at Columbia Zoo. *Int. Zoo Yearb.* 19: 150–155.
- Southwick, C. H. and Siddiqi, M. F. 1974. Contrasts in primate social behaviour. *Bioscience* 24: 398–406.
- Vochteloo, J. D., Timmermans, P. J. A., Duijghuisen, J. A. H. and Vossen, M. H. 1993. Effects of reducing the mother's radius of action on the development of mother-infant relationships in longtailed macaques. *Anim. Behav.* 45: 603–612.