

## Two-Component Signaling Elements and Histidyl-Aspartyl Phosphorelays †

Authors: Schaller, G. Eric, Kieber, Joseph J., and Shiu, Shin-Han

Source: The Arabidopsis Book, 2008(6)

Published By: The American Society of Plant Biologists

URL: <https://doi.org/10.1199/tab.0112>

---

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

First published on July 14, 2008: e0112. doi: 10.1199/tab.0112

This chapter is an updated version of a chapter originally published on March 27, 2002, doi:10.1199/tab.0086

# Two-Component Signaling Elements and Histidyl-Aspartyl Phosphorelays

G. Eric Schaller,<sup>a,1</sup> Joseph J. Kieber,<sup>b</sup> and Shin-Han Shiu<sup>c</sup>

<sup>a</sup>Department of Biological Sciences, Dartmouth College, Hanover, NH 03755

<sup>b</sup>Department of Biology, University of North Carolina, Chapel Hill, NC 27599

<sup>c</sup>Department of Plant Biology, Michigan State University, East Lansing, MI 48824

<sup>1</sup>Address correspondence to george.e.schaller@dartmouth.edu

**Two-component systems are an evolutionarily ancient means for signal transduction. These systems are comprised of a number of distinct elements, namely histidine kinases, response regulators, and in the case of multi-step phosphorelays, histidine-containing phosphotransfer proteins (HPts). Arabidopsis makes use of a two-component signaling system to mediate the response to the plant hormone cytokinin. Two-component signaling elements have also been implicated in plant responses to ethylene, abiotic stresses, and red light, and in regulating various aspects of plant growth and development. Here we present an overview of the two-component signaling elements found in Arabidopsis, including functional and phylogenetic information on both bona-fide and divergent elements.**

## INTRODUCTION

Protein phosphorylation is a key mechanism for regulating signal transduction pathways in both prokaryotes and eukaryotes. In eukaryotes, regulatory phosphorylation predominantly occurs at serine, threonine, and tyrosine residues (Hunter, 1995; Hunter and Plowman, 1997; Plowman et al., 1999). In contrast, many signal transduction pathways in bacteria employ a so-called “two-component system” that relies upon phosphorylation of histidine and aspartic-acid residues (Mizuno, 1997). Plants also make use of two-component systems and these perform important roles in growth and development (Schaller, 2000; Mizuno, 2005; To and Kieber, 2008).


Two-component systems confer upon bacteria the ability to sense and respond to environmental stimuli and are involved in such diverse processes as chemotaxis, osmotic sensing, oxygen sensing, and host recognition (Parkinson, 1993; Stock et al., 2000; Baker et al., 2006). The simplest form of a two-component system employs a receptor with histidine-kinase activity and a response regulator (Figure 1). The receptor is located in the plasma membrane of the bacterium. In response to an environmental stimulus, the histidine kinase autophosphorylates a conserved histidine residue. This phosphate is then transferred to a conserved aspartic acid residue within the receiver domain of the response regulator. Phosphorylation of the response regulator modulates its ability to mediate downstream signaling. In bacteria, many of the response regulators are transcription factors. Thus, two proteins create a signaling circuit, capable of converting an external stimulus into a change in transcription. There are permutations on the two-component system. Of particular relevance to the plant two-component systems are multi-step phosphorelays (Figure 1) (Swanson et al., 1994; Appleby et al., 1996). These make use of

a “hybrid” kinase that contains both histidine kinase and receiver domains in one protein, a histidine-containing phosphotransfer (HPt) protein, and a separate response regulator (Appleby et al., 1996). In these multi-step phosphorelays, the phosphate is transferred from amino acid to amino acid in sequence His → Asp → His → Asp.

Although originally identified in bacteria, two-component signaling elements have also been identified in fungi, slime molds, and plants (Swanson et al., 1994; Loomis et al., 1997; Schaller, 2000; Mizuno, 2005). Interestingly, the canonical histidyl-aspartyl phosphorelay is apparently not found in animals. Analysis using the SMART protein-domain search interface (<http://SMART.embl-heidelberg.de>) (Schultz et al., 2000) indicates that bona-fide two-component signaling elements are lacking from the genome sequences of *Homo sapiens*, *Drosophila melanogaster*, and *Caenorhabditis elegans*.

In Arabidopsis, proteins with significant sequence similarities to all elements of the two-component system have been identified, including histidine kinases, response regulators, and HPt proteins (Figure 2) (Schaller, 2000; Mizuno, 2005). Phosphorylation activity has been confirmed for at least one example in each case (Gamble et al., 1998; Imamura et al., 1998; Miyata et al., 1998; Suzuki et al., 1998; Imamura et al., 1999). In recent years, multiple experimental approaches have demonstrated the action of a histidyl-aspartyl phosphorelay in mediating cytokinin signal transduction (Hwang and Sheen, 2001; Higuchi et al., 2004; Nishimura et al., 2004; Mason et al., 2005; Hutchison et al., 2006; Riefler et al., 2006; Yokoyama et al., 2007).

Arabidopsis also contains divergent two-component-like elements that are unlikely to function in histidyl-aspartyl phosphorelays (e.g. phytochromes and pseudo-response regulators) as they are missing one or more key amino-acid residues involved in a




**Figure 1.** Histidyl-Aspartyl Phosphorelay.

Histidine kinase domains are indicated by rectangles, receiver domains by ovals, HPT proteins by rounded rectangles, and transmembrane domains by black bars. Sites of phosphorylation upon histidine (H) and aspartic acid (D) residues are indicated. Terminology is based on that of Parkinson (1993).

(A) Simple two-component system that employs a histidine kinase and a response regulator.

(B) Multi-step phosphorelay that employs a hybrid histidine kinase with both histidine kinase and receiver domains, a histidine-containing phosphotransfer protein (HPT), and a response regulator.



**Figure 2.** Representative Domain Structures of Arabidopsis Two-Component Signaling Elements.

phosphorelay (Figure 2) (Makino et al., 2000; Schaller, 2000). Plants, like animals, contain pyruvate dehydrogenase kinase; this enzyme is related to histidine kinases but has an altered specificity such that it now phosphorylates serine residues (Popov et al., 1993). Thus, there is evidence from eukaryotes that two-component signaling elements have evolved to fill new functions that no longer rely upon phosphorylation of histidine and aspartic acid residues.

## HISTIDINE KINASES

Analysis of the Arabidopsis genome supports the existence of eight histidine kinases that contain all the conserved residues required for enzymatic activity (Table 1, Figures 2 and 3), as well as additional diverged histidine-kinase like proteins that lack residues essential to histidine kinase activity. Some of the functional histidine-kinases have been identified as receptors for the plant hormones cytokinin (AHK2, AHK3, and AHK4) and ethylene (ETR1 and ERS1), but the ligands for the other three bona-fide histidine kinases (AHK1, CKI1, and CKI2) have yet to be determined.

## Cytokinin Receptor Family


The cytokinin receptor family is composed of three histidine kinases: AHK2, AHK3, and AHK4 (also called WOL1 or CRE1). Initial evidence that this family functions in cytokinin perception came from the study of AHK4 transgenically expressed in bacteria and yeast, where it was shown that AHK4 could bind cytokinins and that ligand-binding stimulated the receptor's ability to signal through a phosphorelay (Inoue et al., 2001; Suzuki et al., 2001; Ueguchi et al., 2001; Yamada et al., 2001). All three receptors contain transmembrane domains, are thought to be localized to the plasma membrane, and contain a CHASE (cyclases/histidine kinases associated sensing extracellular) domain in their predicted extracellular portion that functions in cytokinin binding (Anantharaman and Aravind, 2001; Heyl et al., 2007). The isolation and characterization of T-DNA insertion mutations has demonstrated roles for the cytokinin receptors in diverse cytokinin-regulated processes including cell division,

**Figure 2.** (continued)

Histidine kinase domains are indicated by rectangles, receiver domains by ovals, HPT proteins by rounded rectangles, and transmembrane domains by black bars. Additional domains are as indicated and are described in the text.

(A) Bona-fide signaling elements likely to participate in phosphorelays. Sites of phosphorylation upon histidine (H) and aspartic acid (D) residues are indicated.

(B) Divergent signaling elements from Arabidopsis that lack residues normally associated with participation in a two-component phosphorelay. An X in a histidine-kinase-like domain indicates that it lacks multiple residues implicated in histidine kinase activity. The divergent HPT (AHP1) contains an Asn (N) substitution for the His that normally gets phosphorylated. The divergent response regulators typically contain a Glu (E) substitution for the Asp that normally gets phosphorylated.



**Figure 3.** Phylogenetic Relationship of Histidine Kinases.

Sequences containing conserved histidine-kinase domains (open circles) and diverged histidine kinase domains (filled squares) were identified in Arabidopsis. An unrooted bootstrapped tree is shown based on amino-acid alignment of the histidine-kinase domain sequences. The histidine kinase domain was defined as the sequence extending from the phospho-acceptor domain through the ATPase domain.

vascular differentiation, leaf senescence, seed size, and stress responses (Higuchi et al., 2004; Nishimura et al., 2004; Riefler et al., 2006; Tran et al., 2007).

### Ethylene Receptor Family

The ethylene receptor family of Arabidopsis (ETR1, ERS1, ETR2, ERS2, and EIN4) also contains proteins with histidine kinase domains (Chang et al., 1993; Hua et al., 1995; Hua et al., 1998; Sakai et al., 1998). ETR1 and ERS1 have been demonstrated to bind ethylene when transgenically expressed in yeast (Schaller and Bleecker, 1995; Rodriguez et al., 1999; Hall et al., 2000) and both contain functional histidine kinase domains (Gamble et al., 1998; Moussatche and Klee, 2004). Although histidine-kinase activity has been implicated in subtle modulations of the ethylene response, no major role has yet been identified in ethylene signal transduction (Wang et al., 2003; Binder et al., 2004; Qu and Schaller, 2004). Mutational analysis indicates that, rather than relying on a histidyl-aspartyl phosphorelay, ethylene signal transduction incorporates the Raf-like kinase CTR1, the

membrane-bound Nrap-like protein EIN2, and the EIN3 family of transcription factors (Chen et al., 2005). Histidine kinase activity of the ethylene receptors could play a direct but lesser role in ethylene signal transduction. Alternatively, histidine kinase activity could allow for cross-talk between ethylene perception and other two-component signaling pathways such as cytokinin signal transduction.

The ethylene-receptor family also includes three members (ETR2, ERS2, and EIN4) that contain divergent histidine-kinase domains (Bleecker, 1999; Schaller, 2000). Thus, one protein family of related function contains both bona-fide and divergent histidine kinases. Based on in-vitro phosphorylation assays, ETR2, ERS2, and EIN4 are now thought to function as ser/thr kinases (Moussatche and Klee, 2004). In the same study ERS1 was found to phosphorylate serine in addition to histidine, suggesting that it might be bi-functional.

### CKI1

*CKI1* was initially identified in a mutant screen where it was found that its ectopic expression results in cytokinin-independent greening and shoot induction in callus cultures (Kakimoto, 1996). However *CKI1* is not closely related to the cytokinin-receptor family and lacks the CHASE domain involved in cytokinin binding. Thus, the cytokinin-related phenotype resulting from *CKI1* over-expression may arise due to non-physiological cross-talk with the cytokinin-signaling pathway. Analysis of loss-of-function mutations has clarified the role of *CKI1* in plant growth and development, although a regulatory ligand for this putative receptor has not been identified. The homozygous *cki1* mutation is lethal and examination of the mutant indicates that *CKI1* is required for megagametogenesis, consistent with expression data that indicate *CKI1* is expressed in developing ovules (Pischke et al., 2002; Hejatko et al., 2003).

### CKI2

*CKI2*, like *CKI1*, can induce cytokinin responses when ectopically expressed (Kakimoto, 1996), but is also not thought to be directly involved in cytokinin signaling. *CKI2* shows strong expression in the root and weaker expression in flowers (Iwami et al., 2007). Loss-of-function mutations in *CKI2* exert a subtle phenotype in which root elongation is more sensitive to growth inhibition in response to ethylene, suggesting possible cross-talk with the ethylene receptors (Iwami et al., 2007).

### AHK1

AHK1 was initially proposed to function as a plant osmosensor based on its ability to complement function of the yeast osmosensor *SLN1*, a histidine kinase that regulates the HOG1 MAP-kinase pathway for osmosensing in yeast (Urao et al., 1999). However it was subsequently found that other histidine kinases of Arabidopsis, including the cytokinin receptors, can complement yeast *SLN1* mutations, indicating that this assay is not necessarily diagnostic of plant function (Reiser et al., 2003; Tran et al.,

**Table 1.** Histidine Kinase-like Proteins of Arabidopsis

Gene Names	Synonyms	Chrom. Locus <sup>a</sup>	Features <sup>b</sup>	Family	References <sup>c</sup>
ETR1		At1g66340	C <sub>2</sub> H <sub>4</sub> , GAF, HK, Rec	Ethylene Receptor	Chang et al. 1993
ERS1		At2g40940	C <sub>2</sub> H <sub>4</sub> , GAF, HK	Ethylene Receptor	Hua et al. 1995
ETR2		At3g23150	C <sub>2</sub> H <sub>4</sub> , GAF, HKL, Rec	Ethylene Receptor	Sakai et al. 1998b
ERS2		At1g04310	C <sub>2</sub> H <sub>4</sub> , GAF, HKL	Ethylene Receptor	Hua et al. 1998
EIN4		At3g04580	C <sub>2</sub> H <sub>4</sub> , GAF, HKL, Rec	Ethylene Receptor	Hua et al. 1998
AHK4	WOL1, CRE1	At2g01830	CHASE, HK, Rec	Cytokinin Receptor	Mähönen et al. 2000, Inoue et al. 2001, Suzuki et al. 2001
AHK2		At5g35750	CHASE, HK, Rec	Cytokinin Receptor	Ueguchi et al. 2001
AHK3		At1g27320	CHASE, HK, Rec	Cytokinin Receptor	Ueguchi et al. 2001
CKI1		At2g47430	HK, Rec		Kakimoto, 1996, Pischke et al. 2002
AHK1	ATHK1	At2g17820	HK, Rec		Urao et al. 1999
CKI2	AHK5	At5g10720	HK, Rec		Kakimoto, 1996, Iwama et al. 2007
PHYA		At1g09570	GAF, HKL	Phytochrome	Sharrock and Quail, 1989
PHYB		At2g18790	GAF, HKL	Phytochrome	Sharrock and Quail, 1989
PHYC		At5g35840	GAF, HKL	Phytochrome	Sharrock and Quail, 1989
PHYD		At4g16250	GAF, HKL	Phytochrome	Sharrock et al. 1994
PHYE		At4g18130	GAF, HKL	Phytochrome	Sharrock et al. 1994
PDK		At3g06483	HKL	Pyruvate dehydrogenase kinase	Thelen et al. 2000

<sup>a</sup>Chromosome loci are given by the MIPS designation.

<sup>b</sup>Features noted are conserved histidine-kinase domain (HK), diverged histidine-kinase like domain (HKL), receiver domain (Rec), ethylene-binding domain (C<sub>2</sub>H<sub>4</sub>), CHASE domain for cytokinin binding (CHASE), and GAF domain which in phytochromes serves as the chromophore-binding site.

<sup>c</sup>References are chosen based on initial descriptions of genes and their synonyms

2007). Rather, genetic analysis indicates that *AHK1* modulates plant growth and stress responses based on the finding that an *ahk1,ahk2,ahk3* triple mutant (i.e. a combination of *ahk1* with two cytokinin receptor mutations) is reduced in size compared to the *ahk2,ahk3* double mutant (Tran et al., 2007). *AHK1* is induced by dehydration and acts as a positive regulator of drought and salt stress responses, in contrast to the cytokinin receptors which are also induced by dehydration but act as negative regulators of these responses (Tran et al., 2007).

### Phytochrome Family

The phytochromes, which act as red light receptors (Sharrock and Quail, 1989; Clack et al., 1994; Rockwell et al., 2006), are divergent histidine kinases that possess ser/thr kinase activity rather than histidine kinase activity (Yeh and Lagarias, 1998).

### Pyruvate dehydrogenase kinase

Pyruvate dehydrogenase kinase retains the most highly conserved residues found in histidine kinases, but has nevertheless diverged significantly in other residues, resulting in an ability to phosphorylate its substrate, pyruvate dehydrogenase, at a serine residue (Thelen et al., 2000).

### HISTIDINE-CONTAINING PHOSPHOTRANSFER PROTEINS

Histidine-containing phosphotransfer (HPt) proteins function in multi-step phosphorelays, acting as signaling intermediates between hybrid histidine kinase and response regulators (Figure 1). The Arabidopsis genome encodes five HPt proteins (AHP1 through 5) that contain the conserved residues required for activity, as well as one pseudo-HPt (APHP1/APHP6) that lacks the histidine phosphorylation site (Table 2, Figures 2 and 4) (Miyata et al., 1998; Suzuki et al., 1998; Suzuki et al., 2000). The HPt proteins have been shown capable of participating in a phosphorelay with Arabidopsis response regulators (Suzuki et al., 1998). In addition, two-hybrid analysis has demonstrated their ability to interact with both hybrid histidine kinases and response regulators (Imamura et al., 1999; Urao et al., 2000; Tanaka et al., 2004; Dortay et al., 2007), consistent with an ability to function in a multi-step phosphorelay.


Analysis of loss-of-function mutations has revealed that *AHP1*, *AHP2*, *AHP3*, and *AHP5* function as redundant positive regulators of cytokinin signaling (Hutchison et al., 2006). *AHP4* only contributes slightly to the cytokinin responses, and in some cases appears to act as a negative regulator (Hutchison et al., 2006). The AHPs have been shown to accumulate in the nucleus in response to exogenous cytokinin (Hwang and Sheen, 2001; Yamada et al., 2004). *APHP1*, which encodes a pseudo-HPt lacking the

**Table 2.** HPt Proteins of Arabidopsis

Gene Names	Synonyms	Chrom. Locus <sup>a</sup>	Features <sup>b</sup>	References
AHP1	ATHP3	AT3g21510	HPt	Suzuki et al. 1998, Miyata et al. 1998
AHP2	ATHP1	AT3g29350	HPt	Suzuki et al. 1998, Miyata et al. 1998
AHP3	ATHP2	AT5g39340	HPt	Suzuki et al. 1998, Miyata et al. 1998
AHP4		AT3g16360	HPt	Suzuki et al. 2000
AHP5		At1g03430	HPt	Suzuki et al. 2000
APHP1	AHP6	At1g80100	pseudo-HPt	Suzuki et al. 2000, Mähönen et al. 2006

<sup>a</sup>Chromosome loci are given by the MIPS designation.

<sup>b</sup>Features indicate whether the protein contains a conserved histidine-containing phosphotransfer domain (HPt) or a pseudo-HPt lacking the His that is phosphorylated

**Figure 4.** Phylogenetic Relationship of HPt Proteins.

An unrooted bootstrapped tree is shown based on amino-acid alignment of HPt domain sequences.

phosphorylation site, acts as a negative regulator of cytokinin responses (Mähönen et al., 2006b). *APHP1* expression is induced by cytokinin and thus functions as part of a negative feedback loop to reduce the plants sensitivity to cytokinin, with *APHP1* potentially interacting with the receiver domains of the cytokinin receptors to prevent phospho-transfer to bona-fide AHPs.

## RESPONSE REGULATORS

There are 23 genes in the Arabidopsis genome encoding proteins predicted to be functional response regulators (Table 3, Figures 2 and 5). These authentic response regulators (ARRs) can be divided into three classes based on phylogenetic analysis and func-

tion: type-A type-B, and type-C response regulators (Imamura et al., 1999; Schaller et al., 2007). In addition, there are nine genes encoding response regulators that lack the conserved Asp for phosphorylation, and are called pseudo-response regulators (PRRs) (Makino et al., 2000).


### Type-A response regulators

The type-A response regulators are relatively small, containing a receiver domain along with short N- and C-terminal extensions (Brandstatter and Kieber, 1998; Imamura et al., 1998; Urao et al., 1998). Members of the type-A family are transcriptionally induced to varying extents by cytokinin (Brandstatter and Kieber, 1998; Taniguchi et al., 1998; D'Agostino et al., 2000), and cytokinin also stabilizes some type-A ARR proteins in a phosphorylation-dependent manner (To et al., 2007). Genetic analyses indicate that *ARR3*, *4*, *5*, *6*, *7*, *8*, *9* and *15* function as negative regulators of cytokinin signaling, thus participating in a negative feedback loop to reduce the plant sensitivity to cytokinin (Kiba et al., 2003; To et al., 2004; Leibfried et al., 2005; Lee et al., 2007; To et al., 2007).

Some type-A ARRs are also implicated in other regulatory pathways. *ARR4* interacts with phytochrome B to modulate its activity, allowing for cross-talk between the cytokinin and light signaling pathways (Sweere et al., 2001; Mira-Rodado et al., 2007). *WUSCHEL*, a regulator of stem cells in the shoot apical meristem, negatively regulates transcription of several type-A ARRs, revealing cross-talk between cytokinin signaling and a key meristem identity gene (Leibfried et al., 2005). *ARR3* and *ARR4* regulate the circadian period in a cytokinin-independent manner, with loss of these two genes resulting in a longer clock period (Salomé et al., 2006).

### Type-B response regulators

The type-B ARRs differ from the type-A ARRs in that the type-B ARRs contain long C-terminal extensions with a Myb-like DNA binding domain referred to as the GARP domain (Imamura et al., 1999; Hosoda et al., 2002). Multiple lines of evidence support the role of the type-B ARRs as transcription factors (Sakai et al., 2000; Imamura et al., 2001; Lohrmann et al., 2001; Sakai et al., 2001;



**Figure 5.** Phylogenetic Relationship of Receiver Domains from Response Regulators and Hybrid Kinases.

An unrooted bootstrapped tree is shown based on amino-acid alignment of receiver and pseudoreceiver domain sequences. The different families of receiver domains are highlighted, with subfamilies 1, 2 and 3 being indicated for the type-B ARRs. Note that the tree has some low bootstrap values indicating that regions of this tree should be considered to be simply radiations.

Hosoda et al., 2002; Imamura et al., 2003; Mason et al., 2004; Mason et al., 2005; Rashotte et al., 2006). Type-B ARRs are nuclear-localized in Arabidopsis and capable of transcriptional activation when expressed in yeast (Lohrmann et al., 1999; Sakai et al., 2000; Lohrmann et al., 2001). The ARR1, ARR2, and ARR10 type-B response regulators bind to a core DNA sequence 5'-(G/A)GAT(T/C)-3' (Sakai et al., 2000; Hosoda et al., 2002).

The eleven type-B ARRs of Arabidopsis fall into three subfamilies based on phylogenetic analysis: subfamily 1 contains seven

members (ARR1, ARR2, ARR10, ARR11, ARR12, ARR14, and ARR18); subfamily 2 contains two members (ARR13 and ARR21); and subfamily 3 is also comprised of two members (ARR19 and ARR20) (Mason et al., 2004). Genetic analyses indicate that at least five subfamily-1 members mediate cytokinin signaling, with ARR1, ARR10, and ARR12 appearing to play key roles (Mason et al., 2005; Yokoyama et al., 2007; Ishida et al., 2008). The cytokinin transcriptional response is substantially reduced in type-B mutant backgrounds, supporting a central role of the type-B ARRs

in the cytokinin signaling pathway (Rashotte et al., 2006; Yokoyama et al., 2007). A number of primary response genes directly regulated by the type-B ARR<sub>s</sub> have been identified, including the type-A ARR<sub>s</sub> (Taniguchi et al., 2007). It has been proposed that the subfamily-1 member ARR2 may modulate ethylene signaling, but the effect is subtle and differing results have been obtained in the analysis of *arr2* mutants, perhaps due to differing growth conditions (Hass et al., 2004; Mason et al., 2005).

The functions of the subfamily-2 and subfamily-3 type-B ARR<sub>s</sub> are unclear. They are not as broadly expressed as the subfamily-1 ARR<sub>s</sub> (Mason et al., 2004; Tajima et al., 2004), and the only reported mutant phenotypes arise from overexpression of activated versions of the genes (Tajima et al., 2004; Kiba et al., 2005). Overexpression of activated ARR21 (subfamily-2) results in seedlings in which cell proliferation is activated to form callus-like structures. Overexpression of activated ARR20 (subfamily-


**Table 3.** Response Regulators of Arabidopsis

Gene Name	Synonyms	Chrom Locus <sup>a</sup>	Features <sup>b</sup>	Family	References
ARR3		At1g59940	Rec	Type A	Imamura et al. 1998
ARR4	IBC7, ATR1	At1g10470	Rec	Type A	Imamura et al. 1998, Brandstatter and Kieber 1998, Urao et al. 1998
ARR5	IBC6, ATR2	At3g48100	Rec	Type A	Imamura et al. 1998, Brandstatter and Kieber 1998, Urao et al. 1998
ARR6		At5g62920	Rec	Type A	Imamura et al. 1998
ARR7		At1g19050	Rec	Type A	Imamura et al. 1998
ARR8	ATRR3	At2g41310	Rec	Type A	Imamura et al. 1999, Urao et al. 1998
ARR9	ATRR4	At3g57040	Rec	Type A	Imamura et al. 1999, Urao et al. 1998
ARR15		At1g74890	Rec	Type A	D'Agostino et al. 2000
ARR16		At2g40670	Rec	Type A	D'Agostino et al. 2000
ARR17		At3g56380	Rec	Type A	D'Agostino et al. 2000
ARR1		At3g16857	Rec, Myb	Type B	Sakai et al. 1998a
ARR2		At4g16110	Rec, Myb	Type B	Sakai et al. 1998a
ARR10	ARLP2	At4g31920	Rec, Myb	Type B	Imamura et al. 1999, Lohrmann et al. 1999
ARR11	ARLP1	At1g67710	Rec, Myb	Type B	Imamura et al. 1999, Lohrmann et al. 1999
ARR12		At2g25180	Rec, Myb	Type B	Imamura et al. 1999
ARR13		At2g27070	Rec, Myb	Type B	Imamura et al. 1999
ARR14		At2g01760	Rec, Myb	Type B	Imamura et al. 1999
ARR18		At5g58080	Rec, Myb	Type B	Schaller et al. 2002, Mason et al. 2004, Tajima et al. 2004
ARR19		At1g49190	Rec, Myb	Type B	Schaller et al. 2002, Mason et al. 2004, Tajima et al. 2004
ARR20		At3g62670	Rec, Myb	Type B	Schaller et al. 2002, Mason et al. 2004, Tajima et al. 2004
ARR21		At5g07210	Rec, Myb	Type B	Schaller et al. 2002, Mason et al. 2004, Tajima et al. 2004
ARR22		At3g04280	Rec	Type C	Schaller et al. 2002, Kiba et al., 2004
ARR23		At5g62120	missing 5' end		Schaller et al. 2002
ARR24		At5g26594	Rec	Type C	Gattolin et al., 2006
TOC1	PRR1, APRR1	At5g61380	Pseudo-Rec, CCT	Clock	Makino et al. 2000, Strayer et al. 2000
PRR3	APRR3	At5g60100	Pseudo-Rec, CCT	Clock	Matsushika et al. 2000
PRR5	APRR5	At5g24470	Pseudo-Rec, CCT	Clock	Matsushika et al. 2000
PRR7	APRR7	At5g02810	Pseudo-Rec, CCT	Clock	Matsushika et al. 2000
PRR9	APRR9	At2g46790	Pseudo-Rec, CCT	Clock	Matsushika et al. 2000
PRR2	APRR2	At4g18020	Pseudo-Rec, Myb		Makino et al. 2000
PRR4	APRR4	At5g49240	Pseudo-Rec, Myb		Schaller et al. 2002
PRR6	APRR6	At1g68210	Pseudo-Rec, Myb		Schaller et al. 2002
PRR8	APRR8	At4g00760	Pseudo		Schaller et al. 2002

<sup>a</sup>Chromosome loci are given by the MIPS designation.

<sup>b</sup>Features include receiver domain (Rec), pseudo-receiver domain lacking the phosphorylatable Asp (Pseudo-Rec), Myb-like DNA binding domain (Myb), CCT motif found in clock proteins, and whether the gene is missing essential coding sequence for a receiver.





**Figure 6.** Cytokinin Signal Transduction Occurs Through a Multi-Step Phosphorelay.

Cytokinin receptors, AHPs, and type-B ARRs function as a positive regulatory circuit to relay the cytokinin signal from plasma membrane to nucleus. Type-B ARRs act as transcription factors, one target being genes encoding type-A ARRs. The type-A ARRs feed back to inhibit their own transcription and may also mediate other cytokinin responses. The roles of the listed cytokinin receptors, AHPs, type-B response regulators, and type-A response regulators in cytokinin signal transduction have each been confirmed by the analysis of T-DNA insertion mutations.

3) results in plants that develop small flowers and abnormal siliques with reduced fertility.

### Type-C response regulators

Arabidopsis also contains two additional response regulators (ARR22 and 24). These lack long C-terminal extensions, like the type-A ARRs, but are not closely related to the type-A ARRs based on phylogenetic analysis (Figure 5) (Kiba et al., 2004; Schaller et al., 2007). The type-C ARRs are also not transcriptionally regulated by cytokinin. The type-C ARR sequences are more similar to the hybrid-kinase receiver domains than to other response regulators (Figure 5), raising the possibility that a histidine kinase, rather than an HPT protein, could serve as their phosphodonor. *ARR22* and *ARR24* are predominantly expressed in flowers and siliques (Kiba et al., 2004; Gattolin et al., 2006). Single and double loss-of-function mutants grow similarly to wild-type (Gattolin et al., 2006). Overexpression of *ARR22* inhibits cytokinin signaling based on the transgenic lines having reduced shoot growth, poor root development, reduced cytokinin-responsive gene induction, and insensitivity to cytokinin under conditions for callus production (Kiba et al., 2004; Kiba et al., 2005; Gattolin et al., 2006). Whether the type-C ARRs normally antagonize the cytokinin signaling pathway is not known.

### Pseudo-response regulators

Arabidopsis contains nine pseudo-response regulators termed PRRs (Table 3, Figures 2 and 5). These contain complete receiver domains but are missing essential residues required for activity (Makino et al., 2000). In particular, the aspartate that serves as a site for phosphorylation is missing, in many cases being replaced by a glutamate residue that may mimic the phosphorylated form.

These also contain C-terminal extensions, some members with a CCT-motif (PRR1/TOC1, PRR3, PRR5, PRR7, and PRR9) and others with the Myb-like motif found in the type-B response regulators (PRR2, PRR4, and PRR6).

The pseudoresponse-regulators with the CCT motif are involved in the regulation of circadian rhythms and participate in multiple regulatory feedback loops in recent models for the Arabidopsis clock (Mizuno, 2005; Gardner et al., 2006; McClung, 2006). *PRR1/TOC1* was identified in a forward genetic screen due to a semi-dominant mutation (*timing of cab expression 1-1*) that had shortened circadian periods for leaf movement, stomatal conductance, and several molecular markers (Millar et al., 1995; Somers et al., 1998; Strayer et al., 2000). *PRR1/TOC1* participates in a feedback loop involving the other well-known clock components, CCA1 and LHY (Alabadi et al., 2001). Evidence that all the CCT-motif PRRs participate in the clock came from the discovery that their expression varies in a circadian manner (Matsushika et al., 2000; Makino et al., 2001) and that loss-of-function mutants have altered circadian periods (Kaczorowski and Quail, 2003; Michael et al., 2003; Farré et al., 2005; Nakamichi et al., 2005b; Nakamichi et al., 2005a; Salomé and McClung, 2005; Ito et al., 2008). The current model places PRR5, PRR7, and PRR9 in a feedback loop that also involves CCA1 and LHY. Recent work indicates that protein stability of TOC1, PRR5, PRR7, and PRR9 is post-translationally regulated (Farré and Kay, 2007; Ito et al., 2007; Kiba et al., 2007; Para et al., 2007). Additionally, PRR7 is phosphorylated, presumably on serine/threonine residues since it lacks the canonical phospho-acceptor aspartate, which may serve as a means to regulate its stability and/or function (Farré and Kay, 2007).

### Potential Pseudogenes

Arabidopsis also contains the predicted sequence for a response regulator (ARR23) that, although containing the phosphorylated

aspartate, is predicted to lack the N-terminal domain of the receiver; no EST is reported and *ARR23* could be a pseudogene. The gene product of At3g04270 also shows homology to receiver domains but is missing what would be its C-terminal end.

## SIGNALING THROUGH PHOSPHORELAYS

A multi-step phosphorelay, rather than a simple two-component system, appears to be the major His-Asp signaling circuit employed by Arabidopsis. This possibility was initially raised by analysis of the Arabidopsis genome, which revealed a preponderance of hybrid kinases in Arabidopsis along with the presence of HPT proteins (Schaller et al., 2002). Subsequent analyses of protein-protein interactions using the yeast two-hybrid system supports interactions among the hybrid kinases with HPT proteins, and of the HPT proteins with both type-A and type-B response regulators, consistent with what would be expected in a multi-step phosphorelay (Imamura et al., 1999; Urao et al., 2000; Dortay et al., 2007). Finally, as described below, analysis of the cytokinin signaling pathway supports use of the multi-step phosphorelay *in planta*.

The cytokinin signaling pathway gives us our clearest picture of how two-component signaling elements have been adapted to signaling in plants (Hwang and Sheen, 2001; Haberer and Kieber, 2002; Heyl and Schmölling, 2003; Kakimoto, 2003; To and Kieber, 2008). Genetic analyses using loss-of-function mutations indicate that the primary cytokinin signaling pathway is a positive regulatory circuit that requires hybrid histidine kinases (AHK2, AHK3, and AHK4) (Inoue et al., 2001; Suzuki et al., 2001; Ueguchi et al., 2001; Yamada et al., 2001; Kakimoto, 2003; Kim et al., 2006), HPT proteins (AHP1, AHP2, AHP3, AHP5) (Hutchison et al., 2006), and type-B response regulators (ARR1, ARR2, ARR10, ARR11, and ARR12) (Sakai et al., 2001; Mason et al., 2005; Yokoyama et al., 2007; Ishida et al., 2008) (Figure 6). The function of type-B ARRs as transcription factors indicates that signals may move from histidine kinase to the nucleus solely through elements of a two-component system, the same type of signaling circuit employed by many prokaryotes. The mobile element of this signaling circuit are the AHPs, which relocate from cytosol to nucleus in response to cytokinin (Hwang and Sheen, 2001; Yamada et al., 2004). It should be noted that additional two-component signaling elements (such as other type-B ARRs) may also contribute to this cytokinin signaling circuit but this has not yet been confirmed genetically.

Several negative regulatory circuits are also employed in cytokinin signaling. First, AHK4, like some bacterial histidine kinases, has both kinase and phosphatase activities (Mähönen et al., 2006a). In the absence of cytokinin, AHK4 acts as a phosphatase to dephosphorylate AHPs, thereby decreasing signaling through the phosphorelay. Upon cytokinin binding, AHK4 switches to act as a histidine kinase to initiate the multi-step phosphorelay, resulting in phosphorylation of AHPs and downstream response regulators. Second, one of the initial transcriptional responses mediated by the type-B ARRs in response to cytokinin is to induce the expression of the type-A ARRs (Brandstatter and Kieber, 1998; Imamura et al., 1998; D'Agostino et al., 2000; Mason et al., 2005; Rashotte et al., 2006; Taniguchi et al., 2007; Yokoyama et al., 2007). The type-A ARRs then negatively regulate cytokinin signaling, potentially by competing with the type-B ARRs for phosphorylation by

the AHPs (Kiba et al., 2003; To et al., 2004; Leibfried et al., 2005; Lee et al., 2007; To et al., 2007). Furthermore, multiple type-A proteins are stabilized by cytokinin, further increasing this negative feedback loop (To et al., 2007). Third, the pseudo-HPT, *APHP1*, is induced by cytokinin and acts as a negative regulator of cytokinin responses (Mähönen et al., 2006b).

The presence of additional hybrid histidine-kinases in Arabidopsis, such as ETR1, AHK1, CKI1, and CKI2, suggests that multi-step phosphorelays function in relaying signals other than cytokinin. This raises the question as to which downstream components are involved in relaying the additional signals, given that a majority of the two-component signaling elements have already been implicated in cytokinin signaling. While there exists a formal possibility for unique combinations of signaling elements, it is far more likely that downstream signaling elements are shared among the receptors. In some cases, different input signals may be channeled into the same downstream signaling pathway to regulate a common response. Alternatively, a response tailored to a specific signal input could still be obtained even with shared signaling elements. One way to accomplish this would be to have varying affinities between the receptors and the AHPs, and between the AHPs and the response regulators, such that each signal would differentially activate the downstream components. Yeast two-hybrid analysis does suggest some specificity to the interactions between receptors and the AHPs (Urao et al., 2000), but the physiological relevance of these differences has not been determined. An alternative possibility would be for specificity of the downstream signaling elements to be modified by covalent modification or protein-protein interaction in ways unique to the signal input. It is also possible that protein subsets are sequestered in unique signaling complexes by scaffold proteins. Genetic, molecular, and proteomic approaches should clarify how this network of interactions functions in plants.

## ACKNOWLEDGEMENTS

We thank Dennis Mathews, Mike Gribskov, and John Walker for their assistance in writing the previous version of this review (Schaller et al., 2002). We thank Takeshi Mizuno and Tatsuo Kakimoto for their assistance with naming the two-component signaling elements, which resulted in the unified gene nomenclature adopted in 2002. Research in the authors' laboratories has been supported by grants from the National Science Foundation, Department of Energy, and the U. S. Department of Agriculture.

## REFERENCES

- Alabadi, D., Oyama, T., Yanovsky, M.J., Harmon, F.G., Mas, P., and Kay, S.A. (2001). Reciprocal regulation between TOC1 and LHY/CCA1 within the Arabidopsis circadian clock. *Science* **293**: 880-883.
- Anantharaman, V., and Aravind, L. (2001). The CHASE domain: a predicted ligand-binding module in plant cytokinin receptors and other eukaryotic and bacterial receptors. *Trends Biochem. Sci.* **26**: 579-582.
- Appleby, J.L., Parkinson, J.S., and Bourret, R.B. (1996). Signal transduction via the multi-step phosphorelay: not necessarily a road less traveled. *Cell* **86**: 845-848.
- Baker, M.D., Wolanin, P.M., and Stock, J.B. (2006). Signal transduction in bacterial chemotaxis. *Bioessays* **28**: 9-22.
- Binder, B.M., O'Malley R, C., Wang, W., Moore, J.M., Parks, B.M., Spalding, E.P., and Bleecker, A.B. (2004). Arabidopsis seedling growth re-

- sponse and recovery to ethylene. A kinetic analysis. *Plant Physiol.* **136**: 2913-2920.
- Bleecker, A.B.** (1999). Ethylene perception and signaling: an evolutionary perspective. *Trends Plant Sci.* **4**: 269-274.
- Brandstatter, I., and Kieber, J.J.** (1998). Two genes with similarity to bacterial response regulators are rapidly and specifically induced by cytokinin in *Arabidopsis*. *Plant Cell* **10**: 1009-1019.
- Chang, C., Kwok, S.F., Bleecker, A.B., and Meyerowitz, E.M.** (1993). *Arabidopsis* ethylene response gene *ETR1*: Similarity of product to two-component regulators. *Science* **262**: 539-544.
- Chen, Y.F., Etheridge, N., and Schaller, G.E.** (2005). Ethylene signal transduction. *Ann. Bot. (Lond)* **95**: 901-915.
- Clack, T., Mathews, S., and Sharrock, R.A.** (1994). The phytochrome apoprotein family in *Arabidopsis* is encoded by five genes: the sequences and expression of PHYD and PHYE. *Plant Mol. Biol.* **25**: 413-427.
- D'Agostino, I.B., Deruere, J., and Kieber, J.J.** (2000). Characterization of the response of the *Arabidopsis* response regulator gene family to cytokinin. *Plant Physiol.* **124**: 1706-1717.
- Dortay, H., Mehnert, N., L., B., T., S., and A., H.** (2007). Analysis of protein interactions within the cytokinin-signaling pathway of *Arabidopsis thaliana*. *FEBS J.* **273**: 4631-4644.
- Farré, E.M., and Kay, S.A.** (2007). PRR7 protein levels are regulated by light and the circadian clock in *Arabidopsis*. *Plant J.* **52**: 548-560.
- Farré, E.M., Harmer, S.L., Harmon, F.G., Yanovsky, M.J., and Kay, S.A.** (2005). Overlapping and distinct roles of PRR7 and PRR9 in the *Arabidopsis* circadian clock. *Curr. Biol.* **15**: 47-54.
- Gamble, R.L., Coonfield, M.L., and Schaller, G.E.** (1998). Histidine kinase activity of the ETR1 ethylene receptor from *Arabidopsis*. *Proc. Natl. Acad. Sci. USA* **95**: 7825-7829.
- Gardner, M.J., Hubbard, K.E., Hotta, C.T., Dodd, A.N., and Webb, A.A.** (2006). How plants tell the time. *Biochem. J.* **397**: 15-24.
- Gattolin, S., Alandete-Saez, M., Elliot, K., Gonzalez-Carranza, Z., Naomab, E., Powell, C., and Roberts, J.A.** (2006). Spatial and temporal expression of the response regulators *ARR22* and *ARR24* in *Arabidopsis thaliana*. *J. Exp. Bot.* **57**: 4225-4233.
- Haberer, G., and Kieber, J.J.** (2002). Cytokinins. New insights into a classic phytohormone. *Plant Physiol.* **128**: 354-362.
- Hall, A.E., Findell, J.L., Schaller, G.E., Sisler, E.C., and Bleecker, A.B.** (2000). Ethylene perception by the ERS1 protein in *Arabidopsis*. *Plant Physiol.* **123**: 1449-1458.
- Hass, C., Lohrmann, J., Albrecht, V., Sweere, U., Hummel, F., Yoo, S.D., Hwang, I., Zhu, T., Schafer, E., Kudla, J., and Harter, K.** (2004). The response regulator 2 mediates ethylene signalling and hormone signal integration in *Arabidopsis*. *EMBO J.* **23**: 3290-3302.
- Hejatko, J., Pernisova, M., Eneva, T., Palme, K., and Brzobohaty, B.** (2003). The putative sensor histidine kinase CK11 is involved in female gametophyte development in *Arabidopsis*. *Mol. Genet. Genomics* **269**: 443-453.
- Heyl, A., and Schmölling, T.** (2003). Cytokinin signal perception and transduction. *Curr. Opin. Plant Biol.* **6**: 480-488.
- Heyl, A., Wulfetange, K., Pils, B., Nielsen, N., Romanov, G.A., and T.D., S.** (2007). Evolutionary proteomics identifies amino acids essential for ligand-binding of the cytokinin receptor CHASE domain. *BMC Evol. Biol.* **7**: 62.
- Higuchi, M., Pischke, M.S., Mahonen, A.P., Miyawaki, K., Hashimoto, Y., Seki, M., Kobayashi, M., Shinozaki, K., Kato, T., Tabata, S., Helariutta, Y., Sussman, M.R., and Kakimoto, T.** (2004). In planta functions of the *Arabidopsis* cytokinin receptor family. *Proc. Natl. Acad. Sci. USA* **101**: 8821-8826.
- Hosoda, K., Imamura, A., Katoh, E., Hatta, T., Tachiki, M., Yamada, H., Mizuno, T., and Yamazaki, T.** (2002). Molecular structure of the GARP family of plant Myb-related DNA binding motifs of the *Arabidopsis* response regulators. *Plant Cell* **14**: 2015-2029.
- Hua, J., Chang, C., Sun, Q., and Meyerowitz, E.M.** (1995). Ethylene sensitivity conferred by *Arabidopsis* *ERS* gene. *Science* **269**: 1712-1714.
- Hua, J., Sakai, H., Nourizadeh, S., Chen, Q.G., Bleecker, A.B., Ecker, J.R., and Meyerowitz, E.M.** (1998). Ein4 and ERS2 are members of the putative ethylene receptor family in *Arabidopsis*. *Plant Cell* **10**: 1321-1332.
- Hunter, T.** (1995). Protein kinases and phosphatases: the yin and yang of protein phosphorylation and signaling. *Cell* **80**: 225-236.
- Hunter, T., and Plowman, G.D.** (1997). The protein kinases of budding yeast: six score and more. *Trends Biochem. Sci.* **22**: 18-22.
- Hutchison, C.E., Li, J., Argueso, C., Gonzalez, M., Lee, E., Lewis, M.W., Maxwell, B.B., Perdue, T.D., Schaller, G.E., Alonso, J.M., Ecker, J.R., and Kieber, J.J.** (2006). The *Arabidopsis* histidine phosphotransfer proteins are redundant positive regulators of cytokinin signaling. *Plant Cell* **18**: 3073-3087.
- Hwang, I., and Sheen, J.** (2001). Two-component circuitry in *Arabidopsis* cytokinin signal transduction. *Nature* **413**: 383-389.
- Imamura, A., Yoshino, Y., and Mizuno, T.** (2001). Cellular localization of the signaling components of *Arabidopsis* His-to-Asp phosphorelay. *Biosci. Biotechnol. Biochem.* **65**: 2113-2117.
- Imamura, A., Kiba, T., Tajima, Y., Yamashino, T., and Mizuno, T.** (2003). In vivo and in vitro characterization of the ARR11 response regulator implicated in the His-to-Asp phosphorelay signal transduction in *Arabidopsis thaliana*. *Plant Cell Physiol.* **44**: 122-131.
- Imamura, A., Hanaki, N., Umeda, H., Nakamura, A., Suzuki, T., Ueguchi, C., and Mizuno, T.** (1998). Response regulators implicated in his-to-asp phosphotransfer signaling in *Arabidopsis*. *Proc. Natl. Acad. Sci. USA* **95**: 2691-2696.
- Imamura, A., Hanaki, N., Nakamura, A., Suzuki, T., Taniguchi, M., Kiba, T., Ueguchi, C., Sugiyama, T., and Mizuno, T.** (1999). Compilation and characterization of *Arabidopsis thaliana* response regulators implicated in His-Asp phosphorelay signal transduction. *Plant Cell Physiol.* **40**: 733-742.
- Inoue, T., Higuchi, M., Hashimoto, Y., Seki, M., Kobayashi, M., Kato, T., Tabata, S., Shinozaki, K., and Kakimoto, T.** (2001). Identification of CRE1 as a cytokinin receptor from *Arabidopsis*. *Nature* **409**: 1060-1063.
- Ishida, K., Yamashino, T., Yokoyama, A., and Mizuno, T.** (2008). Three type-B response regulators, ARR1, ARR10, and ARR12, play essential but redundant roles in cytokinin signal transduction throughout the life cycle of *Arabidopsis thaliana*. *Plant Cell Physiol.* **49**: 47-57.
- Ito, S., Nakamichi, N., Kiba, T., Yamashino, T., and Mizuno, T.** (2007). Rhythmic and light-inducible appearance of clock-associated pseudo-response regulator protein PRR9 through programmed degradation in the dark in *Arabidopsis thaliana*. *Plant Cell Physiol.* **48**: 1644-1651.
- Ito, S., Niwa, Y., Nakamichi, N., Kawamura, H., Yamashino, T., and Mizuno, T.** (2008). Insight into missing genetic links between two evening-expressed pseudo-response regulator genes TOC1 and PRR5 in the circadian clock-controlled circuitry in *Arabidopsis thaliana*. *Plant Cell Physiol.* **49**: 201-213.
- Iwami, A., Yamashino, T., Tanaka, Y., Sakakibara, H., Kakimoto, T., Sato, S., Kato, T., Tabata, S., Nagatani, A., and Mizuno, T.** (2007). AHK5 histidine kinase regulates root elongation through an ETR1-dependent abscisic acid and ethylene signaling pathway in *Arabidopsis thaliana*. *Plant Cell Physiol.* **48**: 375-380.
- Kaczorowski, K.A., and Quail, P.H.** (2003). *Arabidopsis* PSEUDO-RESPONSE REGULATOR7 is a signaling intermediate in phytochrome-regulated seedling deetiolation and phasing of the circadian clock. *Plant Cell* **15**: 2654-2665.
- Kakimoto, T.** (1996). CK11, a histidine kinase homologue involved in cytokinin signal transduction. *Science* **274**: 982-985.
- Kakimoto, T.** (2003). Perception and signal transduction of cytokinins. *Annu. Rev. Plant Biol.* **54**: 605-627.
- Kiba, T., Aoki, K., Sakakibara, H., and Mizuno, T.** (2004). *Arabidopsis* response regulator, ARR22, ectopic expression of which results in phenotypes similar to the wol cytokinin-receptor mutant. *Plant Cell Physiol.* **45**: 1063-1077.
- Kiba, T., Henriques, R., Sakakibara, H., and Chua, N.H.** (2007). Targeted degradation of PSEUDO-RESPONSE REGULATOR5 by an SCFZTL complex regulates clock function and photomorphogenesis in *Arabidopsis thaliana*. *Plant Cell* **19**: 2516-2530.
- Kiba, T., Naitou, T., Koizumi, N., Yamashino, T., Sakakibara, H., and Mizuno, T.** (2005). Combinatorial microarray analysis revealing *Arabidopsis* genes implicated in cytokinin responses through the His->Asp Phosphorelay circuitry. *Plant Cell Physiol.* **46**: 339-355.
- Kiba, T., Yamada, H., Sato, S., Kato, T., Tabata, S., Yamashino, T., and Mizuno, T.** (2003). The type-A response regulator, ARR15, acts as a negative regulator in the cytokinin-mediated signal transduction in *Arabidopsis thaliana*. *Plant Cell Physiol.* **44**: 868-874.

- Kim, H.J., Ryu, H., Hong, S.H., Woo, H.R., Lim, P.O., Lee, I.C., Sheen, J., Nam, H.G., and Hwang, I. (2006). Cytokinin-mediated control of leaf longevity by AHK3 through phosphorylation of ARR2 in *Arabidopsis*. *Proc. Natl. Acad. Sci. USA* **103**: 814-819.
- Lee, D.J., Park, J.Y., Ku, S.J., Ha, Y.M., Kim, S., Kim, M.D., Oh, M.H., and Kim, J. (2007). Genome-wide expression profiling of ARABIDOPSIS RESPONSE REGULATOR 7 (ARR7) overexpression in cytokinin response. *Mol. Genet. Genomics* **277**: 115-137.
- Leibfried, A., To, J.P.C., Stehling, S., Kehle, A., Busch, W., Demar, M., Kieber, J.J., and Lohmann, J.U. (2005). WUSCHEL controls meristem size by direct transcriptional regulation of cytokinin inducible response regulators. *Nature* **438**: 1172-1175.
- Lohrmann, J., Buchholz, G., Keitel, C., Sweere, U., Kircher, S., Bäurle, I., Kudla, J., Schäfer, E., and Harter, K. (1999). Differential expression and nuclear localization of response regulator-like proteins from *Arabidopsis thaliana*. *Plant. Biol.* **1**: 495-505.
- Lohrmann, J., Sweere, U., Zabaleta, E., Bäurle, I., Keitel, C., Kozma-Bognar, L., Brennicke, A., Schafer, E., Kudla, J., and Harter, K. (2001). The response regulator ARR2: a pollen-specific transcription factor involved in the expression of nuclear genes for components of mitochondrial complex I in *Arabidopsis*. *Mol. Genet. Genomics* **265**: 2-13.
- Loomis, W.F., Shaulsky, G., and Wang, N. (1997). Histidine kinases in signal transduction pathways of eukaryotes. *J. Cell Science* **110**: 1141-1145.
- Mähönen, A.P., Higuchi, M., Törmäkangas, K., Miyawaki, K., Pischke, M.S., Sussman, M.R., Helariutta, Y., and Kakimoto, T. (2006a). Cytokinins regulate a bidirectional phosphorelay network in *Arabidopsis*. *Curr. Biol.* **16**: 1116-1122.
- Mähönen, A.P., Bishopp, A., Higuchi, M., Nieminen, K.M., Kinoshita, K., Tormakangas, K., Ikeda, Y., Oka, A., Kakimoto, T., and Helariutta, Y. (2006b). Cytokinin signaling and its inhibitor AHP6 regulate cell fate during vascular development. *Science* **311**: 94-98.
- Makino, S., Matsushika, A., Kojima, M., Oda, Y., and Mizuno, T. (2001). Light response of the circadian waves of the APRR1/TOC1 quintet: when does the quintet start singing rhythmically in *Arabidopsis*? *Plant Cell Physiol.* **42**: 334-339.
- Makino, S., Kiba, T., Imamura, A., Hanaki, N., Nakamura, A., Suzuki, T., Taniguchi, M., Ueguchi, C., Sugiyama, T., and Mizuno, T. (2000). Genes encoding pseudo-response regulators: insight into His-to-Asp phosphorelay and circadian rhythm in *Arabidopsis thaliana*. *Plant Cell Physiol.* **41**: 791-803.
- Mason, M.G., Li, J., Mathews, D.E., Kieber, J.J., and Schaller, G.E. (2004). Type-B response regulators display overlapping expression patterns in *Arabidopsis*. *Plant Physiol.* **135**: 927-937.
- Mason, M.G., Mathews, D.E., Argyros, D.A., Maxwell, B.B., Kieber, J.J., Alonso, J.M., Ecker, J.R., and Schaller, G.E. (2005). Multiple type-B response regulators mediate cytokinin signal transduction in *Arabidopsis*. *Plant Cell* **17**: 3007-3018.
- Matsushika, A., Makino, S., Kojima, M., and Mizuno, T. (2000). Circadian waves of expression of the APRR1/TOC1 family of pseudo-response regulators in *Arabidopsis thaliana*: insight into the plant circadian clock. *Plant Cell Physiol.* **41**: 1002-1012.
- McClung, C.R. (2006). Plant circadian rhythms. *Plant Cell* **18**: 792-803.
- Michael, T.P., Salome, P.A., Yu, H.J., Spencer, T.R., Sharp, E.L., McPeck, M.A., Alonso, J.M., Ecker, J.R., and McClung, C.R. (2003). Enhanced fitness conferred by naturally occurring variation in the circadian clock. *Science* **302**: 1049-1053.
- Millar, A.J., Carré, I.A., Strayer, C.A., Chua, N.H., and Kay, S.A. (1995). Circadian clock mutants in *Arabidopsis* identified by luciferase imaging. *Science* **267**: 1161-1163.
- Mira-Rodado, V., Sweere, U., Grefen, C., Kunkel, T., Fejes, E., Nagy, F., Schäfer, E., and Harter, K. (2007). Functional cross-talk between two-component and phytochrome B signal transduction in *Arabidopsis*. *J. Exp. Bot.* **58**: 2595-2607.
- Miyata, S.-i., Urao, T., Yamaguchi-Shinozaki, K., and Shinozaki, K. (1998). Characterization of genes for two-component phosphorelay mediators with a single HPT domain in *Arabidopsis thaliana*. *FEBS Lett.* **437**: 11-14.
- Mizuno, T. (1997). Compilation of all genes encoding two-component phosphotransfer signal transducers in the genome of *Escherichia coli*. *DNA Res.* **4**: 161-168.
- Mizuno, T. (2005). Two-component phosphorelay signal transduction systems in plants: from hormone responses to circadian rhythms. *Biosci. Biotechnol. Biochem.* **69**: 2263-2276.
- Moussatche, P., and Klee, H.J. (2004). Autophosphorylation activity of the *Arabidopsis* ethylene receptor multigene family. *J. Biol. Chem.* **279**: 48734-48741.
- Nakamichi, N., Kita, M., Ito, S., Yamashino, T., and Mizuno, T. (2005a). PSEUDO-RESPONSE REGULATORS, PRR9, PRR7 and PRR5, together play essential roles close to the circadian clock of *Arabidopsis thaliana*. *Plant Cell Physiol.* **46**: 686-698.
- Nakamichi, N., Kita, M., Ito, S., Sato, E., Yamashino, T., and Mizuno, T. (2005b). The *Arabidopsis* pseudo-response regulators, PRR5 and PRR7, coordinately play essential roles for circadian clock function. *Plant Cell Physiol.* **46**: 609-619.
- Nishimura, C., Ohashi, Y., Sato, S., Kato, T., Tabata, S., and Ueguchi, C. (2004). Histidine kinase homologs that act as cytokinin receptors possess overlapping functions in the regulation of shoot and root growth in *Arabidopsis*. *Plant Cell* **16**: 1365-1377.
- Para, A., Farre, E.M., Imaizumi, T., Pruneda-Paz, J.L., Harmon, F.G., and Kay, S.A. (2007). PRR3 is a vascular regulator of TOC1 stability in the *Arabidopsis* circadian clock. *Plant Cell* **19**: 3462-3473.
- Parkinson, J.S. (1993). Signal transduction schemes of bacteria. *Cell* **73**: 857-871.
- Pischke, M.S., Jones, L.G., Otsuga, D., Fernandez, D.E., Drews, G.N., and Sussman, M.R. (2002). An *Arabidopsis* histidine kinase is essential for megagametogenesis. *Proc. Natl. Acad. Sci. USA* **99**: 15800-15805.
- Plowman, G.D., Sudarsanam, S., Bingham, J., Whyte, D., and Hunter, T. (1999). The protein kinases of *Caenorhabditis elegans*: a model for signal transduction in multicellular organisms. *Proc. Natl. Acad. Sci. USA* **96**: 13603-13610.
- Popov, K.M., Kedishvili, N.Y., Zhao, Y., Shimomura, Y., Crabb, D.W., and Harris, R.A. (1993). Primary structure of pyruvate dehydrogenase kinase establishes a new family of eukaryotic protein kinases. *J. Biol. Chem.* **268**: 26602-26606.
- Qu, X., and Schaller, G.E. (2004). Requirement of the histidine kinase domain for signal transduction by the ethylene receptor ETR1. *Plant Physiol.* **136**: 2961-2970.
- Rashotte, A.M., Mason, M.G., Hutchison, C.E., Ferreira, F.J., Schaller, G.E., and Kieber, J.J. (2006). A subset of *Arabidopsis* AP2 transcription factors mediates cytokinin responses in concert with a two-component pathway. *Proc. Natl. Acad. Sci. USA* **103**: 11081-11085.
- Reiser, V., Raitt, D.C., and Saito, H. (2003). Yeast osmosensor SLn1 and plant cytokinin receptor Cre1 respond to changes in turgor pressure. *J. Cell Biol.* **161**: 1035-1040.
- Riefler, M., Novak, O., Strnad, M., and Schumling, T. (2006). *Arabidopsis* cytokinin receptor mutants reveal functions in shoot growth, leaf senescence, seed size, germination, root development, and cytokinin metabolism. *Plant Cell* **18**: 40-54.
- Rockwell, N.C., Su, Y.-S., and Lagarias, J.C. (2006). Phytochrome structure and signaling mechanisms. *Annu. Rev. Plant Biol.* **57**: 837-858.
- Rodriguez, F.I., Esch, J.J., Hall, A.E., Binder, B.M., Schaller, G.E., and Bleeker, A.B. (1999). A copper cofactor for the ethylene receptor ETR1 from *Arabidopsis*. *Science* **283**: 996-998.
- Sakai, H., Aoyama, T., and Oka, A. (2000). *Arabidopsis* ARR1 and ARR2 response regulators operate as transcriptional activators. *Plant J.* **24**: 703-711.
- Sakai, H., Hwa, J., Chen, Q.G., Chang, C., Medrano, L.J., Bleeker, A.B., and Meyerowitz, E.M. (1998). *ETR2* is an *ETR1*-like gene involved in ethylene signaling in *Arabidopsis*. *Proc. Natl. Acad. Sci. USA* **95**: 5812-5817.
- Sakai, H., Honma, T., Aoyama, T., Sato, S., Kato, T., Tabata, S., and Oka, A. (2001). ARR1, a transcription factor for genes immediately responsive to cytokinins. *Science* **294**: 1519-1521.
- Salomé, P.A., and McClung, C.R. (2005). PSEUDO-RESPONSE REGULATOR 7 and 9 are partially redundant genes essential for the temperature responsiveness of the *Arabidopsis* circadian clock. *Plant Cell* **17**: 791-803.
- Salomé, P.A., To, J.P., Kieber, J.J., and McClung, C.R. (2006). *Arabidopsis* response regulators ARR3 and ARR4 play cytokinin-independent roles in the control of circadian period. *Plant Cell* **18**: 55-69.

- Schaller, G.E.** (2000). Histidine kinases and the role of two-component systems in plants. *Adv. Bot. Res.* **32**: 109-148.
- Schaller, G.E., and Bleecker, A.B.** (1995). Ethylene-binding sites generated in yeast expressing the Arabidopsis *ETR1* gene. *Science* **270**: 1809-1811.
- Schaller, G.E., Mathews, D.E., Gribskov, M., and Walker, J.C.** (2002). Two-component signaling elements and histidyl-aspartyl phosphorelays. *The Arabidopsis Book* (C. Somerville, E. Meyerowitz, editors), 1-9.
- Schaller, G.E., Doi, K., Hwang, I., Kieber, J.J., Khurana, J.P., Kurata, N., Mizuno, T., Pareek, A., Shiu, S.-H., Wu, P., and Yip, W.K.** (2007). Letter to the Editor: Nomenclature for two-component signaling elements of *Oryza sativa*. *Plant Physiol.* **143**: 555-557.
- Schultz, J., Copley, R.R., Doerks, T., Ponting, C.P., and Bork, P.** (2000). SMART: a web-based tool for the study of genetically mobile domains. *Nucleic Acids Res.* **28**: 231-234.
- Sharrock, R.A., and Quail, P.H.** (1989). Novel phytochrome sequences in Arabidopsis thaliana: structure, evolution, and differential expression of a plant regulatory photoreceptor family. *Genes Dev.* **3**: 1745-1757.
- Somers, D.E., Webb, A.A., Pearson, M., and Kay, S.A.** (1998). The short-period mutant, *toc1-1*, alters circadian clock regulation of multiple outputs throughout development in Arabidopsis thaliana. *Development* **125**: 485-494.
- Stock, A.M., Robinson, V.L., and Goudreau, P.N.** (2000). Two-component signal transduction. *Annu. Rev. Biochem.* **69**: 183-215.
- Strayer, C., Oyama, T., Schultz, T.F., Raman, R., Somers, D.E., Mas, P., Panda, S., Kreps, J.A., and Kay, S.A.** (2000). Cloning of the Arabidopsis clock gene *TOC1*, an autoregulatory response regulator homolog. *Science* **289**: 768-771.
- Suzuki, T., Imamura, A., Ueguchi, C., and Mizuno, T.** (1998). Histidine-containing phosphotransfer (HPT) signal transducers implicated in His-to-Asp phosphorelay in Arabidopsis. *Plant Cell Physiol.* **39**: 1258-1268.
- Suzuki, T., Zakurai, K., Imamura, A., Nakamura, A., Ueguchi, C., and Mizuno, T.** (2000). Compilation and characterization of histidine-containing phosphotransmitters implicated in His-to-Asp phosphorelay in plants: AHP signal transducers of *Arabidopsis thaliana*. *Biosci. Biotechnol. Biochem.* **64**: 2482-2485.
- Suzuki, T., Miwa, K., Ishikawa, K., Yamada, H., Aiba, H., and Mizuno, T.** (2001). The arabidopsis sensor his-kinase, *ahk4*, can respond to cytokinins. *Plant Cell Physiol.* **42**: 107-113.
- Swanson, R.V., Alex, L.A., and Simon, M.I.** (1994). Histidine and aspartate phosphorylation: two-component systems and the limits of homology. *Trends Biochem.* **19**: 485-490.
- Sweere, U., Eichenberg, K., Lohrmann, J., Mira-Rodado, V., Baurle, I., Kudla, J., Nagy, F., Schafer, E., and Harter, K.** (2001). Interaction of the response regulator *ARR4* with phytochrome B in modulating red light signaling. *Science* **294**: 1108-1111.
- Tajima, Y., Imamura, A., Kiba, T., Amano, Y., Yamashino, T., and Mizuno, T.** (2004). Comparative Studies on the Type-B Response Regulators Revealing their Distinctive Properties in the His-to-Asp Phosphorelay Signal Transduction of Arabidopsis thaliana. *Plant Cell Physiol.* **45**: 28-39.
- Tanaka, Y., Suzuki, T., Yamashino, T., and Mizuno, T.** (2004). Comparative studies of the AHP histidine-containing phosphotransmitters implicated in His-to-Asp phosphorelay in *Arabidopsis thaliana*. *Biosci. Biotechnol. Biochem.* **68**: 462-465.
- Taniguchi, M., Sasaki, N., Tsuge, T., Aoyama, T., and Oka, A.** (2007). *ARR1* directly activates cytokinin response genes that encode proteins with diverse regulatory functions. *Plant Cell Physiol.* **48**: 263-277.
- Taniguchi, M., Kiba, T., Sakakibara, H., Ueguchi, C., Mizuno, T., and Sugiyama, T.** (1998). Expression of Arabidopsis response regulator homologs is induced by cytokinins and nitrate. *FEBS Lett.* **429**: 259-262.
- Thelen, J.J., Miernyk, J.A., and Randall, D.D.** (2000). Pyruvate dehydrogenase kinase from Arabidopsis thaliana: a protein histidine kinase that phosphorylates serine residues. *Biochem J.* **349**: 195-201.
- To, J.P., Haberer, G., Ferreira, F.J., Duere, J., Mason, M.G., Schaller, G.E., Alonso, J.M., Ecker, J.R., and Kieber, J.J.** (2004). Type-A Arabidopsis response regulators are partially redundant negative regulators of cytokinin signaling. *Plant Cell* **16**: 658-671.
- To, J.P.C., and Kieber, J.J.** (2008). Cytokinin signaling: two-components and more. *Trends Plant Sci.* **13**: 85-92.
- To, J.P.C., Duere, J., Maxwell, B.B., Morris, V.F., Hutchison, C.E., Ferreira, F.J., Schaller, G.E., and Kieber, J.J.** (2007). Cytokinin regulates type-A Arabidopsis response regulator activity and protein stability via two-component phosphorelay. *Plant Cell* **19**: 3901-3914.
- Tran, L.-S.P., Urao, T., Qin, F., Maruyama, K., Kakimoto, T., Shinozaki, K., and Yamaguchi-Shinozaki, K.** (2007). Functional analysis of *AHK1/ATHK1* and cytokinin receptor histidine kinases in response to abscisic acid, drought, and salt stress in *Arabidopsis*. *Proc. Natl. Acad. Sci. USA* **104**: 20623-20628.
- Ueguchi, C., Sato, S., Kato, T., and Tabata, S.** (2001). The *AHK4* gene involved in the cytokinin-signaling pathway as a direct receptor molecule in Arabidopsis thaliana. *Plant Cell Physiol.* **42**: 751-755.
- Urao, T., Yakubov, B., Yamaguchi-Shinozaki, K., and Shinozaki, K.** (1998). Stress-responsive expression of genes for two-component response regulator-like proteins in *Arabidopsis thaliana*. *FEBS Lett.* **427**: 175-178.
- Urao, T., Miyata, S., Yamaguchi-Shinozaki, K., and Shinozaki, K.** (2000). Possible His to Asp phosphorelay signaling in an Arabidopsis two-component system. *FEBS Lett.* **478**: 227-232.
- Urao, T., Yakubov, B., Satoh, R., Yamaguchi-Shinozaki, K., Seki, M., Hirayama, T., and Shinozaki, K.** (1999). A transmembrane hybrid-type histidine kinase in Arabidopsis functions as an osmosensor. *Plant Cell* **11**: 1743-1754.
- Wang, W., Hall, A.E., O'Malley, R., and Bleecker, A.B.** (2003). Canonical histidine kinase activity of the transmitter domain of the *ETR1* ethylene receptor from Arabidopsis is not required for signal transmission. *Proc. Natl. Acad. Sci. USA* **100**: 352-357.
- Yamada, H., Koizumi, N., Nakamichi, N., Kiba, T., Yamashino, T., and Mizuno, T.** (2004). Rapid response of *Arabidopsis* T87 cultured cells to cytokinin through His-to-Asp phosphorelay signal transduction. *Biosci. Biotechnol. Biochem.* **68**: 1966-1976.
- Yamada, H., Suzuki, T., Terada, K., Takei, K., Ishikawa, K., Miwa, K., and Mizuno, T.** (2001). The Arabidopsis *AHK4* histidine kinase is a cytokinin-binding receptor that transduces cytokinin signals across the membrane. *Plant Cell Physiol.* **42**: 1017-1023.
- Yeh, K.-C., and Lagarias, J.C.** (1998). Eukaryotic phytochromes: Light-regulated serine/threonine protein kinases with histidine kinase ancestry. *Proc. Natl. Acad. Sci. USA* **95**: 13976-13981.
- Yokoyama, A., Yamashino, T., Amano, Y., Tajima, Y., Imamura, A., Sakakibara, H., and Mizuno, T.** (2007). Type-B ARR Transcription Factors, *ARR10* and *ARR12*, are Implicated in Cytokinin-Mediated Regulation of Protoxylem Differentiation in Roots of Arabidopsis thaliana. *Plant Cell Physiol.* **48**: 84-96.