

## **FORTY-FOURTH SUPPLEMENT TO THE AMERICAN ORNITHOLOGISTS' UNION CHECK-LIST OF NORTH AMERICAN BIRDS**

Authors: Banks, Richard C., Cicero, Carla, Dunn, Jon L., Kratter, Andrew W., Rasmussen, Pamela C., et al.

Source: The Auk, 120(3) : 923-931

Published By: American Ornithological Society

URL: [https://doi.org/10.1642/0004-8038\(2003\)120\[0923:FSTTAO\]2.0.CO;2](https://doi.org/10.1642/0004-8038(2003)120[0923:FSTTAO]2.0.CO;2)

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## FORTY-FOURTH SUPPLEMENT TO THE AMERICAN ORNITHOLOGISTS' UNION CHECK-LIST OF NORTH AMERICAN BIRDS

RICHARD C. BANKS,<sup>1,9,10</sup> CARLA CICERO,<sup>2</sup> JON L. DUNN,<sup>3</sup> ANDREW W. KRATTER,<sup>4</sup> PAMELA  
C. RASMUSSEN,<sup>5</sup> J. V. REMSEN, JR.,<sup>6</sup> JAMES D. RISING,<sup>7</sup> AND DOUGLAS F. STOTZ<sup>8</sup>

<sup>1</sup>U.S. Geological Survey, Patuxent Wildlife Research Center, National Museum of Natural History, MRC-111,  
Washington, DC 20560-0111, USA;

<sup>2</sup>Museum of Vertebrate Zoology, 3101 Valley Life Sciences Building, University of California,  
Berkeley, California 94720-3160, USA;

<sup>3</sup>RR2, Box 52R, Bishop, California 93514, USA;

<sup>4</sup>Florida Museum of Natural History, P. O. Box 117800, University of Florida, Gainesville, Florida 32611, USA;

<sup>5</sup>Michigan State University Museum, West Circle Drive, East Lansing, Michigan 48824-1045, USA;

<sup>6</sup>Museum of Natural Science, Louisiana State University, Foster Hall 119, Baton Rouge, Louisiana 70803, USA;

<sup>7</sup>Department of Zoology, Ramsay Wright Zoological Labs, University of Toronto,  
Toronto, Ontario M5S 3G5, Canada; and

<sup>8</sup>Environmental and Conservation Programs, Field Museum of Natural History, 1400 South Lake Shore Drive,  
Chicago, Illinois 60605-2496, USA

This is the third Supplement since publication of the 7<sup>th</sup> edition of the *Check-list of North American Birds* (American Ornithologists' Union [AOU] 1998). It summarizes decisions made by the AOU's Committee on Classification and Nomenclature between 1 January 2002 and 31 December 2002. The Committee has continued to operate in the manner outlined in the 42<sup>nd</sup> Supplement (AOU 2000), but will now publish Supplements annually so that changes accepted by the Committee may be publicized more quickly. Changes in this Supplement fall into the following categories: (1) one species is added to the main list because of splitting of a species previously on the list (*Loxia megaplaga*); (2) one species replaces another presently on the list because of splitting of an extralimital form (*Picoides dorsalis*); (3) two genera (*Euphonia* and *Chlorophonia*), with their 16 species in our area, are moved from the family Thraupidae and placed in the subfamily Euphoniinae in the Fringillidae; (4) three new generic names are inserted in the list because of splitting of genera previously included (*Patagioenas*, *Megascops*, and *Gymnoglaux*), with the consequent change in generic names of 21 species; (5) one genus is removed from the list (*Nyctea*) because of its merger with another on the list (*Bubo*), with the consequent change of the scientific name of one species; (6) two English names are changed without change in scientific name (Belcher's Gull and Rock Pigeon); (7) the

distribution of one species is changed because of the merger with it of an extralimital form (*Butorides striatus*); (8) one species is added to part 2 of the Appendix (*Oenoenas chiriquensis*); and (9) changes are made in the endings of 9 species names to bring them into conformity with the International Code of Zoological Nomenclature (see David and Gosselin 2002). In addition, several minor changes are made to correct citations of generic names or other errors. The addition to the main list brings the number of species recognized as occurring in the *Check-list* area (main list) to 2,031. Literature that provides the basis for the Committee's decisions is cited at the end of the Supplement, and citations not already in the Literature Cited of the 7<sup>th</sup> edition (with Supplements) become additions to it. An updated list of the bird species known from the AOU *Check-list* area may be accessed at <http://www.AOU.org/aou/birdlist.html>.

A significant decision by the Committee reflected in the list of species posted on the AOU web site but not yet in the text of the *Check-list* is the recognition of a major grouping of birds generally known as the Galloanseres and comprising the orders Anseriformes and Galliformes. Multiple lines of evidence show that the Galloanseres forms a sister group to the rest of the presently recognized Neognathae (p. 3); for a review see Cracraft and Clark (2001). Recognition of this group is based on immunological distances (Ho et al. 1976), amino-acid sequences from conservative alpha-crystallin genes (Caspers et al. 1997), DNA-DNA hybridization (Sibley and Ahlquist 1990), mitochondrial DNA gene sequences (Mindell et al. 1997, van Tuinen et al. 2000), nuclear gene sequences (Groth and Barrowclough 1999), and morphological characters (Dzerhinsky 1995, Livezey 1997, Cracraft 1998,

<sup>9</sup>Authors are members of the Committee on Classification and Nomenclature of the American Ornithologists' Union, listed alphabetically after the Chairman.

<sup>10</sup>E-mail: [banks.rc@nrmnh.si.edu](mailto:banks.rc@nrmnh.si.edu) or  
[richard\\_banks@usgs.gov](mailto:richard_banks@usgs.gov)

Cracraft and Clark 2001). We do not give formal nomenclatural recognition to this group at this time because of problems caused in the overall classification and because we anticipate that ongoing work in avian molecular genetics will result in additional changes in higher level classification. These changes will be incorporated into the next edition of the *Check-list* but cannot readily be incorporated into Supplements. The major present effect of recognition of the group Galloanseres is the move of the Anseriformes and Galliformes, in that sequence, from their present positions in the list of species on pp. xvii–liv to a position between the Tinamiformes and Gaviiformes. The rest of the ordinal sequence is unchanged.

A recent series of papers on genetic relationships of members of the nine-primaried oscines has shown that some species and genera long classified in established family groups actually, or probably, are more closely related to members of other family groups. It has long been recognized that the distinction between “tanagers” and “finches” is problematical; see Notes under Emberizidae in AOU 1998:591. Recent studies of mitochondrial DNA (Burns 1997, Burns et al. 2002, Klicka et al. 2000, García-Moreno et al. 2001, Lovette and Bermingham 2002, Yuri and Mindell 2002) indicate that some species normally considered to be members of the Thraupidae are actually more closely related to the Cardinalidae, Emberizidae, or Fringillidae. Other groups of species seem not to belong in the Thraupidae, but relationships with other families are not obvious. Further, some species traditionally placed in the Emberizidae or Parulidae may make these families paraphyletic or polyphyletic in respect to other families in the nine-primaried oscines. In many instances these studies complement and support earlier morphological work that questioned traditional placement but that were inconclusive. The combination of several lines of evidence often provide compelling arguments that species and genera are misplaced in the current classification, but do not provide equally compelling arguments about where they should be placed. The primary reason for this is the limited sampling of taxa either within the misplaced groups or among the potential recipient groups. Another related reason is that different studies may lead to different placement, partly because of differences in taxon sampling. These studies leave us with varying degrees of uncertainty about the relationships of groups that have been studied—not to mention those that have not yet been tested. We anticipate that additional studies, some already under way, will lead eventually to definitive answers to questions raised by past studies. Meantime, we are faced with several options. First, we can leave the present classification alone, continuing with a system we know to be flawed but that is at least familiar. Second, we can remove genera from families where

genetic data show that they do not belong and place them into a large and growing group of uncertain position (*incertae sedis*), which admits our ignorance but results in a mere list rather than a classification. Third, we can remove genera from families where they do not belong and place them tentatively in other families on the basis of genetic evidence, which risks an unstable classification that may change when more data become available. We have chosen what we believe is a middle ground, to retain the present sequence of families and species, but to mark those species that studies have shown or suggested should be transferred to another, but still indefinite, position. In the list of species on pp. xvii–liv of AOU (1998), and on the AOU web site, we suggest using the symbol \* to mark such species. When additional studies resolve the relationship of these problematical taxa, formal changes will be proposed and acted on.

The following changes to the 7<sup>th</sup> edition (page numbers refer thereto) result from the Committee’s actions:

pp. xvii–liv. In the list of bird species known from the *Check-list* area, change 2030 (from 43<sup>rd</sup> Supplement, Banks et. al. 2002) to 2031. Add to the Notes: The symbol \* indicates a species that is probably misplaced in the current phylogenetic listing, but for which data indicating proper placement are not yet available.

In the list, insert the following species in the proper position as indicated by the text of this Supplement:

*Gymnoglaux lawrencii* Bare-legged Owl.  
*Picoides dorsalis* American Three-toed Woodpecker.  
*Loxia megaplaga* Hispaniolan Crossbill.

Remove the following names:  
*Otus lawrencii* Cuban Screech-Owl.  
*Picoides tridactylus* Three-toed Woodpecker.

Change the following scientific names, with no change in English names:

*Neocrex colombianus* to *Neocrex colombiana*  
*Chlidonias hybridus* to *Chlidonias hybrida*  
*Columba cayennensis* to *Patagioenas cayennensis*  
*Columba speciosa* to *Patagioenas speciosa*  
*Columba squamosa* to *Patagioenas squamosa*  
*Columba leucocephala* to *Patagioenas leucocephala*  
*Columba flavirostris* to *Patagioenas flavirostris*  
*Columba inornata* to *Patagioenas inornata*  
*Columba fasciata* to *Patagioenas fasciata*  
*Columba caribaea* to *Patagioenas caribaea*  
*Columba subvinacea* to *Patagioenas subvinacea*  
*Columba nigriristris* to *Patagioenas nigriristris*  
*Otus kennicottii* to *Megascops kennicottii*  
*Otus asio* to *Megascops asio*  
*Otus seductus* to *Megascops seductus*

*Otus cooperi* to *Megascops cooperi*  
*Otus trichopsis* to *Megascops trichopsis*  
*Otus choliba* to *Megascops choliba*  
*Otus barbarus* to *Megascops barbarus*  
*Otus guatemalae* to *Megascops guatemalae*  
*Otus clarkii* to *Megascops clarkii*  
*Otus nudipes* to *Megascops nudipes*  
*Nyctea scandiaca* to *Bubo scandiacus*  
*Chaetura spinicauda* to *Chaetura spinicaudus*  
*Ornithion brunneicapillum* to *Ornithion brunnei-*  
*capillus*  
*Vireo atricapillus* to *Vireo atricapilla*  
*Poecile atricapilla* to *Poecile atricapillus*  
*Seiurus aurocapillus* to *Seiurus aurocapilla*  
*Chrysothlypis chrysomelaena* to *Chrysothlypis*  
*chrysomelas*

Change the following English names:

*Larus belcheri* Belcher's Gull.  
*Columba livia* Rock Pigeon.

Move the species in Anseriformes and Galliformes to a position immediately following those in the Tinamiformes.

Move the species from *Euphonia jamaica* through *Chlorophonia callophrys* to a position following *Fringilla montifringilla*, under the new heading *Euphoniinae*.

Change the following annotation:

*Gracula religiosa* Hill Myna. (I)

Add the symbol \* before each of the following names:

*Microligea palustris* Green-tailed Warbler.  
*Teretistris fernandinae* Yellow-headed Warbler.  
*Teretistris fornsi* Oriente Warbler.  
*Zeledonia coronata* Wrenthrush.  
*Icteria virens* Yellow-breasted Chat.  
*Granatellus venustus* Red-breasted Chat.  
*Granatellus sallaei* Gray-throated Chat.  
*Xenoligea montana* White-winged Warbler.  
*Coereba flaveola* Bananaquit.  
*Nesospingus speculiferus* Puerto Rican Tanager.  
*Chlorospingus ophthalmicus* Common Bush-Tanager.  
*Chlorospingus tacarcunae* Tacarcuna Bush-Tanager.  
*Chlorospingus inornatus* Pirre Bush-Tanager.  
*Chlorospingus pileatus* Sooty-capped Bush-Tanager.  
*Chlorospingus flavigularis* Yellow-throated Bush-Tanager.  
*Chlorospingus canigularis* Ashy-throated Bush-Tanager.  
*Phaenicophilus palmarum* Black-crowned Palm-Tanager.  
*Phaenicophilus poliocephalus* Gray-crowned Palm-Tanager.

*Calyptophilus tertius* Western Chat-Tanager.  
*Calyptophilus frugivorus* Eastern Chat-Tanager.  
*Rhodinocichla rosea* Rosy Thrush-Tanager.  
*Mitrospingus cassinii* Dusky-faced Tanager.  
*Chlorothraupis carmioli* Olive Tanager.  
*Chlorothraupis olivacea* Lemon-spectacled Tanager.  
*Habia rubica* Red-crowned Ant-Tanager.  
*Habia fuscicauda* Red-throated Ant-Tanager.  
*Habia atrimaxillaris* Black-cheeked Ant-Tanager.  
*Piranga roseogularis* Rose-throated Tanager.  
*Piranga flava* Hepatic Tanager.  
*Piranga rubra* Summer Tanager.  
*Piranga olivacea* Scarlet Tanager.  
*Piranga ludoviciana* Western Tanager.  
*Piranga bidentata* Flame-colored Tanager.  
*Piranga leucoptera* White-winged Tanager.  
*Piranga erythrocephala* Red-headed Tanager.  
*Spindalis zena* Western Spindalis.  
*Spindalis nigricephala* Jamaican Spindalis.  
*Spindalis dominicensis* Hispaniolan Spindalis.  
*Spindalis portoricensis* Puerto Rican Spindalis.  
*Volatinia jacarina* Blue-black Grassquit.  
*Sporophila schistacea* Slate-colored Seedeater.  
*Sporophila americana* Variable Seedeater.  
*Sporophila torqueola* White-collared Seedeater.  
*Sporophila nigricollis* Yellow-bellied Seedeater.  
*Sporophila minuta* Ruddy-breasted Seedeater.  
*Oryzoborus nuttingi* Nicaraguan Seed-Finch.  
*Oryzoborus funereus* Thick-billed Seed-Finch.  
*Amaurospiza concolor* Blue Seedeater.  
*Melopyrrha nigra* Cuban Bullfinch.  
*Tiaris canora* Cuban Grassquit.  
*Tiaris olivacea* Yellow-faced Grassquit.  
*Tiaris bicolor* Black-faced Grassquit.  
*Loxipasser anoxanthus* Yellow-shouldered Grassquit.  
*Loxigilla portoricensis* Puerto Rican Bullfinch.  
*Loxigilla violacea* Greater Antillean Bullfinch.  
*Loxigilla noctis* Lesser Antillean Bullfinch.  
*Euneornis campestris* Orangequit.  
*Melanoospiza richardsoni* St. Lucia Black Finch.  
*Pinaroloxias inornata* Cocos Finch.  
*Haplospiza rustica* Slaty Finch.  
*Acanthidops bairdii* Peg-billed Finch.  
*Diglossa baritula* Cinnamon-bellied Flowerpiercer.  
*Diglossa plumbea* Slaty Flowerpiercer.  
*Sicalis flaveola* Saffron Finch.  
*Sicalis luteola* Grassland Yellow-Finch.  
*Emberizoides herbicola* Wedge-tailed Grass-Finch.  
*Paroaria coronata* Red-crested Cardinal. (H, I)  
*Paroaria capitata* Yellow-billed Cardinal. (H, I)  
*Calcarius mccownii* McCown's Longspur.  
*Calcarius lapponicus* Lapland Longspur.  
*Calcarius pictus* Smith's Longspur.  
*Calcarius ornatus* Chestnut-collared Longspur.  
*Plectrophenax nivialis* Snow Bunting.  
*Plectrophenax hyperboreus* McKay's Bunting.  
*Saltator albicollis* Lesser Antillean Saltator.

*Saltator striatipectus* Streaked Saltator.  
*Saltator coerulescens* Grayish Saltator.  
*Saltator maximus* Buff-throated Saltator.  
*Saltator atriceps* Black-headed Saltator.  
*Saltator grossus* Slate-colored Grosbeak.

p. 16. In the account for *Pterodroma longirostris*, the California record should be 53 rather than 35 miles southwest of Point Reyes.

p. 45. *Butorides sundevalli* of the Galapagos Islands is considered to be conspecific with *B. striatus*, following Payne in Mayr and Cottrell (1979) and most other sources. In the Distribution section of *B. striatus*, insert "(striatus Group)" after the words *Resident* and *Wanders*. To the *Resident* paragraph, add: "and (*sundevalli* Group) in the Galapagos Islands." Change the last sentence of Notes to: Groups: *B. striatus* [Striated Heron] and *B. sundevalli* (Reichenow, 1877) [Lava Heron]. The latter Group is sometimes (e.g., Sibley and Monroe 1990) considered a distinct species. The extent of global variation in *B. striatus* suggests that more than one species may be involved.

p. 62. Remove the Notes section from *Cygnus olor* and place it at the end of the account for *Cygnus buccinator*. Change "the next three" to "the next two."

p. 108. Change the citation for *Planofalco*, in the synonymy of *Falco*, to: Oberholser, 1925, Amer. Midl. Nat. 9: 601, fn. Type, by original designation, *Falco mexicanus* Schlegel.

p. 135. *Neocrex colombianus* should be *N. colombiana* (fide David and Gosselin 2002).

p. 187. Change the English name of *Larus belcheri* from Band-tailed Gull to Belcher's Gull, a name parallel to that of Olrog's Gull for the sister species *L. atlanticus* and used for *L. belcheri* by Murphy (1936). Change the last sentence of the Notes to: Also known as Band-tailed Gull.

p. 205. *Chlidonias hybridus* should be *C. hybrida* (fide David and Gosselin 2002).

p. 218. In the synonymy of the genus *Columba*, *Ænoenas* should be *Cenoenas*.

p. 218. Change the English name of *Columba livia* to Rock Pigeon, to conform to the recent name change by the British Ornithologists' Union (1992), and modify the Notes accordingly.

On the basis of studies by Johnson and Clayton (2000) and Johnson et al. (2001) of nuclear and mitochondrial DNA, and a review of morphological (Ridgway 1916), serological (Cumley and Irwin 1944),

and behavioral (Johnston 1962) characters, we place New World pigeons formerly included in *Columba* in a separate genus, *Patagioenas* Reichenbach, 1853.

p. 218. After the account of *Columba livia*, insert a heading:

#### Genus *Patagioenas* Reichenbach

Follow this heading with the citations for the generic names *Patagioenas*, *Chloroenas*, *Lepidoenas*, and *Oenoenas* presently listed as synonyms under *Columba* and remove these citations from the synonymy of *Columba*.

Delete the Notes under the generic synonymy of *Columba* and insert the following after the synonymy of *Patagioenas*:

**Notes.**—For the use of *Oenoenas* as a distinct genus, see Johnston (1962); for a contrary opinion, see Corbin (1968). Reichenbach (1853) simultaneously provided three new generic names for American species of pigeon, as indicated above. The name *Patagioenas* was used first and has priority if *Chloroenas* and *Lepidoenas* are considered synonyms of it, as here and as implied by Johnson et al. (2001).

Change the headings for the remaining species now listed in *Columba* as follows, and change generic names and abbreviations in Notes accordingly:

*Patagioenas cayennensis* (Bonnaterre). Pale-vented Pigeon.  
*Patagioenas speciosa* (Gmelin). Scaled Pigeon.  
*Patagioenas squamosa* (Bonnaterre). Scaly-naped Pigeon.  
*Patagioenas leucocephala* (Linnaeus). White-crowned Pigeon.  
*Patagioenas flavirostris* (Wagler). Red-billed Pigeon.  
*Patagioenas inornata* (Vigors). Plain Pigeon.  
*Patagioenas fasciata* (Say). Band-tailed Pigeon.  
*Patagioenas caribaea* (Jacquin). Ring-tailed Pigeon.  
*Patagioenas subvinacea* (Lawrence). Ruddy Pigeon.  
*Patagioenas nigrirostris* (Sclater). Short-billed Pigeon.

p. 254. The subgenus *Megascops*, recognized for New World species of *Otus* except *O. flammeolus* (Marshall and King in Amadon and Bull 1988), is elevated to full generic status on the basis of mitochondrial DNA and vocal data (König et al. 1999). *Otus flammeolus* is retained within *Otus* because of vocal similarity with some Old World species.

After *Otus sunia*, insert:

#### Genus *Megascops* Kaup

*Megascops* Kaup, 1848, Isis 14:769. Type, by subsequent designation (Gray 1855), *Strix asio* Linnaeus.



Move the citation for *Gymnasio* from the synonymy of *Otus* (on p. 253) to the synonymy of *Megascops*. Add the following under the generic heading and synonymy:

**Notes.**—Formerly treated as a subgenus within *Otus* (Marshall and King in Amadon and Bull 1988), but mitochondrial DNA and vocal differences with Old World species indicate that generic status is warranted (König et al. 1999).

Change the headings for the following species now listed in *Otus* as follows, and change generic names and abbreviations in Notes accordingly:

*Megascops kennicottii* (Elliot). Western Screech-Owl.

*Megascops asio* (Linnaeus). Eastern Screech-Owl.

*Megascops seductus* (Moore). Balsas Screech-Owl.

*Megascops cooperi* (Ridgway). Pacific Screech-Owl.

*Megascops trichopsis* (Wagler). Whiskered Screech-Owl.

*Megascops choliba* (Vieillot). Tropical Screech-Owl.

*Megascops barbarus* (Sclater and Salvin). Bearded Screech-Owl.

*Megascops guatemalae* (Sharpe). Vermiculated Screech-Owl.

*Megascops clarki* (Kelso and Kelso). Bare-shanked Screech-Owl.

*Megascops nudipes* (Daudin). Puerto Rican Screech-Owl.

p. 256. In Notes under *Megascops guatemalae*, *M. atricapillus* should be *M. atricapilla* (fide David and Gosselin 2002).

p. 257. The monotypic genus *Gymnoglaux* is reinstated for *Otus lawrencii* on the basis of strong differences in morphology and vocal patterns, and because no justification was given for the merger of this species into *Otus*. Accordingly, the English name of this species is changed to Bare-legged Owl.

After *Otus nudipes*, insert the heading:

Genus *Gymnoglaux* Cabanis

Move the citation for the generic name from the top of p. 254, in synonymy of *Otus*.

Replace the species heading with:

*Gymnoglaux lawrencii* Sclater and Salvin. Bare-legged Owl.

Retain the species account for *Otus lawrencii* in 7<sup>th</sup> edition, but change Notes to: Formerly merged into *Otus*, following Marshall and King in Amadon and Bull (1988), as Cuban Screech-Owl, but separated on the basis of strong differences in morphology and vo-

cal patterns. Also known as Cuban Bare-legged Owl or Cuban Screech-Owl.

p. 258. The genus *Nyctea* is merged into *Bubo* on the basis of genetic studies (Wink and Heidrich 1999). Move the heading and citation for *Nyctea* to the synonymy of the genus *Bubo* on p. 257.

Change the species heading *Nyctea scandiaca* (Linnaeus) to *Bubo scandiacus* (Linnaeus).

Add the following to the account of *Bubo scandiacus*: **Notes.**—Former treatment of this species in the monotypic genus *Nyctea* was based on distinct plumage and weak osteological differences (Ford 1967). Genetic studies, however, indicate that it is closely related to *Bubo* (Sibley and Ahlquist 1990) and in fact is nested within the genus (Wink and Heidrich 1999). The specific name is an adjective and changes to agree with the gender of the generic name.

p. 274. Following Cleere (2002), the citation for the genus *Steatornis* should be changed to: Humboldt, 1814, in Humboldt and Bonpland, Voy. Nouv. Cont., Pt. 1, 1:416. The type species remains unchanged.

p. 278. *Chaetura spinicauda* should be *C. spinicaudus* (fide David and Gosselin 2002).

p. 341. New World and Old World populations of Three-toed Woodpeckers are split on the basis of differences in mitochondrial DNA (Zink et al. 1995, 2002) and voice (Winkler and Short 1978, Short 1982). Ridgway (1914) considered New World and Old World populations to be separate species, and the merger of New World *dorsalis* into Old World *tridactylus* (e.g., AOU 1931, Peters 1948) was never explained. Replace the account for *Picoides tridactylus* with the following:

*Picoides dorsalis* Baird. American Three-toed Woodpecker.

*Picoides dorsalis* Baird, 1858, in Baird, Cassin and Lawrence, Rep. Explor. and Surv. R. R. Pacific, vol. 9, pt. 2, xxviii, 97, 100. (Laramie Peak, Rocky Mountains = Albany County, Wyoming.)

**Habitat.**—Coniferous forest, mixed coniferous-deciduous forest, willows in riparian areas; favors areas with trees killed by fire or beetles.

**Distribution.**—As that listed for *P. tridactylus*, ending with "Nova Scotia," deleting the clause beginning "and in Eurasia . . ."

**Notes.**—Formerly considered conspecific with the Old World *P. tridactylus* (Linnaeus) [Eurasian Three-toed Woodpecker], but separated because of significant differences in mitochondrial DNA sequences (Zink et al. 1995, 2002) and call (Winkler and Short 1978, Short 1982).

p. 372. Add to the Notes under *Scytalopus panamensis*: Also known as Pale-throated Tapaculo.

p. 373. *Ornithion brunneicapillum* should be *O. brunneicapillus* (fide David and Gosselin 2002).

p. 432. *Vireo atricapilla* should be *V. atricapilla* (fide David and Gosselin 2002).

p. 463. *Poecile atricapilla* (as changed by AOU 2000) should be *P. atricapillus* (fide David and Gosselin 2002).

p. 508. In the account for *Turdus iliacus*, the date of the record at St. Anthony, Newfoundland, should be 1980 rather than 1950.

p. 554. *Seiurus aurocapillus* should be *S. aurocapilla* (fide David and Gosselin 2002).

p. 571. *Chrysothlypis chrysomelaena* should be *C. chrysomelas* (fide David and Gosselin 2002); remove Notes from the account.

p. 582–586, 659. Studies of mitochondrial DNA (Burns 1997, Klicka et al. 2000, Burns et al. 2002, Yuri and Mindell 2002) show some genera traditionally considered to be members of the Thraupidae are more closely related to members of other families. The genera *Euphonia* and *Chlorophonia*, always considered close to each other, are shown to fall well outside the limits of the Thraupidae and, among taxa sampled, closest to members of the Fringillidae (sensu AOU 1998). Because of incomplete sampling of species in the Fringillidae, placement of these genera within that family is uncertain. We resurrect the subfamily Euphoniinae (Cabanis 1847), previously used to separate these genera within the Thraupidae (Sclater 1886), and transfer it to the Fringillidae, where it is tentatively placed between the Fringillinae and Carduelinae.

Remove the genera *Euphonia* and *Chlorophonia*, and included species, from pages 582–586 and transfer them to a position in the Fringillidae on p. 659; see below.

p. 585. In the citation for the genus *Chlorophonia*, *Pipra cyanea* Vieillot should read *Pipra cyanea* Thunberg.

p. 659. After the account for *Fringilla montifringilla*, insert the following heading:

Subfamily EUPHONIINAE: Euphonioid Finches

Insert the accounts for the genera *Euphonia* and

*Chlorophonia*, and included species, from pages 582–586.

p. 663. Crossbills on the island of Hispaniola in the Greater Antilles are separated as a species on the basis of vocal and morphological differences that seem not to have been adequately considered when the species was merged with *Loxia leucoptera* many years ago (see Benkman 1994, Smith 1997). After the account for *Loxia curvirostra*, insert the following:

*Loxia megaplaga* Riley. Hispaniolan Crossbill.

*Loxia megaplaga* Riley, 1916, Smiths. Misc. Coll. 66, no. 15, p. 1. (El Rio, 4,000 feet, Santo Domingo [Dominican Republic].)

**Habitat.**—Pine forests.

**Distribution.**—*Resident* on Hispaniola, in the mountains of the Dominican Republic and the Massif de La Selle of southeastern Haiti.

**Notes.**—Formerly considered conspecific with *L. leucoptera*, but separated on the basis of vocal and morphological differences (Benkman 1994, Smith 1997).

In the account for *Loxia leucoptera*, delete the Greater Antilles portion of the breeding distribution and the Hispaniola portion of the winter distribution. Add the following sentence to the Notes: “Formerly included populations resident on Hispaniola, now separated as *L. megaplaga*.”

p. 697. In the heading and account for *Garrulax caeruleus*, change Laughing-thrush to Laughingthrush, to agree with use elsewhere in the text.

p. 699. The following species, discussed under *Columba nigriceps* in the 7<sup>th</sup> edition, is added to Part 2 of the Appendix. Insert the following after the account for *Larus nelsoni*:

*Oenoas chiriensis* Ridgway. Chiriqui Pigeon.

*Oenoas chiriensis* Ridgway, 1915, Proc. Biol. Soc. Wash. 28:139. (Chiriquí, Panama; Volcán de Chiriquí suggested by Deignan, Bull. U. S. Nat'l. Mus. 221, 1961.)

This species was based on a unique type. Conover (in Hellmayr and Conover 1942) suggested that the locality was in error and that the bird was *Columba purpureotincta* of the Guianas. Johnston (1962) believed it to be an aberrant individual of *C. subvinacea*, but Wetmore (1968) declared it to be *C. nigriceps*. A hybrid origin has not been ruled out. With the generic changes accepted above, the name would be *Patagioenas chiriensis*.

pp. 705–730. In the list of French Names of North American Birds, insert the following species in the proper position as indicated by the text of this Supplement:

*Picoides dorsalis* Pic à dos rayé  
*Loxia megalaga* Bec-croisé d'Hispaniola

Delete the entry for the following name:  
*Picoides tridactylus*

Change the following scientific names, with no change in French names:

*Neocrex colombianus* to *Neocrex colombiana*  
*Chlidonias hybridus* to *Chlidonias hybrida*  
*Columba cayennensis* to *Patagioenas cayennensis*  
*Columba speciosa* to *Patagioenas speciosa*  
*Columba squamosa* to *Patagioenas squamosa*  
*Columba leucocephala* to *Patagioenas leucocephala*  
*Columba flavirostris* to *Patagioenas flavirostris*  
*Columba inornata* to *Patagioenas inornata*  
*Columba fasciata* to *Patagioenas fasciata*  
*Columba caribaea* to *Patagioenas caribaea*  
*Columba subvinacea* to *Patagioenas subvinacea*  
*Columba nigristrostris* to *Patagioenas nigristrostris*  
*Otus kennicottii* to *Megascops kennicottii*  
*Otus asio* to *Megascops asio*  
*Otus seductus* to *Megascops seductus*  
*Otus cooperi* to *Megascops cooperi*  
*Otus trichopsis* to *Megascops trichopsis*  
*Otus choliba* to *Megascops choliba*  
*Otus barbarus* to *Megascops barbarus*  
*Otus guatemalae* to *Megascops guatemalae*  
*Otus clarkii* to *Megascops clarkii*  
*Otus nudipes* to *Megascops nudipes*  
*Otus lawrencii* to *Gymnoglaux lawrencii*  
*Nyctea scandiaca* to *Bubo scandiacus*  
*Chaetura spinicauda* to *Chaetura spinicaudus*  
*Ornithion brunneicapillum* to *Ornithion brunneicapillus*  
*Vireo atricapillus* to *Vireo atricapilla*  
*Poecile atricapilla* to *Poecile atricapillus*  
*Seiurus aurocapillus* to *Seiurus aurocapilla*  
*Chrysothlypis chrysomelaena* to *Chrysothlypis chrysomelas*

Change the French name of one entry as follows:  
*Rhytipterna holerythra* Tyran plaintif

Move the species in the Anatidae and in the Cracidae, Phasianidae, and Odontophoridae, in that sequence, to a position between the Tinamidae and Gaviidae.

Move the species from *Euphonia jamaica* through *Chlorophonia callophrys* to a position following *Fringilla montifringilla*.

Add the following to the list in Appendix, part 2:  
*Oenoenas chiriquensis* Pigeon du Chiriqui

Taxonomic proposals considered but not yet accepted by the committee include the transfer of the species *clamator* from the genus *Pseudoscops* to *Asio*, the merger of *Ciccaba* into *Strix*, and the division of *Ammodramus maritimus* into two (or more) species. We considered and rejected suggestions to change the English names of the Nazca Booby (*Sula granti*; see AOU 2000) and the prairie-chickens. Still under consideration is the proper placement of the species now considered *incertae sedis* between the Tyrannidae and Cotingidae, and the proper classification of genera and species of Tetraoninae. We are aware of reports in our area of several species not now on our list, but are awaiting consideration of these reports by our sister committee of the American Birding Association.

#### ACKNOWLEDGMENTS

M. Gosselin serves as the Committee's authority for French names, and N. David serves as authority for classical languages, especially relative to scientific names. N. Bahr, P. Davis, A. L. Edwards, D. D. Gibson, J. Heindel, S. N. G. Howell, I. Paulson, and T. S. Schulenberg either called matters to our attention or provided helpful advice, or both.

#### LITERATURE CITED

- AMADON, D., AND J. BULL. 1988. Hawks and owls of the world: A distributional and taxonomic list. *Proceedings of the Western Foundation of Vertebrate Zoology* 3:296–357.
- AMERICAN ORNITHOLOGISTS' UNION. 1931. *Check-list of North American Birds*. 4th edition. American Ornithologists' Union, Lancaster, Pennsylvania.
- AMERICAN ORNITHOLOGISTS' UNION. 1998. *Check-list of North American Birds*. 7th edition. American Ornithologists' Union, Washington, D.C.
- AMERICAN ORNITHOLOGISTS' UNION. 2000. Forty-second supplement to the American Ornithologists' Union *Check-list of North American Birds*. *Auk* 117:847–858.
- BANKS, R. C., C. CICERO, J. L. DUNN, A. W. KRATTER, P. C. RASMUSSEN, J. V. REMSEN, JR., J. D. RISING, AND D. F. STOTZ. 2002. Forty-third Supplement to the American Ornithologists' Union *Check-list of North American Birds*. *Auk* 119:897–906.
- BENKMAN, C. W. 1994. Comments on the ecology and status of the Hispaniolan Crossbill (*Loxia leucoptera megalaga*), with recommendations for its conservation. *Caribbean Journal of Science* 30:250–254.
- BRITISH ORNITHOLOGISTS' UNION. 1992. Checklist of



- Birds of Britain and Ireland. 6<sup>th</sup> ed. British Ornithologists' Union, Tring, Hertfordshire, United Kingdom.
- BURNS, K. J. 1997. Molecular systematics of tanagers (Thraupidae): Evolution and biogeography of a diverse radiation of Neotropical birds. *Molecular Phylogenetics and Evolution* 8: 334–348.
- BURNS, K. J., S. J. HACKETT, AND N. K. KLEIN. 2002. Phylogenetic relationships and morphological diversity in Darwin's finches and their relatives. *Evolution* 56:1240–1252.
- CABANIS, J. 1847. Ornithologische notizen. *Archiv für Naturgeschichte* 13:308–352.
- CASPERS, G.-J., D. U. DE WEERD, J. WATTEL, AND W. W. DE JONG. 1997.  $\alpha$ -crystallin sequences support a galliform/anseriform clade. *Molecular Phylogenetics and Evolution* 7:185–188.
- CLEERE, N. 2002. Notes on the generic citation of the Oilbird *Steatornis caripensis* (Steatornithidae). *Bulletin of the British Ornithologists' Club* 122: 71–73.
- CORBIN, K. W. 1968. Taxonomic relationships of some *Columba* species. *Condor* 70:1–13.
- CRACRAFT, J. 1998. The major clades of birds. Pages 339–361 in *The Phylogeny and Classification of the Tetrapods. Volume 1, Amphibians, Reptiles, Birds* (M. J. Benton, Ed.). Clarendon Press, Oxford.
- CRACRAFT, J., AND J. CLARK. 2001. The basal clades of modern birds. Pages 143–156 in *New Perspectives on the Origin and Early Evolution of Birds: Proceedings of the International Symposium in honor of John H. Ostrom* (J. Gauthier and L. F. Gall, Eds.). Peabody Museum of Natural History, Yale University, New Haven, Connecticut.
- CUMLEY, R. W., AND M. R. IRWIN. 1944. The correlation between antigenic composition and geographic range in the Old or the New World of some species of *Columba*. *American Naturalist* 78:238–256.
- DAVID, N., AND M. GOSSELIN. 2002. Gender agreement of avian species names. *Bulletin of the British Ornithologists' Club* 122:14–49.
- DZERHINSKY, R. Y. 1995. Evidence for common ancestry of Galliformes and Anseriformes. *Courier Forschungsinstitut Senckenberg* 181: 325–336.
- FORD, N. L. 1967. A systematic study of the owls based on comparative osteology. Ph.D. Dissertation, University of Michigan, Ann Arbor.
- GARCÍA-MORENO, J., J. OHLSON, AND J. FJELDSÅ. 2001. MtDNA sequences support monophyly of *Hemispingus* tanagers. *Molecular Phylogenetics and Evolution* 21:424–435.
- GROTH, J. G., AND G. F. BARROWCLOUGH. 1999. Basal divergences in birds and the phylogenetic utility of the nuclear RAG-1 gene. *Molecular Phylogenetics and Evolution* 12:115–123.
- HELLMAYR, C. E., AND B. CONOVER. 1942. *Catalogue of Birds of the Americas*. Field Museum of Natural History Publications, Zoological Series, vol. 13, part 1, no. 1.
- HO, C. Y.-K., E. M. PRAGER, A. C. WILSON, D. T. OSUGA, AND R. E. FEENEY. 1976. Penguin evolution: Protein comparisons demonstrate phylogenetic relationship to flying aquatic birds. *Journal of Molecular Evolution* 8:271–282.
- JOHNSON, K. P., AND D. H. CLAYTON. 2000. Nuclear and mitochondrial genes contain similar phylogenetic signal for pigeons and doves (Aves: Columbiformes). *Molecular Phylogenetics and Evolution* 14:141–151.
- JOHNSON, K. P., S. DE CORT, K. DINWOODEY, A. C. MATEMAN, C. TEN CATE, C. M. LESSELLS, AND D. H. CLAYTON. 2001. A molecular phylogeny of the dove genera *Streptopelia* and *Columba*. *Auk* 118:874–887.
- JOHNSTON, R. F. 1962. The taxonomy of pigeons. *Condor* 64:69–74.
- KLICKA, J., K. P. JOHNSON, AND S. M. LANYON. 2000. New World nine-primaried oscine relationships: Constructing a mitochondrial DNA framework. *Auk* 117:321–336.
- KÖNIG, C., F. WEICK, AND J.-H. BECKING. 1999. *Owls: A Guide to the Owls of the World*. Yale University Press, New Haven, Connecticut.
- LIVEZEY, B. C. 1997. A phylogenetic analysis of basal Anseriformes, the fossil *Presbyornis*, and the interordinal relationships of waterfowl. *Zoological Journal of the Linnean Society* 121:361–428.
- LOVETTE, I. J., AND E. BERMINGHAM. 2002. What is a wood-warbler? Molecular characterization of a monophyletic Parulidae. *Auk* 119:695–714.
- MAYR, E., AND G. W. COTTRELL. (Eds.) 1979. *Check-list of Birds of the World*, vol. 1, 2nd ed. Museum of Comparative Zoology, Cambridge, Massachusetts.
- MINDELL, D. P., M. D. SORENSON, C. J. HUDDLESTON, H. C. MIRANDA, JR., A. KNIGHT, S. J. SAWCHUK, AND T. YURI. 1997. Phylogenetic relationships among and within select avian orders based on mitochondrial DNA. Pages 213–247 in *Avian Molecular Evolution and Systematics* (D. P. Mindell, Ed.). Academic Press, San Diego.
- MURPHY, R. C. 1936. *Oceanic Birds of South America*. 2 vols. McMillan Co., New York.
- PETERS, J. L. 1948. *Check-list of Birds of the World*, vol. 6. Harvard University Press, Cambridge, Massachusetts.
- REICHENBACH, H. G. L. 1852 (1853). *Handbuch der Speziellen Ornithologie, Die Vögel*, part 3. Dresden.

- RIDGWAY, R. 1914. The Birds of North and Middle America. Bulletin of the U.S. National Museum, no. 50, part 6.
- RIDGWAY, R. 1916. The Birds of North and Middle America. Bulletin of the U.S. National Museum, no. 50, part 7.
- SCLATER, P. L. 1886. Catalogue of the Birds in the British Museum. Vol. XI. British Museum, London.
- SHORT, L. L. 1982. Woodpeckers of the World. Monograph Series no. 4, Delaware Museum of Natural History. Greenville, Delaware.
- SIBLEY, C. G., AND J. E. AHLQUIST. 1990. Phylogeny and Classification of Birds. Yale University Press, New Haven, Connecticut.
- SIBLEY, C. G., AND B. L. MONROE, JR. 1990. Distribution and Taxonomy of Birds of the World. Yale University Press, New Haven, Connecticut.
- SMITH, P. W. 1997. The history and taxonomic status of the Hispaniolan Crossbill *Loxia megaloplaga*. Bulletin of the British Ornithologists' Club 117:264–271.
- VAN TUINEN, M., C. G. SIBLEY, AND S. B. HEDGES. 2000. The early history of modern birds inferred from DNA sequences of nuclear and mitochondrial ribosomal genes. Molecular Biology and Evolution 17:451–457.
- WETMORE, A. 1968. The Birds of the Republic of Panamá, part 2. Smithsonian Miscellaneous Collections, vol. 150.
- WINK, M., AND P. HEIDRICH. 1999. Molecular evolution and systematics of the owls (Strigiformes). Pages 39–57 in *Owls: A Guide to Owls of the World*. Yale University Press, New Haven, Connecticut.
- WINKLER, H., AND L. L. SHORT. 1978. A comparative analysis of acoustical signals in pied woodpeckers (Aves, *Picoides*). Bulletin of the American Museum of Natural History 160: 1–110.
- YURI, T., AND D. P. MINDELL. 2002. Molecular phylogenetic analysis of Fringillidae, “New World nine-primaried oscines” (Aves: Passeriformes). Molecular Phylogenetics and Evolution 23: 229–243.
- ZINK, R. M., S. ROHWER, A. V. ANDREEV, AND D. L. DITTMANN. 1995. Trans-Beringia comparisons of mitochondrial DNA differentiation in birds. Condor 97:639–649.
- ZINK, R. M., S. ROHWER, S. DROVETSKI, R. C. BLACKWELL-RAGO, AND S. L. FARRELL. 2002. Holarctic phylogeography and species limits of three-toed woodpeckers. Condor 104: 167–170.

This Supplement is a publication of the American Ornithologists' Union. Copies are available for \$3.00 from Buteo Books, 3130 Laurel Road, Shipman, VA 22971, USA. Buteo Books is the official sales outlet for publications of the AOU.

DATE OF ISSUE (Vol. 120, No. 3): 7 August 2003