

## **Exotic C4 Grasses Have Increased Tolerance to Glyphosate under Elevated Carbon Dioxide**

Authors: Manea, A, Leishman, M. R, and Downey, P. O

Source: Weed Science, 59(1) : 28-36

Published By: Weed Science Society of America

URL: <https://doi.org/10.1614/WS-D-10-00080.1>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## Exotic C<sub>4</sub> Grasses Have Increased Tolerance to Glyphosate under Elevated Carbon Dioxide

A. Manea, M. R. Leishman, and P. O. Downey\*

The increase in atmospheric CO<sub>2</sub> levels can influence the growth of many invasive exotic plant species. However, it is not well-documented, especially for C<sub>4</sub> plants, how these growth responses will alter the effectiveness of the world's most widely used herbicide for weed control, glyphosate. We aimed to address this question by carrying out a series of glasshouse experiments to determine if tolerance to glyphosate is increased in four C<sub>4</sub> invasive exotic grasses grown under elevated CO<sub>2</sub> in nonlimiting water conditions. In addition, traits including specific leaf area, leaf weight ratio, leaf area ratio, root : shoot ratio, total leaf area, and total biomass were measured in order to assess their contribution to glyphosate response under ambient and elevated CO<sub>2</sub> levels. Three of the four mature grass species that were treated with the recommended concentration of glyphosate displayed increased tolerance to glyphosate under elevated CO<sub>2</sub>. This was due to increased biomass production resulting in a dilution effect on the glyphosate within the plant. From this study, we can conclude that as atmospheric CO<sub>2</sub> levels increase, application rates of glyphosate might need to be increased to counteract the growth stimulation of invasive exotic plants.

**Nomenclature:** Glyphosate.

**Key words:** Alien plants, biomass, climate change, herbicide, photosynthetic pathway.

The nonselective herbicide glyphosate has been utilized extensively on a global basis since 1974 (Perez and Kogan 2003). Its ability to control a broad range of weeds has made it the world's most important herbicide (Baylis 2000; Powles 2008; Woodburn 2000). Glyphosate works by inhibiting 5-enolpyruvylshikimate-3-phosphate synthase, an enzyme in the shikimic acid pathway. The inhibition of this enzyme prevents the biosynthesis of aromatic amino acid, which causes a halt in the formation of proteins and secondary compounds (Bradshaw et al. 1997).

With the emergence of climate change as a global issue, studying the response of plants to different climate change components and the resulting effects on the efficacy of glyphosate should be a priority. One critical component of climate change is the rising concentration of atmospheric CO<sub>2</sub>. Over the last two decades, the amount of CO<sub>2</sub> available to plants has increased significantly (IPCC 2007). This is due mainly to the burning of fossil fuels and changes in land use that have caused an increase in the amount of atmospheric CO<sub>2</sub> from a preindustrial era concentration of 280 parts per million (ppm) to 379 ppm in 2005 (IPCC 2007). This trend is set to continue with atmospheric CO<sub>2</sub> predicted to reach 700 ppm by 2100 (IPCC 2007). An increase in atmospheric CO<sub>2</sub> could result in anatomical, morphological, and physiological changes in many plants that could influence uptake rates, transport, and the overall effectiveness of glyphosate (Bradshaw et al. 1997; Ziska and Teasdale 2000).

Plants that utilize the C<sub>3</sub> photosynthetic pathway are predicted to be favored under increased atmospheric CO<sub>2</sub> because the current CO<sub>2</sub> concentration is suboptimal for C<sub>3</sub> photosynthesis (Leegood 2002). Additional CO<sub>2</sub> causes a "fertilization effect," resulting in higher growth rates (Belote et al. 2003; Dukes 2002; Erickson et al. 2007; Poorter and Navas 2003; Sasek and Strain 1988, 1991; Smith et al. 2000;

Wray and Strain 1987; Ziska 2002; Ziska et al. 2004, 2005, 2007) and greater total leaf area (Sasek and Strain 1991; Wray and Strain 1987; Ziska et al. 2007). These changes might have a dilution effect on the glyphosate rendering it less effective than under current atmospheric CO<sub>2</sub> concentrations. In addition, C<sub>3</sub> plants display decreased stomatal conductance and stomatal number and increased leaf thickness that may further compromise glyphosate's effectiveness by restricting its foliar uptake (Ainsworth and Long 2005; Nowak et al. 2004; Ziska and Teasdale 2000).

Similar to C<sub>3</sub> plants, stomatal conductance in C<sub>4</sub> plants consistently is reduced under elevated CO<sub>2</sub> (Ainsworth and Long 2005; Wand et al. 1999). However, for C<sub>4</sub> plants, CO<sub>2</sub> fixation is saturated at 360 ppm, so these plants are less likely to show a positive response to additional CO<sub>2</sub> availability (Leegood 2002). This makes predicting the response of C<sub>4</sub> plants to increased atmospheric CO<sub>2</sub> more difficult (Dukes 2000). This unpredictability is demonstrated by studies that show the growth of C<sub>4</sub> plants can be stimulated (Bazzaz et al. 1989; Owensby et al. 1993, 1999), not affected (Erickson et al. 2007; Wray and Strain 1987), or inhibited (Belote et al. 2003) under elevated CO<sub>2</sub>. These variable responses could result from variation in environmental conditions such as water availability (Owensby et al. 1993, 1999), making accurate predictions about the effectiveness of glyphosate on C<sub>4</sub> plants under elevated CO<sub>2</sub> more difficult in comparison to C<sub>3</sub> plants. For this reason we chose to focus on the efficacy of glyphosate under elevated CO<sub>2</sub> in C<sub>4</sub> plants.

We are aware of only three studies that have looked at changes in tolerance of plant species to glyphosate under elevated CO<sub>2</sub>. Ziska et al. (1999) studied two weedy species, *Amaranthus retroflexus* L. and *Chenopodium album* L., which are C<sub>4</sub> and C<sub>3</sub>, respectively. Irrespective of CO<sub>2</sub> level, *A. retroflexus* displayed a reduction in growth and subsequent elimination as a result of glyphosate application. In contrast, the growth of *C. album* was not affected by glyphosate application under elevated CO<sub>2</sub>. From this, the authors concluded that C<sub>3</sub> plants have an increased tolerance to glyphosate under elevated CO<sub>2</sub>. These results were reinforced by Ziska and Teasdale (2000) and Ziska et al. (2004) who studied the C<sub>3</sub> perennial weeds *Elytrigia repens* (L.) Desv. ex

DOI: 10.1614/WS-D-10-00080.1

\*First and second authors: Research Assistant and Associate Professor, Department of Biological Sciences, Macquarie University, North Ryde, New South Wales 2109, Australia; third author: Weed Ecologist, Pest Management Unit, Parks and Wildlife, Department of Environment, Climate Change and Water, P.O. Box 1967, Hurstville, New South Wales 2220, Australia. Corresponding author's E-mail: michelle.leishman@mq.edu.au

B.D. Jackson and *Cirsium arvense* (L.) Scop., respectively. The reason for this increased tolerance to glyphosate under elevated CO<sub>2</sub> is unclear, but it has been suggested that an increase in the root : shoot ratio might play a role (Ziska et al. 2004). Thus, it is apparent that more research is required on two fronts: (1) the tolerance of plants, especially C<sub>4</sub> plants, to glyphosate under elevated CO<sub>2</sub>; and (2) the underlying mechanism responsible for changes in tolerance.

In the current study, we examined four C<sub>4</sub> grass species to determine: (1) the efficacy of glyphosate on different age cohorts (seedling, juvenile, mature) of the plants under two CO<sub>2</sub> concentrations (ambient and elevated), (2) regrowth responses after glyphosate application under ambient and elevated CO<sub>2</sub>, and (3) growth and allocation responses that might be responsible for altering the effectiveness of glyphosate under elevated CO<sub>2</sub>. The grasses selected for this study all have been introduced to Australia and are invasive along the east coast from southeastern Queensland to northeastern Victoria (Harden 1993). The experiment was carried out in nonlimiting water conditions. As stomatal conductance is decreased in C<sub>4</sub> plants under elevated CO<sub>2</sub> (Ainsworth and Long 2005; Wand et al. 1999), nonlimiting water conditions should not affect the relative success of the grasses under elevated CO<sub>2</sub>.

## Materials and Methods

**Species Selection.** The four grass species selected were *Chloris gayana* Kunth, *Eragrostis curvula* (Schrad.) Nees, *Paspalum dilatatum* Poir., and *Sporobolus indicus* (L.) R. Br. Species selection was based on two criteria: (1) they had to be invasive exotics within Australia, and (2) they are chemically controlled with glyphosate for land management. All species are C<sub>4</sub> grasses and have a fairly rapid development so that mature developmental stages could be incorporated in the study.

**Experimental Design.** Each species was grown in a series of glasshouse experiments under ambient (380 to 420 ppm) and elevated (675 to 715 ppm) CO<sub>2</sub> conditions. These CO<sub>2</sub> concentration ([CO<sub>2</sub>]) ranges were maintained by a CO<sub>2</sub> dosing and monitoring system.<sup>1</sup> The lower concentrations of these ranges tended to occur at nighttime, and the higher concentrations tended to occur during the daytime. The ambient CO<sub>2</sub> treatment represents the atmospheric CO<sub>2</sub> concentration during the turn of the 21st century (IPCC 2007). The elevated CO<sub>2</sub> treatment represents the potential atmospheric CO<sub>2</sub> concentration by 2100 (IPCC 2007). For each CO<sub>2</sub> treatment, individual plants were planted at three intervals (1, 2, and 3 mo) prior to glyphosate application. These planting intervals represent the postemergent developmental stages (seedling, juvenile, mature) of the grasses. Two glyphosate application rates or concentrations ([glyphosate]) were applied to each [CO<sub>2</sub>] by cohort treatment: recommended (10 ml L<sup>-1</sup>) and double recommended (20 ml L<sup>-1</sup>). There were eight replicates for each of the 12 treatments (two [CO<sub>2</sub>] by three cohorts by two [glyphosate] = 12 treatments). In addition, eight extra control plants per grass species were grown for each [CO<sub>2</sub>] by cohort treatment to be harvested prior to glyphosate application. This design resulted in 576 plants (four grass species by two [CO<sub>2</sub>] by three cohorts by two [glyphosate] by eight replicates + four grass species by two [CO<sub>2</sub>] by three cohorts by eight replicates). In total, four

Table 1. Results from the log rank tests of overall survival rates for each cohort and glyphosate treatment or concentration ([glyphosate]) combination. Values shown are P values and significant differences are shown in bold. Letter in last column indicates under which CO<sub>2</sub> level (A = Ambient, E = Elevated) the grass species had a significantly (P = 0.05) higher survival rate.

| [Glyphosate] | Cohort   | df | χ <sup>2</sup> | P value          |
|--------------|----------|----|----------------|------------------|
| Recommended  | Young    | 1  | 0.509          | 0.476            |
| Double       | Young    | 1  | 0.509          | 0.476            |
| Recommended  | Juvenile | 1  | 0.220          | 0.639            |
| Double       | Juvenile | 1  | 0.220          | 0.639            |
| Recommended  | Mature   | 1  | 39.290         | < <b>0.001 E</b> |
| Double       | Mature   | 1  | 2.392          | 0.122            |

glasshouses were used in this study, two for each CO<sub>2</sub> level. The pots within a CO<sub>2</sub> level were evenly split between the two glasshouses so each glasshouse contained the same number of replicates for each cohort by [glyphosate] treatment. The reason for doing this was twofold: (1) to act as insurance in case one glasshouse malfunctioned during the experiment, and (2) to average out the potentially differing conditions between the glasshouses. The temperature of the glasshouses was set for a maximum of 25 C and a minimum of 19 C. This temperature range was maintained with a reverse cycle inverter-ducted air conditioning unit. To avoid variation in growing conditions within each glasshouse, pots within each glasshouse were randomly moved to new positions on a fortnightly basis.

**Seed Collection and Germination.** Seeds for each of the four grass species were collected from a range of individual plants from sites in the Hawkesbury region of western Sydney. Once collected, the seeds for each of the grass species were germinated on moist filter paper within petri dishes.

**Planting and Growth Conditions.** Planting occurred 3, 2, and 1 mo prior to glyphosate application in order to produce three age cohorts of individuals: emergent seedlings, juveniles, and mature plants. For each cohort, seedlings were transplanted into pots in the glasshouses at the stage of cotyledon emergence. Multiple seedlings were transplanted into each pot to act as insurance against seedling mortality. After 3 d, any excess seedlings were removed from the pot. The seedlings were transplanted into pots 175 mm diam and 195 mm deep. The soil mixture used was obtained from a commercial supplier<sup>2</sup> and consisted of soil, double washed sand, composted sawdust, and graded ash in a ratio of 5 : 2 : 2 : 1. Each pot contained 2.4 L of this mixture. The pots were lined with newspaper to prevent soil loss through the drainage holes in the pots. The grasses were mist watered for 2 min three times daily. To counteract the nutrient loss resulting from this daily watering, 6.5 ± 0.2 g of slow release native plant fertilizer<sup>3</sup> (23N : 2P : 17K) was added to each pot.

**Glyphosate Application and Harvesting.** One d prior to glyphosate<sup>4</sup> application, the additional control plants from each cohort by [CO<sub>2</sub>] treatment were harvested. These plants were separated into the following components: three fully expanded leaves, remaining leaf biomass, remaining above-ground biomass, and belowground biomass. The plant components were then washed free of soil before being oven-dried at 80 C for 48 h and weighed using a Mettler Toledo B-S electronic balance.




Figure 1. Survival rates of all the age cohort by glyphosate combinations grown under ambient and elevated CO<sub>2</sub> glasshouse conditions for 6 wk. The lines indicate the mean of eight plants grown under each CO<sub>2</sub> treatment (solid = ambient, dashed = elevated) for the four grass species combined.

On the day following the initial harvest, the glyphosate treatment (10 ml per plant) was applied to all remaining plants. After 42 d, the surviving plants were harvested and oven-dried as previously described but were not split into their components.

During the 42-d period after glyphosate application, the mortality of the grasses was recorded on a weekly basis. This involved classifying each individual grass as alive or dead. A grass was classified as dead if all its leaves became discolored and died. A plant was classified as alive if it showed signs of regeneration by the end of the 42 d. If a grass was classified as dead it remained in the experiment for the entire 42-d period and continued to be watered. Grasses that were classified as dead still were harvested after the 42-d period to ensure there was no living root biomass. There were no instances where

living root biomass was found in the pots of the grasses classified as dead, so we can assume this classification system was reliable.

**Measuring Plant and Allocation Traits.** In order to determine if there was any relationship between plant traits and tolerance to glyphosate, the following range of traits were measured or calculated for each grass species: Specific leaf area (SLA) was measured as the leaf area per unit leaf mass for three randomly selected outer canopy leaves. Leaf area was measured using a LI-3100C Area Meter<sup>5</sup> prior to oven-drying the leaves for weighing. Leaf weight ratio (LWR) was calculated as total plant leaf mass divided by total plant mass.




Figure 2. Survival rates of all the age cohort by glyphosate combinations grown under ambient and elevated CO<sub>2</sub> glasshouse conditions for 6 wk. The lines indicate the mean of eight plants grown under each CO<sub>2</sub> treatment (solid = ambient, dashed = elevated) for *C. gayana*.

Leaf area ratio (LAR) was calculated as SLA times LWR. Root-to-shoot ratio (R : S) was calculated as total root biomass divided by total shoot biomass. Total leaf area ( $L_T$ ) was measured as the sum of the leaf area of all leaves on a plant. Total biomass ( $B_T$ ) was calculated as the sum of the dry weight of all of the harvested components.

**Across-Species Data Analysis.** To determine the survival function of all the grass species in response to CO<sub>2</sub> level, Kaplan-Meier survival curves were generated. The survival distributions for each of the grass species under ambient and elevated CO<sub>2</sub> were compared using a log rank test, with significance determined at  $P < 0.05$ . These analyses were carried out for each cohort by glyphosate treatment across all grass species.

**Species-Level Data Analysis.** To determine the survival function of each grass species in response to CO<sub>2</sub> level, Kaplan-Meier survival curves were generated. The survival distributions for each of the grass species under ambient and elevated CO<sub>2</sub> were compared using a log rank test, with significance determined at  $P < 0.05$ . These analyses were carried out for each cohort by glyphosate treatment within each grass species.

Species that survived glyphosate treatment had total leaf senescence followed by resprouting. To assess this regrowth response for each grass species under ambient and elevated CO<sub>2</sub>, the regrowth biomass data was analysed using two sample t-tests. This analysis was carried out for both the recommended and double glyphosate treatments in each cohort for each of the four grass species.




Figure 3. Survival rates of all the age cohort by glyphosate combinations grown under ambient and elevated CO<sub>2</sub> glasshouse conditions for 6 wk. The lines indicate the mean of eight plants grown under each CO<sub>2</sub> treatment (solid = ambient, dashed = elevated) for *E. curvula*.

To determine if the plant growth and allocation traits were influenced by CO<sub>2</sub> level, the trait data was analysed using two sample *t*-tests. This analysis was carried out for all three cohorts for each of the four grass species.

For both the overall and species-level data analyses, the significance level was set at 0.05. The regrowth biomass and trait data analyses were carried out using Minitab 15 Statistical Software (Minitab, Inc. 2007). The survival analysis was carried out using SAS 9.2 (SAS Institute, Inc. 2008).

## Results and Discussion

In this study we tested whether the efficacy of glyphosate on invasive exotic grass species is affected under elevated CO<sub>2</sub> conditions. Our results showed that the mature invasive exotic grasses that were sprayed with the recommended concentra-

tion of glyphosate had a significantly higher survival rate under elevated CO<sub>2</sub> compared with ambient CO<sub>2</sub> ( $\chi^2 = 39.290$ ,  $P < 0.001$ ; Table 1; Figure 1). To determine if particular species were driving this result, the data were analyzed at the species level. This analysis showed that *C. gayana* ( $\chi^2 = 8.641$ ,  $P = 0.003$ ; Figure 2), *E. curvula* ( $\chi^2 = 7.752$ ,  $P = 0.005$ ; Figure 3), and *P. dilatatum* ( $\chi^2 = 31.200$ ,  $P < 0.001$ ; Figure 4) had significantly higher survival rates under elevated CO<sub>2</sub> conditions, whereas *S. indicus* ( $\chi^2 = 0.830$ ,  $P = 0.362$ ; Figure 5) showed no difference in response between CO<sub>2</sub> treatments. Sufficient individuals survived of only two species to enable analysis of this regrowth response (the mature cohort of *C. gayana* and *S. indicus* at both glyphosate treatments). The regrowth responses of *C. gayana* ( $t_6 = -0.240$ ,  $P_{\text{Recommended}} = 0.818$ ;  $t_5 = 0.752$ ,  $P_{\text{Double}} = 0.486$ ) and *S. indicus* ( $t_7 = 0.096$ ,  $P_{\text{Recommended}} = 0.926$ ;




Figure 4. Survival rates of all the age cohort by glyphosate combinations grown under ambient and elevated CO<sub>2</sub> glasshouse conditions for 6 wk. The lines indicate the mean of eight plants grown under each CO<sub>2</sub> treatment (solid = ambient, dashed = elevated) for *P. dilatum*.

$t_2 = 1.709$ ,  $P_{\text{Double}} = 0.229$ ) did not differ significantly between ambient and elevated CO<sub>2</sub> for either of the glyphosate treatments.

We examined growth and allocation traits of the four species before glyphosate application in order to aid interpretation of the survival data (Table 2). The three grass species (*C. gayana*, *E. curvula*, and *P. dilatum*) that had a significantly higher survival rate under elevated CO<sub>2</sub> also produced more biomass (39, 83, and 59% increase, respectively) and leaf area (40, 67, and 24% increase, respectively) under these conditions. These trait differences between ambient and elevated CO<sub>2</sub> were significant for the biomass production of *E. curvula* ( $t_{14} = 3.475$ ,  $P = 0.005$ ) and *P. dilatum* ( $t_{14} = 4.134$ ,  $P = 0.001$ ), and total leaf area of *E. curvula* ( $t_{14} = 5.309$ ,  $P < 0.001$ ). In contrast, the biomass (5% decrease) and total leaf area (34% decrease)

produced by *S. indicus*, the only species not to have a significantly higher survival rate under elevated CO<sub>2</sub>, decreased between the two CO<sub>2</sub> levels. This suggests that the effectiveness of glyphosate might be proportional to the amount of plant tissue upon which it has to act. That is, a larger amount of biomass might dilute the glyphosate within the plant, rendering it less effective. In the future, if invasive exotic plant growth is stimulated under elevated CO<sub>2</sub> conditions, it might be necessary to increase the concentration at which glyphosate is applied to these plants to counteract the increase in biomass production.

The results of this study are consistent with the few previous studies that have found an increase in tolerance of invasive exotic C<sub>3</sub> plant species to glyphosate under elevated CO<sub>2</sub> levels compared to ambient CO<sub>2</sub> levels (Ziska et al. 1999, 2004; Ziska and Teasdale 2000). Our study has




Figure 5. Survival rates of all the age cohort by glyphosate combinations grown under ambient and elevated CO<sub>2</sub> glasshouse conditions for 6 wk. The lines indicate the mean of eight plants grown under each CO<sub>2</sub> treatment (solid = ambient, dashed = elevated) for *S. indicus*.

expanded this knowledge base to include a larger number of C<sub>4</sub> plant species. Ziska et al. (1999) is the only study of which we are aware that has tested the efficacy of glyphosate on a C<sub>4</sub> plant under ambient and elevated CO<sub>2</sub> levels. They found that glyphosate tolerance of the C<sub>4</sub> weedy species *A. retroflexus* was not affected by CO<sub>2</sub> concentration. Thus of the five C<sub>4</sub> species studied so far, three have shown increased tolerance of glyphosate under elevated CO<sub>2</sub> and two have shown no difference in tolerance.

This study suggests that the differing response in survival among species to glyphosate application between ambient and elevated CO<sub>2</sub> is related to their biomass production. The high mortality of seedlings and juvenile plants of all species under both glyphosate and CO<sub>2</sub> treatments is consistent with this.

However it should be stressed that there are other CO<sub>2</sub>-induced effects that were not measured by this study which might have influenced our results. Stomatal number and cuticle thickness are consistently reduced under elevated CO<sub>2</sub>, which might have limited the uptake of glyphosate into the plant tissue (Ainsworth and Long 2005; Wand et al. 1999). Protein content per gram of tissue also can be reduced under elevated CO<sub>2</sub>, which might reduce the need for aromatic amino acids (Bowes 1996). Glyphosate inhibits the production of these aromatic amino acids, but if there is not a large demand for them, glyphosate will be less effective (Ziska et al. 1999).

This study focused on the response of invasive exotic C<sub>4</sub> grasses to glyphosate application under ambient and elevated CO<sub>2</sub> for plants grown as individual plants under nonresource-



Table 2. Species-level trait means and standard errors for the four species grown under ambient and elevated CO<sub>2</sub> and harvested at 3 mo after planting. Significant P values from *t*-tests are shown in bold.

| Species                   | Trait              | Ambient CO <sub>2</sub> |         | Elevated CO <sub>2</sub> |         | df | <i>t</i> | P                 |
|---------------------------|--------------------|-------------------------|---------|--------------------------|---------|----|----------|-------------------|
|                           |                    | Mean                    | SE      | Mean                     | SE      |    |          |                   |
| <i>Chloris gayana</i>     | Total biomass      | 17.240                  | 2.790   | 23.922                   | 5.448   | 14 | 1.798    | 0.094             |
|                           | Specific leaf area | 16.373                  | 1.786   | 20.662                   | 0.502   | 14 | 5.253    | <b>&lt; 0.001</b> |
|                           | Leaf weight ratio  | 0.306                   | 0.016   | 0.278                    | 0.020   | 14 | -1.232   | 0.246             |
|                           | Leaf area ratio    | 5.054                   | 0.852   | 5.732                    | 0.627   | 14 | 1.367    | 0.199             |
|                           | Root : shoot ratio | 0.390                   | 0.027   | 0.417                    | 0.010   | 14 | 0.403    | 0.693             |
| <i>Eragrostis curvula</i> | Total leaf area    | 872.411                 | 175.187 | 1,222.533                | 138.408 | 14 | 2.155    | 0.052             |
|                           | Total biomass      | 5.303                   | 0.465   | 9.697                    | 0.708   | 14 | 3.475    | <b>0.005</b>      |
|                           | Specific leaf area | 16.838                  | 1.969   | 18.224                   | 2.324   | 14 | 1.109    | 0.289             |
|                           | Leaf weight ratio  | 0.527                   | 0.025   | 0.658                    | 0.017   | 14 | 0.299    | 0.317             |
|                           | Leaf area ratio    | 5.364                   | 1.177   | 5.340                    | 0.961   | 14 | -0.048   | 0.962             |
| <i>Paspalum dilatatum</i> | Root : shoot ratio | 0.193                   | 0.018   | 0.240                    | 0.032   | 14 | 1.056    | 0.310             |
|                           | Total leaf area    | 287.283                 | 19.068  | 479.019                  | 42.303  | 14 | 5.309    | <b>&lt; 0.001</b> |
|                           | Total biomass      | 30.335                  | 1.379   | 48.128                   | 0.900   | 14 | 4.134    | <b>0.001</b>      |
|                           | Specific leaf area | 27.693                  | 3.061   | 28.789                   | 1.627   | 14 | 0.574    | 0.575             |
|                           | Leaf weight ratio  | 0.283                   | 0.017   | 0.202                    | 0.019   | 14 | -6.051   | <b>&lt; 0.001</b> |
| <i>Sporobolus indicus</i> | Leaf area ratio    | 7.899                   | 1.414   | 5.825                    | 0.240   | 14 | -2.920   | <b>0.014</b>      |
|                           | Root : shoot ratio | 0.308                   | 0.024   | 0.375                    | 0.046   | 14 | 1.870    | 0.098             |
|                           | Total leaf area    | 1,990.130               | 97.082  | 2,463.355                | 90.617  | 14 | 1.562    | 0.142             |
|                           | Total biomass      | 2.629                   | 0.620   | 2.496                    | 0.583   | 14 | -0.156   | 0.878             |
|                           | Specific leaf area | 19.576                  | 0.912   | 17.045                   | 0.717   | 14 | -2.182   | <b>0.048</b>      |
| <i>Sporobolus indicus</i> | Leaf weight ratio  | 0.378                   | 0.018   | 0.465                    | 0.053   | 14 | 1.570    | 0.151             |
|                           | Leaf area ratio    | 7.409                   | 0.511   | 7.930                    | 0.934   | 14 | 0.489    | 0.635             |
|                           | Root : shoot ratio | 0.327                   | 0.034   | 0.176                    | 0.045   | 14 | -2.673   | <b>0.019</b>      |
|                           | Total leaf area    | 196.101                 | 44.775  | 129.657                  | 41.737  | 14 | -1.086   | 0.296             |

limited glasshouse conditions. These conditions might not accurately represent those experienced in the grasses' natural environment where there is competition for resources such as soil moisture, nutrients, and light. Previous work has shown that plant response to elevated CO<sub>2</sub> is constrained by resource availability (Ören et al. 2001; Poorter et al. 1996; Reich et al. 2006). Nevertheless, our study has shown clearly that when plants are able to respond to elevated CO<sub>2</sub> with greater growth and biomass production, particularly leafy biomass, this will increase their tolerance of glyphosate application. This suggests that increasing atmospheric concentrations of CO<sub>2</sub> might require an increase in application rates of glyphosate, which could have significant economic and environmental consequences.

### Sources of Materials

<sup>1</sup> CO<sub>2</sub> dosing and monitoring system, Canary Company Pty. Ltd., 1/163 Burns Bay Rd., Lane Cove, New South Wales 2066, Australia.

<sup>2</sup> Coarse river sand and organic garden mix, Australian Native Landscapes, Terrey Hills, New South Wales 2084, Australia.

<sup>3</sup> Slow release low phosphorus fertilizer, J. R. Somplot Co., P.O. Box 27, Boise, ID 83707.

<sup>4</sup> Glyphosate as isopropylamine salt, Monsanto Company, P.O. Box 1750, Columbus, OH 43216-1750.

<sup>5</sup> LI-3100C Area Meter, Li-Cor Corporation, 4808 Progressive Avenue, Lincoln, NE 68504.

### Acknowledgments

We thank the Plant Invasion and Restoration Ecology Laboratory (PIREL) of Macquarie University for their input throughout the experiment and Muhammad Masood for assistance with the glasshouses. This research was funded by an Australian Research Council Linkage grant (LP0776758).

### Literature Cited

- Ainsworth, E. A. and S. P. Long. 2005. What have we learned from 15 years of free-air CO<sub>2</sub> enrichment (FACE)? A meta-analytic review of the responses of photosynthesis, canopy properties and plant production to rising CO<sub>2</sub>. *New Phytol.* 165:351-372.
- Baylis, A. D. 2000. Why glyphosate is a global herbicide: strengths, weaknesses and prospects. *Pest Manag. Sci.* 56:299-308.
- Bazzaz, F. A., K. Garbutt, E. G. Reekie, and W. E. Williams. 1989. Using growth analysis to interpret competition between a C<sub>3</sub> and a C<sub>4</sub> annual under ambient and elevated CO<sub>2</sub>. *Oecologia* 79:223-235.
- Belote, R. T., J. F. Weltzin, and R. J. Norby. 2003. Response of an understory plant community to elevated [CO<sub>2</sub>] depends on differential responses of dominant invasive species and is mediated by soil water availability. *New Phytol.* 161:827-835.
- Bowes, G. 1996. Photosynthetic responses to changing atmospheric carbon dioxide concentration. Pages 397-407 in N. R. Baker, ed. *Photosynthesis and the Environment*. Dordrecht, Netherlands: Kluwer Academic Publishers.
- Bradshaw, L. D., S. R. Padgett, S. L. Kimball, and B. H. Wells. 1997. Perspectives on glyphosate resistance. *Weed Technol.* 11:189-198.
- Dukes, J. S. 2000. Will the increasing atmospheric CO<sub>2</sub> concentration affect the success of invasive species? Pages 95-113 in H. A. Mooney and R. J. Hobbs, ed. *Invasive Species in a Changing World*. Washington, D.C.: Island Press.
- Dukes, J. S. 2002. Comparison of the effect of elevated CO<sub>2</sub> on an invasive species (*Centaurea solstitialis*) in monoculture and community settings. *Plant Ecol.* 160:225-234.
- Erickson, J. E., J. P. Megeonigal, G. Peresta, and B. G. Drake. 2007. Salinity and sea level mediate elevated CO<sub>2</sub> effects on C<sub>3</sub>-C<sub>4</sub> plant interactions and tissue nitrogen in a Chesapeake Bay tidal wetland. *Glob. Change Biol.* 13:202-215.
- Harden, G. W. 1993. *Flora of New South Wales*. Chapter 213: *Poaceae*. Sydney, New South Wales, Australia: New South Wales University Press, pp. 410-656.
- IPCC. 2007. *Climate change 2007: a physical science basis*. In S. Solomon, M. Manning, Z. Chen, M. Marquis, K. B. Averyt, M. Tignor, and H. L. Miller, eds. *Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge, UK and New York: Cambridge University Press. 996 p.
- Leegood, R. C. 2002. C<sub>4</sub> photosynthesis: principles of CO<sub>2</sub> concentration and prospects for its introduction into C<sub>3</sub> plants. *J. Exp. Bot.* 53:581-590.
- Minitab, Inc. 2007. *Minitab 15: Statistical Software*. State College, PA: Publisher.
- Nowak, R. S., D. S. Ellsworth, and S. D. Smith. 2004. Functional responses of plants to elevated atmospheric CO<sub>2</sub>: do photosynthetic and productivity data from FACE experiments support early predictions? *New Phytol.* 162:253-280.
- Oren, R., D. S. Ellsworth, and K. H. Johnsen, et al. 2001. Soil fertility limits carbon sequestration by forest ecosystems in a CO<sub>2</sub>-enriched atmosphere. *Nature* 411:469-472.

- Owensby, C. E., P. I. Coyne, J. M. Ham, L. M. Auen, and A. K. Knapp. 1993. Biomass production in a tallgrass prairie ecosystem exposed to ambient and elevated CO<sub>2</sub>. *Ecol. Appl.* 3:644–653.
- Owensby, C. E., J. M. Ham, A. K. Knapp, and L. M. Auen. 1999. Biomass production and species composition change in a tallgrass prairie ecosystem after long-term exposure to elevated atmospheric CO<sub>2</sub>. *Glob. Change Biol.* 5:497–506.
- Perez, A. and M. Kogan. 2003. Glyphosate-resistant *Lolium multiflorum* in Chilean orchards. *Weed Res.* 43:12–19.
- Poorter, H. and M. L. Navas. 2003. Plant growth and competition at elevated CO<sub>2</sub>: on winners, losers and functional groups. *New Phytol.* 157:175–198.
- Poorter, H., C. Roumet, and B. D. Campbell. 1996. Interspecific variation in the growth response of plants to elevated CO<sub>2</sub>: a search for functional types. Pages 375–412 in C. Korner and F. A. Bazzaz, eds. *Carbon Dioxide, Populations and Communities*. San Diego, CA: Academic.
- Powles, S. B. 2008. Evolved glyphosate-resistant weeds around the world: lessons to be learnt. *Pest Manag. Sci.* 64:360–365.
- Reich, P. B., S. E. Hobbie, T. Lee, D. S. Ellsworth, J. B. West, D. Tilman, J.M.H. Knops, S. Naeem, and J. Trost. 2006. Nitrogen limitation constrains sustainability of ecosystem response to CO<sub>2</sub>. *Nature* 440:922–925.
- SAS. 2008. SAS 9.2. Cary, NC: SAS Institute, Inc.
- Sasek, T. W. and B. R. Strain. 1988. Effects of carbon dioxide enrichment on the growth and morphology of kudzu (*Pueraria lobata*). *Weed Sci.* 36:28–36.
- Sasek, T. W. and B. R. Strain. 1991. Effects of CO<sub>2</sub> enrichment on the growth and morphology of a native and an introduced honeysuckle vine. *Am. J. Bot.* 78:69–75.
- Smith, S. D., T. E. Huxman, S. F. Zitzer, T. N. Charlet, D. C. Housman, J. S. Coleman, L. K. Fenstermaker, J. R. Seemann, and R. S. Nowak. 2000. Elevated CO<sub>2</sub> increases productivity and invasive species success in an arid ecosystem. *Nature* 408:79–82.
- Wand, S.J.E., G. F. Midgley, M. H. Jones, and P. S. Curtis. 1999. Responses of wild C<sub>4</sub> and C<sub>3</sub> grass (Poaceae) species to elevated atmospheric CO<sub>2</sub> concentration: a meta-analytic test of current theories and perceptions. *Glob. Change Biol.* 5:723.
- Woodburn, A. T. 2000. Glyphosate: production, pricing and use worldwide. *Pest Manag. Sci.* 56:309–312.
- Wray, S. M. and B. R. Strain. 1987. Interaction of age and competition under CO<sub>2</sub> enrichment. *Funct. Ecol.* 1:145–149.
- Ziska, L. H. 2002. Influence of rising atmospheric CO<sub>2</sub> since 1900 on early growth and photosynthetic response of a noxious invasive weed, Canada thistle (*Cirsium arvense*). *Funct. Plant Biol.* 29:1387–1392.
- Ziska, L. H., S. Faulkner, and J. Lydon. 2004. Changes in biomass and root : shoot ratio of field-grown Canada thistle (*Cirsium arvense*), a noxious, invasive weed, with elevated CO<sub>2</sub>: implications for control with glyphosate. *Weed Sci.* 52:584–588.
- Ziska, L. H., J. B. Reeves, and B. Blank. 2005. The impact of recent increases in atmospheric CO<sub>2</sub> on biomass production and vegetative retention of cheatgrass (*Bromus tectorum*): implications for fire disturbance. *Glob. Change Biol.* 11:1325–1332.
- Ziska, L. H., R. C. Sicher, K. George, and J. E. Mohan. 2007. Rising atmospheric carbon dioxide and potential impacts on the growth and toxicity of poison ivy (*Toxicodendron radicans*). *Weed Sci.* 55:288–292.
- Ziska, L. H. and J. R. Teasdale. 2000. Sustained growth and increased tolerance to glyphosate observed in a C<sub>3</sub> perennial weed, quackgrass (*Elytrigia repens*), grown at elevated carbon dioxide. *Funct. Plant Biol.* 27:159–166.
- Ziska, L. H., J. R. Teasdale, and J. A. Bunce. 1999. Future atmospheric carbon dioxide may increase tolerance to glyphosate. *Weed Sci.* 47:608–615.

*Received May 26, 2010, and approved July 28, 2010.*