

Scirpus hattorianus (Cyperaceae), newly reported for Europe, naturalized in France

Author: Verloove, Filip

Source: Willdenowia, 44(1) : 51-55

Published By: Botanic Garden and Botanical Museum Berlin (BGBM)

URL: <https://doi.org/10.3372/wi.44.44108>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

FILIP VERLOOVE¹

Scirpus hattorianus (Cyperaceae), newly reported for Europe, naturalized in France

Abstract

Verloove F.: *Scirpus hattorianus* (Cyperaceae), newly reported for Europe, naturalized in France. – Willdenowia 44: 51–55. 14 March 2014. – Version of record published online ahead of inclusion in April 2014 issue; ISSN 1868-6397; © 2014 BGBM Berlin-Dahlem.

DOI: <http://dx.doi.org/10.3372/wi.44.44108>

The identity of the North American xenophyte *Scirpus atrovirens* in France was critically reassessed. Nearly all known populations (notably those from the departments Aisne, Jura, Pas-de-Calais and Saône-et-Loire) turned out to be referable to a closely related species, *S. hattorianus*. Distinguishing features for three species from the *S. atrovirens* complex (i.e. *S. atrovirens* s.str., *S. georgianus* and *S. hattorianus*) are compared and discussed. An identification key and SEM photographs of their achenes are provided. The ecology and degree of invasiveness of *S. hattorianus* in France are also briefly discussed.

Additional key words: ecology, invasive, North America, *Scirpus atrovirens*, *Scirpus georgianus*, taxonomy, xenophyte

Introduction

As currently circumscribed, *Scirpus* L. is a genus of c. 35 subcosmopolitan species (Mabberley 2008). It apparently is most diverse in North America, where 18 species occur (Schuyler 1967; Whittemore & Schuyler 2002). Only two species are originally native in Europe (DeFilipps 1980, as *S. sect. Scirpus*): *S. radicans* Schkuhr and *S. sylvaticus* L. A few others, all originating in North America, are locally naturalized: *S. cyperinus* (L.) Kunth in Germany (Lanski 1985; Kiffe 1998) and *S. pendulus* Muhl. in Switzerland (Bornand & Clerc 2011) and the former republic of Czechoslovakia (DeFilipps l.c.). However, a third American species is by far the most widespread in Europe: *S. atrovirens* Willd. It has been known from parts of C Europe and France for quite some time (DeFilipps l.c.) and over the past several decades it has been discovered in other European countries as well, for instance Belgium (Giot 2004; Lannoy 2004) and Italy (Pignotti 2003).

Scirpus atrovirens belongs to a group of closely related species that includes *S. flaccidifolius* (Fernald) Schuyler, *S. georgianus* R. M. Harper, *S. hattorianus* Makino and *S. pallidus* (Britton) Fernald (Tucker 1987). These four species are largely allopatric, but hybridize in areas where they occur sympatrically (Schuyler 1961; Schuyler 1967; Whittemore & Schuyler 2002). The identity of some of the European populations of *S. atrovirens* has been questioned. Some German plants were re-identified as *S. georgianus* (Schnittler & Niedbala 1993, as *S. atrovirens* var. *georgianus* (R. M. Harper) Fernald) and Kiffe (1998) even suggested that genuine *S. atrovirens* might not have been documented in Germany. Recent reports from Slovenia (Zelnik 2004) and Belgium (Verloove 2006; Verloove & Lambinon 2011) were also ascribed to *S. georgianus*. Similarly, plants from E France have been assigned to *S. pallidus* (Litzler 2000; Prost 2000; Bizot & Parent 2005, as *S. atrovirens* var. *pallidus* Britton, apparently following Riomet & Bournérias 1952–1961).

¹ Botanic Garden Meise, Domein van Bouchout, B-1860 Meise, Belgium; e-mail: filip.verloove@br.fgov.be

The present study, mainly based on a critical reassessment of herbarium specimens of *Scirpus* from Europe (herbaria BR, GENT, LG and P; herbarium codes following Thiers 2014+), found that nearly all collections from France were identified incorrectly and in fact belong to another species, *S. hattorianus*. In the present paper its distinguishing features are discussed and compared with related species; also some additional information on its distribution, ecology and degree of naturalization in France is provided.

The *Scirpus atrovirens* complex in France

Scirpus atrovirens s.l. has been known in France for more than a century. It was probably first collected in Nanteuil-Notre-Dame in the department Aisne where, by 1903, it was already known to occur (Riomet & Bournérias 1952–1961) and lately confirmed by Bizot & Parent (2005). From the 1970s onwards it was reported from various locations in the department Jura (Prost 1983; Litzler 2000; Prost 2000, 2001) and since the 1980s also from the department Marne (Lambinon & Duvigneaud 1988; Worms 1989). Bizot & Parent (2005) confirmed its presence in the aforementioned regions and added a first record from the department Saône-et-Loire. It was recently detected for the first time in the department Pas-de-Calais (B. Toussaint pers. comm. Dec 2011) and Tela Botanica (2012+) added a record from the department Ain.

Although usually referred to as “*Scirpus atrovirens*” in French floras or catalogues (e.g. Kerguélen 1999), populations from the department Aisne were ascribed to *S. atrovirens* var. *pallidus* (\equiv *S. pallidus*) by Riomet & Bournérias (1952–1961). This point of view was followed subsequently by Litzler (2000) and Bizot & Parent (2005). These authors drew attention to the (initially) pale brownish, small spikelet scales (c. 2 mm or slightly longer), characteristic features of *S. pallidus*. However, all specimens seen from France (including those referred to by Litzler l.c. and Bizot & Parent l.c.) clearly differ from this species: the scales are much darker and ultimately turn blackish brown, are much smaller (always less than 1.5 mm long) and have a mucro of 0.1–0.2 mm long, whereas *S. pallidus* has scales 1.6–2.8 mm long with a pale midrib and a distinct mucro 0.4–0.6(–1.2) mm long (Whittemore & Schuyler 2002). Thus, *S. pallidus* obviously does not compare well. A single population from a military camp site in Belgium (Marche-en-Famenne) belongs to *S. georgianus* (\equiv *S. atrovirens* var. *georgianus* \equiv *S. atrovirens* subsp. *georgianus* (R. M. Harper) Verloove & Lambinon). This species is readily told apart by the perianth bristles, which are either absent or at most 3, always much shorter than the achene (often rudimentary) and nearly smooth throughout (or with a few teeth near the apex only). Most collections examined from France have the following characteristics: leaf sheaths not or

only inconspicuously nodulose-septate; spikelet scales blackish brown in distal half, less than 1.5 mm long, with a mucro less than 0.2 mm long; perianth bristles always present, usually 4–6, all straight (never contorted), distinctly shorter than achene; and achenes at most 1 mm long. These features are typical of *S. hattorianus* and clearly exclude *S. atrovirens* s.str. A few collections from the French departments Bas-Rhin and Marne correspond with *S. georgianus* (see below for an overview).

The two species of *Scirpus* hitherto reliably recorded in W Europe and *S. atrovirens* s.str. can be separated using the following key:

1. Perianth bristles (not to be confused with filaments, which often persist!) absent or rudimentary, at most 3 if present, up to $\frac{3}{4}$ of achene length, smooth or with a few teeth near apex *Scirpus georgianus*
- Perianth bristles always present, (4 or)5 or 6, shorter than to slightly longer than achene, with retrorse teeth in distal $\frac{1}{2}$ 2
2. Perianth bristles (4 or)5 or 6, all distinctly shorter than achene; spikelet scales blackish brown in distal $\frac{1}{2}$, less than 1.5 mm long and with mucro less than 0.2 mm long; achenes mostly less than 1 mm long; leaf sheaths smooth to somewhat nodulose-septate *Scirpus hattorianus*
- Perianth bristles nearly always 6 (rarely 5), longer ones frequently exceeding achene; spikelet scales brownish, to 2.1 mm long and with mucro often more than 0.2 mm long; achenes mostly 1–1.3 mm long; lowermost leaf blades and sheaths distinctly nodulose-septate *Scirpus atrovirens*

SEM photos of achenes of the three species are compared in Fig. 1. Other useful SEM photos of achenes are provided by Strong (1994). Line drawings are found in Whittemore & Schuyler (2002).

The identity of other European populations of *Scirpus atrovirens* should be critically assessed. Achenes illustrated by Schnittler & Niedbala (1993), with at least five perianth bristles, possibly also represent *S. hattorianus* and certainly not *S. georgianus* as claimed. *Scirpus hattorianus* appears to be by far the most widespread species, at least in France. Despite being a North American species, it was originally described from Japan (Makino 1933), which clearly demonstrates its potential as an environmental weed.

Specimens examined

Scirpus hattorianus

FRANCE: DEPARTMENT AISNE: Nanteuil-Notre-Dame, dans un fossé voisin de la voie ferrée, adventice, Jul 1919, *Riomet s.n.* (P); Nanteuil (ex cultu), trouvé à l'état adventif dans les fossés avoisinant la voie ferrée, 12 Aug 1923, *G. Desplantes s.n.* (P). — DEPARTMENT




Fig. 1. SEM photographs of achenes of *Scirpus* – A: *S. hattorianus* from France (F. Verloove 9564, see specimens examined); B: *S. georgianus* from Belgium; C: *S. atrovirens* from North America.

JURA: Dole, Forêt Domaniale de Chaux, laies forestières fraîches, mi-ombragées, 4 Aug 1998, P. Litzler 98/659 (P); *ibid.*, 18 Jun 1999, P. Litzler 99/425 (P); Rye, Forêt de Rye, sentier forestier sur sol argilo-siliceux détrempe en hiver, partant de la D 214, abondant, 1 Jul 2001, J.-M. Tison *s.n.* (BR, LG); Bois de Champrougier, en Forêt d'Amont-Aval, layon forestier entre les routes Les Deux Fays-Foulénay et Les Deux Fays-Champrougier, 15 Jul 2004, G. H. Parent 04/120 (BR); Bois de Rye (Forêt d'Amont-Aval), 16 Jul 2004, G. H. Parent 04/121G (BR); Bois de Rye, vers Le Ferment (Forêt d'Amont-

Aval), 16 Jul 2004, G. H. Parent 04/121 (BR); Sergenon, Forêt d'Amont-Aval, bois à l'W du village, 16 Jul 2004, G. H. Parent 04/121 (BR); Sergenon, Bois en Bief Vallon, 16 Jul 2004, G. H. Parent 04/121 (BR); Tassenières, Bois d'Amont-Aval, bord des chemins dans le bois au NW de la N475, parcelle 30, 16 Jul 2004, G. H. Parent 04/121 (BR); Le Villey (Forêt d'Amont), bordure d'un chemin forestier, 16 Jul 2004, G. H. Parent 04/121 (BR). — DEPARTMENT PAS-DE-CALAIS: Bournonville, Forêt Domaniale de Desvres, close to RD 253, damp track in open woodland, locally frequent, 18 Jul 2012, F. Verloove 9564 (BR, LG, GENT). — DEPARTMENT SAÔNE-ET-LOIRE: À l'E de Verdun-sur-le-Doubs, forêt, *Caricetum pendulae*, argileux-humide, Jul 1974, M. Bournérias 577bis (P); Dissey [Mouthier-en-Bresse], Forêt de Dissey, station inédite nouveau pour ce département, 16 Jul 2004, G. H. Parent 04/122 (BR).

Scirpus georgianus

FRANCE: DEPARTMENT BAS-RHIN: Plobsheim, près Strasbourg, plan d'eau installé entre 1965 et 1970, Sep 1975, Kapp *s.n.* (P); Plobsheim, près de Strasbourg, bord du plan d'eau installé 1965 et 1970 (ouvriers méridionaux et nord-africains), Sep 1975, Institut Botanique de Strasbourg *s.n.* (P). — DEPARTMENT MARNE: Ecueil, Jul 1987, C. Worms *s.n.* (BR); [Saint-Martin-d'Ablois], Forêt de Bruguy, chemin forestier humide, Jul 1989, L. Delvosalle *s.n.* (BR) (see Worms 1985). – A claim from Saint-Imoges (also in the department Marne) possibly also belongs here (Lambinon & Duvigneaud 1988; Worms 1989).

Ecology and invasive status in France

Scirpus hattorianus appears to be confined in France to rather damp, shady spots in (open) woodland, nearly always by tracks. This is exactly the sort of habitat where it is typically found in the NE United States. More rarely it has also been observed in forest margins and by ponds. All locations are on sandy or clayish substratum.

Most species of the *Scirpus atrovirens* complex are weedy and their small achenes are well adapted for accidental transport by humans and animals. Once established, plants are readily dispersed to adjoining suitable habitats. Bizot & Parent (2005) suggested that “*S. atrovirens*” was spread by hunters. All French authors have remarked on the abundance of the species in most of its locations: “très nombreuses touffes”, “abondante”, “constante progression”, etc. (see Litzler 2000; Prost 2000; Bizot & Parent 2005). Bizot & Parent (*l.c.*) described its presence in the departments Jura and Saône-et-Loire in an area that by now covers approximately 100 km². In the sense of Richardson & al. (2000), *S. hattorianus* is in France doubtless an invasive species (*i.e.* naturalized plants that produce reproductive offspring at considerable distances from parent plants and thus

have the potential to spread over a considerable area). However, it is unknown whether or not native, vulnerable species are outcompeted by its presence. Additional research on its ecology is therefore required.

Acknowledgements

Dr Ernie Schuyler (Philadelphia, U.S.A.) is thanked for confirming the identity of our collection from Bournonville, Iris Van der Beeten (Meise, Belgium) is thanked for preparing the SEM photographs, and Henry Engledow (Meise, Belgium) is thanked for preparing the distribution map. Jean-Marc Tison (l'Isle d'Abeau, France) and Benoît Toussaint (Bailleul, France) are acknowledged for providing information about French localities of *Scirpus hattorianus*. The curator of the herbarium of the Muséum National d'Histoire Naturelle, Paris (P) is thanked for sending on loan relevant herbarium collections. Finally, two anonymous reviewers are thanked for their comments on an earlier draft of this paper.

References

- Bizot A. & Parent G. H. 2005: Trois plantes américaines du nord-est de la France: 1. *Carex vulpinoidea* Michx., 2. *Glyceria striata* (Lam.) Hitchcock, 3. *Scirpus atrovirens* Willd. – Bull. Soc. Hist. Nat. Ardennes **94**: 24–34.
- Bornand C. & Clerc C. 2011: *Scirpus pendulus* Muhl.: un néophyte nord-américain et ses différents noms dans les flores suisses. – Bull. Cercle Vaud. Bot. **40**: 87–90.
- DeFilipps R. A. 1980: *Scirpus* L. – Pp. 277–280 in: Tutin T. G., Heywood V. H., Burges N. A., Moore D. M., Valentine D. H., Walters S. M. & Webb D. A. (ed.), Flora europaea **5**. – Cambridge: University Press.
- Giot J.-L. 2004: Notes et réflexions militaro-botaniques concernant *Scirpus atrovirens* trouvé sur le camp militaire de Marche. – Les Barbouillons **216**: 1–3.
- Kerguélén M. 1999: Index synonymique de la flore de France. – Published at: <http://www2.dijon.inra.fr/flore-france/index.htm> [accessed 13 Feb 2012].
- Kiffe K. 1998: *Scirpus* L. – Pp. 463–464 in: Wisskirchen R. & Häupler H. (ed.), Standardliste der Farn- und Blütenpflanzen Deutschlands. – Stuttgart: Ulmer.
- Lambinon J. & Duvigneaud J. 1988: Mises au point taxonomiques et nomenclaturales et additions floristiques (documents pour une 4e édition de la “Nouvelle Flore de la Belgique et des régions voisines”). – Dumortiera **40**: 5–22.
- Lannoy M. 2004: Présence de *Scirpus atrovirens* Willd. à Marche-en-Famenne. – Les Barbouillons **216**: 4–6.
- Lenski H. 1985: Ein Fund von *Scirpus cyperinus* (L.) Kunth in West-Niedersachsen. – Göttinger Florist. Rundbr. **19**: 43.
- Litzler P. 2000: Nouveaux loisirs botaniques de retraite. – Monde Pl. **468**: 12.
- Mabberley D. J. 2008: Mabberley's plant-book, ed. 3. – Cambridge: University Press.
- Makino T. 1933: A contribution to the knowledge of the flora of Nippon – J. Jap. Bot. **8**: 44–45.
- Pignotti L. 2003: *Scirpus* L. and related genera (*Cyperaceae*) in Italy. – *Webbia* **58**: 281–400.
- Prost J.-F. 1983: 1970–1979: dix années d'herborisations dans le Jura. – Monde Pl. **354**: 5–6 [erroneously as *Scirpus holoschoenus*].
- Prost J.-F. 2000: *Scirpus atrovirens* Willd. dans le Jura. – Monde Pl. **470**: 15.
- Prost J.-F. 2001: Catalogue des plantes vasculaires de la chaîne jurassienne. – Lyon: Société Linéenne de Lyon [erroneously as *Scirpus holoschoenus*].
- Richardson D. M., Pyšek P., Rejmánek M., Barbour M. G., Panetta F. D. & West C. J. 2000: Naturalization and invasion of alien plants: concepts and definitions. – *Diversity and Distrib.* **6**: 93–107.
- Riomet L. B. & Bournérias M. 1952–1961: Flore de l'Aisne. Catalogue des plantes vasculaires du département. – Laon: Société d'Histoire Naturelle de l'Aisne.
- Schnittler M. & Niedbala J. 1993: Zur Unterscheidung von *Scirpus atrovirens* Willd., *Scirpus sylvaticus* L. und ähnlichen Cyperaceen. – Florist. Rundbr. **27**(1): 25–31.
- Schuyler E. 1961: Evidence for the hybrid origin of *Scirpus peckii*. – *Rhodora* **63**: 237–243.
- Schuyler E. 1967: A taxonomic revision of North American leafy species of *Scirpus*. – Proc. Acad. Nat. Sci. Philadelphia **119**: 295–323.
- Strong M. T. 1994: Taxonomy of *Scirpus*, *Trichophorum*, and *Schoenoplectus* (*Cyperaceae*) in Virginia. – *Bartonia* **58**: 29–68.




Fig. 2. Distribution by department of *Scirpus hattorianus* (dark grey) and *S. georgianus* (pale grey) in France.

- Tela Botanica 2012+ [continuously updated]: Tela Botanica. Le réseau de la botanique francophone. Botanique. *Scirpus atrovirens* Willd. – Published at <http://www.tela-botanica.org/eflore/BDNFF/4.02/nn/61550/chorologie> [accessed 13 Feb 2012].
- Thiers B. 2014+ [continuously updated]: Index herbariorum: a global directory of public herbaria and associated staff. – New York Botanical Garden: published at <http://sweetgum.nybg.org/ih/> [accessed 20 Feb 2014].
- Tucker G. C. 1987: The genera of *Cyperaceae* in the southeastern United States. – *J. Arnold Arbor.* **68**: 361–445.
- Verloove F. 2006: Catalogue of neophytes in Belgium (1800–2005). – *Scripta Bot. Belg.* **39**.
- Verloove F. & Lambinon J. 2011: The non-native vascular flora of Belgium: new combinations and a new variety. – *New J. Bot.* **1**: 38–42.
- Whittemore A. T. & Schuyler A. E. 2002: *Scirpus* L. – Pp. 8–21 in: Flora of North America Editorial Committee (eds.), Flora of North America north of Mexico **23**. – New York & Oxford: Oxford University Press.
- Worms C. 1985: Notes floristiques sur le département de la Marne (France: district champenois et district du nord-est de l'île de France). – *Nat. Mosana* **38(1)**: 18–21.
- Worms C. 1989: Notes floristiques sur la région de Reims (France: district champenois et district du nord-est de l'île de France). – *Nat. Mosana* **42(3)**: 102–108.
- Zelnik I. 2004: *Scirpus georgianus* Harper – a new species in Slovenian flora and character species of the association *Dactylorhiza majalis-Scirpetum georgiani* ass. nova. – *Hacquetia* **3(2)**: 94–105.