



## **The Taxonomic Status of Jean-Jacques Kieffer's Type Specimens of Afrotropical Cecidomyiidae (Diptera)**

Authors: Harris, Keith M., and Dorchin, Netta

Source: African Invertebrates, 53(1) : 169-174

Published By: KwaZulu-Natal Museum

URL: <https://doi.org/10.5733/afin.053.0110>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## The taxonomic status of Jean-Jacques Kieffer's type specimens of Afrotropical Cecidomyiidae (Diptera)

Keith M. Harris<sup>1</sup> and Netta Dorchin<sup>2</sup>

<sup>1</sup>81 Linden Way, Ripley, Woking, Surrey, GU23 6LP, UK; kmharris@lineone.net

<sup>2</sup>Department of Zoology, Tel Aviv University, Tel Aviv, 69978 Israel; ndorchin@post.tau.ac.il

### ABSTRACT

Holotypes and syntypes of cecidomyiid genera and species described from the Seychelles and East Africa by J.-J. Kieffer in the early 20<sup>th</sup> century are held in the collections of the Natural History Museum, London and the Muséum national d'Histoire naturelle, Paris, France. All are specimens caught in flight, so without associated biological data and most are unidentifiable females. Many are in poor condition and inadequate for detailed study and are of little assistance in advancing understanding of the Afrotropical Cecidomyiidae fauna. Following examination of all extant specimens, 16 genera and 59 species are listed and assigned to *nomina dubia*. Six other species that are represented by holotypes or syntypes that still have intact genitalia (which may possibly be recognised in future revisionary studies), are also listed.

KEY WORDS: Afrotropical, Cecidomyiidae, gall midges, Jean-Jacques Kieffer, *nomina dubia*, types.

### INTRODUCTION

R.J. Gagné (1994), in an account of the gall midges of the Neotropical Region, provided a brief biography of Jean-Jacques Kieffer (1857–1925), in which he noted that Kieffer published 470 scientific papers, including taxonomic studies of the Cecidomyiidae, Ceratopogonidae, Chironomidae and parasitic Hymenoptera. Gagné also noted that no collection survived at Kieffer's work place in Bitche, Lorraine, France, an observation that was first made by F.W. Edwards in 1938 and confirmed by later searchers (including Gagné himself in 1987). It is, therefore, surprising that types of almost all of the Cecidomyiidae described by Kieffer from tropical Africa have survived, either in Muséum national d'Histoire naturelle, Paris (MNHN) or in the Natural History Museum, London (BMNH). These specimens were collected by two expeditions: the Percy Sladen Trust expedition to the Indian Ocean in 1905, which collected in the Seychelles, and the Alluaud-Jeannel expedition in 1911–12, which collected in East Africa (Kenya and Tanzania). Kieffer's accounts of this material were published in various papers (Kieffer 1911, 1912, 1913a–f). All of the specimens were caught in flight, so nothing is known regarding their biology, and most are females, without associated males. These types are, therefore, of very little value in understanding the Afrotropical Cecidomyiidae fauna, but the genus and species names based on them were validly published under provisions of the International Code of Zoological Nomenclature (ICZN 1999).

The specimens in the MNHN were originally stored in alcohol in small glass tubes and those in the BMNH were staged on pins. In the late 1970s they were mostly mounted on microscope slides by KMH and examined during preparation of the Cecidomyiidae entries for the *Catalogue of the Diptera of the Afrotropical Region* (Harris 1980). No revisionary studies were conducted at that time, but some generic names were treated as junior synonyms and some species were reassigned. In particular, the following genera were placed in synonymy: *Properrisia* Kieffer, 1912, *Sphaerolauthia* Kieffer, 1912 and *Nanolauthia* Kieffer, 1913 with *Ledomyia* Kieffer, 1894; *Parasphondylia* Kieffer, 1913 with *Schizomyia* Kieffer, 1889; *Tristephanus* Kieffer, 1913 with *Karshomyia* Felt, 1908;

*Plagioidiplosis* Kieffer, 1913 with *Trisopsis* Kieffer, 1912; and *Baeomyza* Kieffer, 1913 and *Jeannellomyia* Kieffer, 1913 with *Stomatosema* Kieffer, 1904. Most of the other genera described by Kieffer from the Seychelles and East Africa were listed by Harris (1980) as valid taxa, pending further study. We have now reassessed the situation during preparation of the chapter on Cecidomyiidae for the forthcoming *Manual of Afrotropical Diptera* and have decided that most of these genera and species named by Kieffer, which have remained unrecognised for almost a century, are unlikely to be recognised with any degree of certainty now or in the future and are best treated as *nomina dubia*.

This is especially true of the many monotypic genera erected by Kieffer on the basis of single, caught, rather than reared, specimens (mostly females); a practice that was acceptable at that time, but not acceptable today. In most cases, Kieffer provided minimal generic descriptions, or no descriptions at all, and almost no comparative diagnoses and, although the type specimens have survived, they are mostly damaged and in poor condition. We, therefore, formally designate all of the genera and species listed below as *nomina dubia*. A few male specimens have distinctive genitalia that may or may not be diagnostic of species. We list them separately, but do not assign them to *nomina dubia*.

Full biographical and other information for all of the species and genera dealt with in this paper is provided in *A Catalog of the Cecidomyiidae (Diptera) of the World* (Gagné 2004) and a digital update, which is available on open access (Gagné 2010). For consistency and ease of amendment, we have listed species names in the format of this catalogue, with the species name preceding the generic assignment. Many cecidomyiid genera erected in sole or joint authorship by Kieffer, mainly in earlier publications, which include Afrotropical representatives, are retained as recognised taxa, notably *Baldratia* Kieffer, 1897; *Clinodiplosis* Kieffer, 1894; *Dicrodiplosis* Kieffer, 1895; *Endaphis* Kieffer, 1896; *Lauthia* Kieffer, 1912; *Ledomyia* Kieffer, 1894; *Lestodiplosis* Kieffer, 1894; *Mitodiplosis* Kieffer, 1914; *Ochnephila* Kieffer, 1914; *Orseolia* Kieffer & Massalongo, 1902; *Procontarinia* Kieffer & Cecconi, 1906; *Schizomyia* Kieffer, 1889; *Stefaniola* Kieffer, 1913; *Stomatosema* Kieffer, 1904; *Trisopsis* Kieffer, 1912; *Trotteria* Kieffer, 1902; and *Zeuxidiplosis* Kieffer, 1904.

#### TAXONOMY

The following listed genera and species are here regarded as *nomina dubia*:

##### Subfamily Micromyinae Rondani, 1856

*parvula* Kieffer, 1919: 198. (*Tricampylomyza*). Cameroon, holotype ♀ (lost).

##### Subfamily Porricondylinae Kieffer, 1913

*abbreviata* Kieffer, 1913e: 114. (*Epidosis*). Kenya, holotype ♀ (MNHN).

*anemotis* Kieffer, 1913e: 114. (*Epidosis*). Kenya, holotype ♀ (MNHN).

*monticola* Kieffer, 1913e: 113. (*Epidosis*). Kenya, holotype ♀ (MNHN).

*mahensis* Kieffer, 1911: 327. (*Prowinnwertzia*). Seychelles, holotype ♀ (BMNH).

##### Subfamily Cecidomyiinae Newman, 1834

*APLECUS* Kieffer, 1913e: 105. Type species: *Aplecus validus* Kieffer (by original designation). Monotypic genus.

- BAEODIPLOSI* Kieffer, 1913e: 111. Type species: *Baeodiplosis tropicalis* Kieffer (by original designation). Monotypic genus.
- CACOPLECUS* Kieffer, 1913e: 112. Type species: *Cacoplecus brevitarsis* Kieffer (by original designation). Monotypic genus.
- CHAETODIPLOSI* Kieffer, 1913e: 103. Type species: *Chaetodiplosis tropica* Kieffer (by original designation). Genus based on one caught female; the later addition from Nigeria of *C. gymnastica* Collin, 1922: 507, is currently in unplaced Cecidomyiidi (Gagné 2010).
- CONODIPLOSI* Kieffer, 1913e: 102. Type species: *Conodiplosis dolichopsis* Kieffer (by original designation). Monotypic genus.
- CTENODIPLOSI* Kieffer, 1913e: 109. Type species: *Ctenodiplosis elongatus* Kieffer (by original designation). Two species included.
- HOLODIPLOSI* Kieffer, 1913e: 103. Type species: *Holodiplosis alticola* Kieffer (by original designation).
- LASIODIPLOSI* Kieffer, 1913a: 232, as replacement name for *Lepidodiplosis* Kieffer, 1912: 2 (preoccupied by *Lepidodiplosis* Kieffer, 1911: 319). Type species: *Chrysodiplosis homotomus* Kieffer, 1911: 319. Monotypic genus.
- LEPIDOBREMIA* Kieffer, 1913b: x. Type species: *Bremia mahensis* Kieffer (by original designation). Monotypic genus.
- LEPIDODIPLOSI* Kieffer, 1911: 319. Type species: *Lepidodiplosis mahensis* Kieffer (by subsequent designation of Kieffer 1913f: 227).
- ORTHODIPLOSI* Kieffer, 1913e: 109. Type species: *Orthodiplosis macrurus* Kieffer (by original designation). Monotypic genus.
- PACHYDIPLOSI* Kieffer, 1913e: 108. Type species: *Pachydiplosis apricans* Kieffer (by original designation). Monotypic genus. This genus was subsequently widely and incorrectly used in applied entomology to accommodate the Asian rice gall midge, *Orseolia oryzae* (Wood-Mason, 1889) (see Krishnaiah 2004).
- PERODIPLOSI* Kieffer, 1913e: 100. Type species: *Perodiplosis coronata* Kieffer (by original designation). Monotypic genus.
- PLANODIPLOSI* Kieffer, 1911: 323. Type species: *Planodiplosis insularis* Kieffer (by monotypy). Monotypic genus.
- PLECOPHORUS* Kieffer, 1913e: 104. Type species: *Plecophorus fuscocinctus* Kieffer (by original designation). Monotypic genus.
- TRISSODIPLOSI* Kieffer, 1913e: 101. Type species: *Trissodiplosis longicornis* Kieffer (by original designation). Monotypic genus.
- albiventris* Kieffer, 1913e: 93 (*Asphondylia*). Kenya, holotype ♀ (MNHN).
- alluaudi* Kieffer, 1913e: 94 (*Trisopsis*). Kenya, holotype ♀ (MNHN).
- alticola* Kieffer, 1913e: 104 (*Holodiplosis*). Kenya, holotype ♂ (MNHN). Type species of *Holodiplosis* Kieffer.
- apricans* Kieffer, 1913e: 109 (*Pachydiplosis*). Kenya, holotype ♀ (MNHN). Type species of *Pachydiplosis* Kieffer.
- aurofulgens* Kieffer, 1911: 316 (*Ledomylia*). Seychelles, syntypes 1♂, 1♀ (MNHN). Assigned to *Lauthia* Kieffer by Gagné (2004).
- brevipalpis* Kieffer, 1913e: 92 (*Asphondylia*). Kenya, holotype ♂ (MNHN).
- brevitarsis* Kieffer, 1913e: 113 (*Cacoplecus*). Kenya, syntypes 4♀ (MNHN). Type species of *Cacoplecus* Kieffer.

- calopus* Kieffer, 1911: 323 (*Lepidodiplosis* Kieffer, 1911). Seychelles, holotype ♀ (BMNH).
- ciliatipennis* Kieffer, 1911: 325 (*Lestodiplosis*). Seychelles, holotype ♀ (BMNH).
- cincticornis* Kieffer, 1913e: 99 (*Lestodiplosis*). Tanzania, holotype ♀ (MNHN).
- coronatus* Kieffer, 1913e: 100 (*Perodiplosis*). Kenya, holotype ♀ (MNHN). Type species of *Perodiplosis* Kieffer.
- crassinervis* Kieffer, 1913e: 92 (*Asphondylia*). Kenya, holotype ♀ (MNHN).
- cubitalis* Kieffer, 1913e: 88 (*Lasioptera*). Kenya, holotype ♀ (MNHN).
- dolichopsis* Kieffer, 1913e: 102 (*Conodiplosis*). Kenya, holotype ♀ (MNHN). Type species of *Conodiplosis* Kieffer.
- dubia* Kieffer, 1913e: 95 (*Lamprodiplosis*). Tanzania, holotype ♀ (MNHN).
- elongata* Kieffer, 1913e: 110 (*Ctenodiplosis*). Kenya, holotype ♀ (MNHN). Type species of *Ctenodiplosis* Kieffer.
- filipes* Kieffer, 1911: 320 (*Lepidodiplosis* Kieffer, 1911). Seychelles, holotype ♀ (BMNH).
- fuscocinctus* Kieffer, 1913e: 105 (*Plecophorus*). Tanzania, holotype ♀ (MNHN). Type species of *Plecophorus* Kieffer.
- gracilipes* Kieffer, 1913c: 41 (*Jeannellomyia*). Kenya, holotype ♀ (MNHN). Type species of *Jeannellomyia* Kieffer.
- gracilis* Kieffer, 1913e: 97 (*Coprodiplosis*). Tanzania, holotype ♀ (MNHN).
- heterotarsus* Kieffer, 1913e: 113 (*Cacoplecus*). Kenya, holotype ♀ (MNHN).
- hirta* Kieffer, 1913e: 96 (*Coprodiplosis*). Kenya, holotype ♀ (MNHN).
- homotomus* Kieffer, 1911: 319 (*Chrysodiplosis*). Seychelles, holotype ♂ (BMNH).
- imicola* Kieffer, 1913e: 104 (*Holodiplosis*). Kenya, holotype ♂ (MNHN).
- insularis* Kieffer, 1911: 323 (*Planodiplosis*). Seychelles, holotype ♂ (BMNH). Type species of *Planodiplosis* Kieffer.
- insularum* Kieffer, 1911: 326 (*Clinodiplosis*). Seychelles, syntypes 1♂, 1♀ (BMNH).
- longicauda* Kieffer, 1913e: 99 (*Coprodiplosis*). Tanzania, syntypes 2♀ (MNHN).
- longicornis* Kieffer, 1913e: 101 (*Trissodiplosis*). Kenya, holotype ♀ (MNHN). Type species of *Trissodiplosis* Kieffer.
- lumiensis* Kieffer, 1913e: 97 (*Coprodiplosis*). Tanzania, holotype ♀ (MNHN).
- macrurus* Kieffer, 1913e: 109 (*Orthodiplosis*). Kenya, holotype ♀ (MNHN). Type species of *Orthodiplosis* Kieffer.
- mahensis* Kieffer, 1911: 326 (*Bremia*). Seychelles, holotype ♂ not found (BMNH). Type species of *Lepidobremia* Kieffer.
- mahensis* Kieffer, 1911: 320 (*Lepidodiplosis* Kieffer, 1911). Seychelles, syntypes 1♂, 1♀ (BMNH). Type species of *Lepidodiplosis* Kieffer, 1911.
- monilicornis* Kieffer, 1911: 318 (*Ledomyia*). Seychelles, holotype ♂ (BMNH). Assigned to *Lauthia* Kieffer by Gagné (2004).
- monticola* Kieffer, 1913e: 90 (*Baeomyza*). Tanzania, holotype ♀ (MNHN).
- nana* Kieffer, 1911: 325 (*Trisopsis*). Seychelles, syntypes 2♂, 2♀ (BMNH).
- nocticolor* Kieffer, 1911: 322 (*Lepidodiplosis* Kieffer, 1911). Seychelles, holotype ♂ (BMNH).
- nympha* Kieffer, 1913e: 96 (*Coprodiplosis*). Tanzania, holotype ♀ (MNHN).
- pedicellatum* Kieffer, 1913e: 90 (*Baeomyza*). Kenya, holotype ♀ (MNHN).
- pictipennis* Kieffer, 1913e: 91 (*Asphondylia*). Kenya, holotype ♂ (MNHN).
- scotti* Kieffer, 1911: 326 (*Clinodiplosis*). Seychelles, holotype ♀ (BMNH).

- scottianus* Kieffer, 1911: 324 (*Lestodiplosis*). Seychelles, holotype ♀ (BMNH).  
*setosa* Kieffer, 1913e: 92 (*Asphondylia*). Kenya, holotype ♀ (MNHN).  
*seychellensis* Kieffer, 1911: 320 (*Lepidodiplosis* Kieffer, 1911). Seychelles, syntypes 8♀ (BMNH).  
*squamosipes* Kieffer, 1911: 321 (*Lepidodiplosis* Kieffer, 1911). Seychelles, holotype ♂ (BMNH).  
*styloptera* Kieffer, 1911: 317 (*Ledomyia*). Seychelles, holotype ♀ (BMNH).  
*subrotunda* Kieffer, 1913e: 110 (*Ctenodiplosis*). Tanzania, holotype ♀ (MNHN).  
*subsessile* Kieffer, 1913e: 89 (*Baeomyza*). Kenya, holotype ♀ (MNHN).  
*tritonus* Kieffer, 1913e: 98 (*Coprodiplosis*). Tanzania, holotype ♀ (MNHN).  
*tropica* Kieffer, 1913e: 103 (*Chaetodiplosis*). Tanzania, holotype ♀ (MNHN). Type species of *Chaetodiplosis* Kieffer.  
*tropicalis*, Kieffer, 1913e: 112 (*Baeodiplosis*). Tanzania, holotype ♀ (MNHN). Type species of *Baeodiplosis* Kieffer.  
*tropicalis* Kieffer, 1913e: 98 (*Coprodiplosis*). Tanzania, holotype ♀ (MNHN).  
*tuberosa* Kieffer, 1913e: 88 (*Lasioptera*). Tanzania, holotype ♀ (MNHN).  
*validus* Kieffer, 1913e: 105 (*Aplecus*). Tanzania, holotype ♀ (MNHN). Type species of *Aplecus* Kieffer.  
*variicornis* Kieffer, 1913e: 93 (*Parasphondylia*). Kenya, holotype ♀ (MNHN). Type species of *Parasphondylia* Kieffer.

Types of six of the Cecidomyiinae species described by Kieffer from the Seychelles and East Africa have male genitalia that are reasonably intact and may possibly be recognised again in future revisionary studies. These are not here assigned to *nomina dubia*, but are listed below.

- atricauda* Kieffer, 1913e: 107 (*Tristephanus*). Kenya, syntypes 2♂, 1♀ (MNHN). Type species of *Tristephanus* Kieffer, treated as a junior synonym of *Karshomyia* Felt by Harris (1980).  
*brevipalpis* Kieffer, 1913e: 108 (*Microplecus*). Tanzania, holotype ♂ (MNHN). Type species of *Microplecus* Kieffer, 1913.  
*mahensis* Kieffer, 1911: 317 (*Ledomyia*). Seychelles, holotype ♂ (BMNH). Assigned to *Lauthia* Kieffer by Gagné (2004).  
*monticola* Kieffer, 1913e: 111 (*Clinodiplosis*). Tanzania, holotype ♂ (MNHN).  
*pulchricornis* Kieffer, 1911: 318 (*Chrysodiplosis*). Seychelles, holotype ♂ (BMNH). Type species of *Chrysodiplosis* Kieffer, 1911.  
*squamosus* Kieffer, 1911: 323 (*Nanodiplosis*). Seychelles, holotype ♂ (BMNH). Type species of *Nanodiplosis* Kieffer, 1913.

The study of Afrotropical Cecidomyiidae is still at a very early stage, probably with most species and genera unknown and a great need for thorough new systematic studies.

#### ACKNOWLEDGEMENTS

This paper is dedicated to the memory of Brian Stuckenberg, who, as early as the 1960s, was aware of the need for new and comprehensive taxonomic studies of the virtually unknown African Cecidomyiidae. We are grateful to Mathias Jaschhof (Senckenberg Deutsches Entomologisches Institut, Müncheberg, Germany) for advising on the status of the Micromyinae and Porricondylinae species and to Erica McAlister (BMNH) and the late Loïc Matile (formerly MNHN) for enabling examination of relevant specimens in their care. Mathias

Jaschhof, Peter Kolesik and Raymond J. Gagné (United States Department of Agriculture, Washington DC, USA) also kindly commented on the draft manuscript.

## REFERENCES

- COLLIN, J.E. 1922. Description of a new genus and two new species of Cecidomyiidae [*sic*], and six new species of acalyptate Muscidae (Ephydriidae and Milichiidae). *Transactions of the Entomological Society of London* **1921**: 504–517.
- GAGNÉ, R.J. 1994. *The gall midges of the Neotropical Region*. Ithaca, New York: Cornell Univ. Press.
- 2004. A catalog of the Cecidomyiidae (Diptera) of the world. *Memoirs of the Entomological Society of Washington* **25**: 1–408.
- 2010. *Update for a catalog of the Cecidomyiidae of the world*. Digital version 1. ([http://www.ars.usda.gov/SP2UserFiles/Place/12754100/Gagne\\_2010\\_World\\_Catalog\\_Cecidomyiidae.pdf](http://www.ars.usda.gov/SP2UserFiles/Place/12754100/Gagne_2010_World_Catalog_Cecidomyiidae.pdf); accessed 26/07/2011)
- HARRIS, K.M. 1980. Family Cecidomyiidae. In: Crosskey, R.W., ed., *Catalogue of the Diptera of the Afro-tropical Region*. London: British Museum (Natural History), pp. 238–251.
- INTERNATIONAL COMMISSION ON ZOOLOGICAL NOMENCLATURE (ICZN). 1999. *International Code of Zoological Nomenclature*. London: International Trust for Zoological Nomenclature.
- KIEFFER, J.J. 1911. No. XIV. – Diptera, Cecidomyiidae, der Seychellen-Inseln aus der Sammlung von H. Scott. *Transactions of the Linnean Society of London, Zoology* (2) **14**: 315–330.
- 1913a. Nouvelle contribution à la connaissance des cécidomyies. *Marcellia* **11** (1912): 219–235.
- 1913b. Neue Gallmücken-Gattungen. Bitsch, France. In: Trotter, A., *Bibliographia e recensionei. Marcellia* **11** (1912): x–xi.
- 1913c. Insectes Diptères I. Chironomidae et Cecidomyiidae [*sic*]. In: *Voyage de Ch. Alluaud et R. Jeannel en Afrique orientale (1911–1912), Résultats scientifiques*. Paris: A. Schultz, pp. 32–43.
- 1913d. Glanures diptérologiques. *Bulletin de la Société d'Histoire naturelle de Metz* **28**: 45–55.
- 1913e. Cécidomyies de l'Afrique Orientale. *Bulletin de la Société d'Histoire naturelle de Metz* **28**: 87–114.
- 1913f. Diptera. Fam. Cecidomyiidae. In: Wytzman, P., ed., *Genera Insectorum*. Fasc. 152. Bruxelles: V. Verteneuil & L. Desmet, pp. 1–346.
- 1919. Microdiptères d'Afrique. *Bulletin de la Société d'Histoire naturelle de l'Afrique du Nord* **10**: 191–207.
- KRISHNAIAH, K. 2004. Rice gall midge, *Orseolia oryzae* — an overview. In: Bennett, J., Bentur, J.S., Pasalu, I.C. & Krishnaiah, K., eds, *New approaches to gall midge resistance in rice. Proceedings of the International Workshop, 22–24 November 1998, Hyderabad, India*. Los Baños, Philippines: International Rice Research Institute and Indian Council of Agricultural Research, pp. 1–5.