



## **The Complete Chloroplast Genome of *Capsicum frutescens* (Solanaceae)**

Authors: Shim, Donghwan, Raveendar, Sebastin, Lee, Jung-Ro, Lee, Gi-An, Ro, Na-Young, et al.

Source: Applications in Plant Sciences, 4(5)

Published By: Botanical Society of America

URL: <https://doi.org/10.3732/apps.1600002>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## THE COMPLETE CHLOROPLAST GENOME OF *CAPSICUM FRUTESCENS* (SOLANACEAE)<sup>1</sup>

DONGHWAN SHIM<sup>2,5</sup>, SEBASTIN RAVEENDAR<sup>3,5</sup>, JUNG-RO LEE<sup>3</sup>, GI-AN LEE<sup>3</sup>, NA-YOUNG RO<sup>3</sup>,  
YOUNG-AH JEON<sup>3</sup>, GYU-TAEK CHO<sup>3</sup>, HO-SUN LEE<sup>3</sup>, KYUNG-HO MA<sup>3</sup>, AND JONG-WOOK  
CHUNG<sup>3,4,6</sup>

<sup>2</sup>Department of Forest Genetic Resources, Korea Forest Research Institute, Suwon 441-350, Republic of Korea; <sup>3</sup>National Agrobiodiversity Center, National Institute of Agricultural Science, Rural Development Administration, Jeonju 54874, Republic of Korea; and <sup>4</sup>Department of Industrial Plant Science and Technology, Chungbuk National University, Cheongju, Chungbuk 28644, Republic of Korea

- **Premise of the study:** We report the complete sequence of the chloroplast genome of *Capsicum frutescens* (Solanaceae), a species of chili pepper.
- **Methods and Results:** Using an Illumina platform, we sequenced the chloroplast genome of *C. frutescens*. The total length of the genome is 156,817 bp, and the overall GC content is 37.7%. A pair of 25,792-bp inverted repeats is separated by small (17,853 bp) and large (87,380 bp) single-copy regions. The *C. frutescens* chloroplast genome encodes 132 unique genes, including 87 protein-coding genes, 37 transfer RNA (tRNA) genes, and eight ribosomal RNA (rRNA) genes. Of these, seven genes are duplicated in the inverted repeats and 12 genes contain one or two introns. Comparative analysis with the reference chloroplast genome revealed 125 simple sequence repeat motifs and 34 variants, mostly located in the noncoding regions.
- **Conclusions:** The complete chloroplast genome sequence of *C. frutescens* reported here is a valuable genetic resource for *Capsicum* species.

**Key words:** *Capsicum frutescens*; chili pepper; chloroplast genome; next-generation sequencing; Solanaceae.

A chloroplast is an organelle with its own genome encoding a number of chloroplast-specific components (Sugiura et al., 1998). Owing to its tractable size and high level of conservation, the chloroplast genome can be used to characterize genetic relationships among species. Furthermore, plant taxonomists have widely adopted the sequence variability of two loci in land plants, consisting of portions of the chloroplast *rbcL* and *matK* genes, as an effective DNA barcode (Vijayan and Tsou, 2010). Chloroplast DNA contains many of the genes necessary for proper functioning of the organelle. The analysis of chloroplast DNA sequences has proven useful in studying plant evolution (Shaw et al., 2007), and the field of chloroplast genome characterization is growing rapidly (Timmis et al., 2004). The size of the genome, which has been determined for a number of plants and algae, ranges from 85 to 292 kbp. The complete DNA sequences of several different chloroplast genomes of plants and algae have been reported. Many chloroplast DNAs contain two inverted repeats (IRs), which separate a large single-copy region (LSC) from a small single-copy region (SSC) (Palmer

and Thompson, 1982). The IRs vary in length from 4 to 25 kbp (Robinson et al., 2009).

*Capsicum frutescens* L. (Solanaceae), a name that is generally applied to all cultivated peppers in the United States, is also known as *C. annuum* L. (Smith and Heiser, 1951). Cultivars of *C. frutescens* can be annual or short-lived perennial plants. The flowers have a greenish white or greenish yellow corolla, and they are either insect- or self-pollinated. The fruit is usually very pungent, growing to 1.0–8.0 cm long and 0.6–3.0 cm in diameter (Smith and Heiser, 1951). The fruit is typically pale yellow as it matures to a bright red, but it can also be other colors (Heiser and Smith, 1953; Stummel and Bosland, 2006). More recently, *C. frutescens* has been bred to produce ornamental strains with a large number of erect peppers growing in colorful ripening patterns (Stummel and Bosland, 2006). *Capsicum frutescens* likely originated in South or Central America (Heiser, 1979; Clement et al., 2010) and spread quickly throughout the tropical and subtropical regions in this area, where it still grows wild today (Purseglove, 1976). It is also believed that *C. frutescens* is the ancestor of *C. chinense* Jacq. (Bosland, 1996; Basu et al., 2003).

In this study, using Illumina technology, the complete chloroplast genome of *C. frutescens* was sequenced, assembled, annotated, and mined for simple sequence repeat (SSR) markers and for single-nucleotide polymorphism (SNP) and insertion/deletion (indel) variants. The resultant data have been made publicly available as a resource for genetic information for *Capsicum* L. species, which will facilitate investigations into

<sup>1</sup>Manuscript received 12 January 2016; revision accepted 5 April 2016.

This study was performed with the support of the Research Program for Agricultural Science and Technology Development (Project no. PJ008623), National Institute of Agricultural Science, Rural Development Administration, Republic of Korea.

<sup>5</sup>These authors contributed equally to this work.

<sup>6</sup>Author for correspondence: jwchung73@chungbuk.ac.kr

doi:10.3732/apps.1600002

*Applications in Plant Sciences* 2016 4(5): 1600002; <http://www.bioone.org/loi/apps> © 2016 Shim et al. Published by the Botanical Society of America. This work is licensed under a Creative Commons Attribution License (CC-BY-NC-SA).




Fig. 1. Gene map of the *Capsicum frutescens* chloroplast genome. Genes drawn inside the circle are transcribed clockwise, while those drawn outside are transcribed counterclockwise (marked with two arrows). Different functional gene groups are color-coded. Variation in the GC content of the genome is shown in the middle circle. The map was drawn using OGDRAW version 1.2 (Lohse et al., 2007).

genetic variation and phylogenetic relationships of closely related *Capsicum* species.

## METHODS AND RESULTS

For this study, *C. frutescens* seeds (accession no. IT158639) were obtained from the National Agrobiodiversity Center, Rural Development Administration, Republic of Korea. Seeds were germinated and grown in a greenhouse,

fresh leaves were collected from 40-d-old seedlings, and DNA was extracted using a DNeasy Plant Mini Kit (QIAGEN, Valencia, California, USA) according to the manufacturer's instructions to construct chloroplast DNA libraries. An Illumina paired-end DNA library (average insert size of 500 bp) was constructed using the Illumina TruSeq library preparation kit following the manufacturer's instructions (Illumina, San Diego, California, USA).

The library was sequenced with 2 × 300 bp on the MiSeq instrument at LabGenomics (<http://www.labgenomics.co.kr/>). Prior to chloroplast de novo assembly, low-quality sequences (quality score < 20; Q20) were filtered out, and the remaining high-quality reads were assembled using the CLC Genome

TABLE 1. SSR candidates of the *Capsicum frutescens* chloroplast genome.

SSR type	SSR abundances	Percentage abundance (%)
Dinucleotide		
TA/AT	9	7.2
Trinucleotide		
TTC/TCT/CTT	9	7.2
TTA/TAT/ATT	23	18.4
Tetranucleotide		
TTTG/TTGT/TGTT	9	7.2
TCTT/CTTT/TTTC	9	7.2
ATAA/TAAA/AAAT	16	12.8
AATT/ATTA/TTAA	9	7.2
AAAT/AATA/ATAA	19	15.2
Pentanucleotide		
TTTTA/TTTAT/TTATT	11	8.8
TTATT/TATTT/ATTTT	11	8.8
Total	125	100

Assembler (version beta 4.6; CLC bio, Aarhus, Denmark) with a minimum overlap size of 200 bp and maximum bubble size of 50 bp for the de Bruijn graph. Chloroplast contigs were selected from the initial assembly by performing a BLAST (version 2.2.31) search against the reference chloroplast genome of *C. annuum* (GenBank accession NC\_018552) using CLC software with parameters of 0.5 for fraction, 0.8 for similarity, and 200–600 bp of overlap size (Jo et al., 2011). The selected chloroplast contigs were merged into a total of four contigs, and iterative contig extensions were performed to construct a complete *C. frutescens* chloroplast genome by mapping raw reads to the contigs. Dual Organellar GenoMe Annotator (DOGMA; Wyman et al., 2004) and CpGAVAS (Liu et al., 2012) were used to annotate the chloroplast genome. All transfer RNA (tRNA) genes were amended with tRNAscan-SE (Lowe and Eddy, 1997). OGDRAW (Lohse et al., 2007) was used to produce a map of the genome.

Sputnik software (Cardle et al., 2000) was used to find the SSR markers present in the chloroplast genome of *C. frutescens*. It uses a recursive algorithm to search for repeats with lengths between two and five, and finds perfect, compound, and imperfect repeats. Sputnik has been applied for SSR identification in many species, including *Arabidopsis* and barley (Cardle et al., 2000). To identify SNP and indel variants in the *C. frutescens* chloroplast genome, we used BWA (Li and Durbin, 2009) with ‘mem’ command line options ‘-k19 -w100 -d100 -r1.5 -y20 -c500 -D0.5 -W0 -m50’ and SAMtools (Li et al., 2009) software with ‘mpileup’ command line options ‘-uf -d250 -q0 -e20 -h100 -L250 -m1 -o40.’ A more detailed method is described at <http://samtools.sourceforge.net/mpileup.shtml>.

Illumina paired-end (2 × 300 bp) sequencing produced a total of 8,272,114 paired-end reads, with an average fragment length of 256 bp, which were then analyzed to generate 1,796,432,923 bp of sequence. The results contain 31,772,592 mapped nucleotides with an average coverage of 202× on the chloroplast genome. Contig alignment and scaffolding based on paired-end data resulted in a complete circular *C. frutescens* chloroplast genome sequence (Fig. 1). The chloroplast genome of *C. frutescens* has been deposited in GenBank (accession no. KR078312; National Center for Biotechnology Information [NCBI]). It has a total length of 156,817 bp and is composed of an LSC of 87,380 bp, two IRs of 25,792 bp, and an SSC of 17,853 bp. The overall GC content of the *C. frutescens* chloroplast genome is 37.7%, with the IRs having a higher GC content (43.1%) than the LSC (35.7%) and SSC (32.0%) due to the presence of GC-rich ribosomal RNA (rRNA) genes. The *C. frutescens* chloroplast genome encodes 132 unique genes (Appendix 1), including 87 protein-coding genes, 37 tRNA genes, and eight rRNA genes. Seven of these genes are duplicated in the IR regions, nine genes (*rps16*, *atpF*, *rpoC1*, *petB*, *petD*, *rpl16*, *rpl2*(IR), *ndhB*(IR), *ndhA*) and six tRNA genes contain one intron, and two genes (*clpP*, *rps12*) and one *ycf* (*ycf3*) contain two introns (Fig. 1).

The size of the *C. frutescens* chloroplast genome (156,817 bp) was larger than reported for *C. frutescens* species such as *C. annuum* var. *glabriusculum* (Dunal) Heiser & Pickersgill (GenBank accession no. KJ619462) and *C. annuum* (GenBank accession no. NC\_018552). The lengths of the LSC and IRs in *C. frutescens* differed from those in the other two species and contributed to the variation of chloroplast genome size. For example, the *C. frutescens* chloroplast genome was 36 bp longer than the reported *C. annuum* chloroplast genome and 205 bp longer than the *C. annuum* var. *glabriusculum* chloroplast genome. Furthermore, the SSC and IR regions of *C. frutescens* were 3 and 9 bp longer, respectively, and the LSC region was 14 bp shorter and 167 bp longer, respectively, than those of the previously reported chloroplast genomes. The average GC content in the *C. frutescens* chloroplast genome is 37.7%, similar to other *Capsicum* species.

The organization and gene order of the *Capsicum* chloroplast genome exhibited the general chloroplast genome structure seen in angiosperms (Sugiura, 1992). The *Capsicum* chloroplast genome contains 132 genes (Appendix 2), of which there were eight rRNA genes, 37 tRNA genes, 21 ribosomal subunit genes (12 small subunit and nine large subunit), and four DNA-directed RNA polymerase genes. Forty-six genes were involved in photosynthesis, of which 11 encoded subunits of the NADH-oxidoreductase, seven for photosystem I, 15 for photosystem II, six for the cytochrome *b<sub>6</sub>/f* complex, six for different subunits of ATP synthase, and one for the large chain of ribulose biphosphate carboxylase/oxygenase (RuBisCO). Five genes were involved in different functions, and three genes were of unknown function. As shown in Fig. 1 and Appendix 2, genome organization appeared to be more conserved with unique gene sequences, as discovered previously in *Capsicum* species (Jo et al., 2011; Zeng et al., 2014; Raveendar et al., 2015a). However, in this newly determined chloroplast genome, we found 132 predicted genes and size variations were observed in the IR and LSC regions.

TABLE 2. SNP markers of the *Capsicum frutescens* chloroplast genome.

No.	REF ( <i>C. annuum</i> )	ALT ( <i>C. frutescens</i> )	Coding region	QUAL	Region
1	T	C	noncoding region	222	LSC
2	A	T	noncoding region	222	LSC
3	T	G	noncoding region	4.77	LSC
4	T	C	noncoding region	19.1	LSC
5	G	T	noncoding region	222	LSC
6	G	A	noncoding region	222	LSC
7	C	A	noncoding region	222	LSC
8	T	A	noncoding region	222	LSC
9	A	G	noncoding region	222	LSC
10	G	C	gene ( <i>petA</i> )	222	LSC
11	A	G	gene ( <i>petA</i> )	222	LSC
12	C	A	gene ( <i>petA</i> )	222	LSC
13	C	A	gene ( <i>petA</i> )	222	LSC
14	A	T	noncoding region	222	LSC
15	G	A	gene ( <i>rpl32</i> )	164	SSC
16	G	T	gene ( <i>rpl32</i> )	124	SSC
17	A	T	noncoding region	222	SSC
18	T	G	noncoding region	222	SSC

Note: ALT = alteration; LSC = large single-copy; QUAL = Phred-scaled quality score; REF = reference; SSC = small single-copy.



TABLE 3. Indel markers of the *Capsicum frutescens* chloroplast genome.

No.	REF ( <i>C. annuum</i> )	ALT ( <i>C. frutescens</i> )	Coding region	QUAL	Region
1	ATTTTTTTTT	ATTTTTTTTTT, ATTTTTTTTTTT	noncoding region	48.5	LSC
2	TAAAAAA	TAAAAAA	noncoding region	178	LSC
3	GAAAAAAAAAAAAAAAA	GAAAAAAAAAAAAAAAAA, GAAAAAAAAAAAA, GAAAAAAAAAAAAAAAAA	noncoding region	18.5	LSC
4	CTTTTTT	CTTTTTT	noncoding region	152	LSC
5	TCAACTCATTTTA	T	noncoding region	214	LSC
6	TATTTTTAATTTTAATTTT AATATATTTTAATTTTAAT ATAAATAAATAATTTTAAT ATATTAATATAAATAAATA AATAAT	TAT	noncoding region	217	LSC
7	CAAAAAAAAA	CAAAAAAAAAA, CAAAAAAAAAAA, CAAAAAAAAA	noncoding region	65.5	LSC
8	ATTTTTTTTTT	ATTTTTTTTTT, ATTTTTTTTTTT	noncoding region	68.5	LSC
9	TAAAAAAAA	TAAAAAAAAA, TAAAAAAAAAA	noncoding region	48.5	LSC
10	TCCGGTAAAGACTCCGG TAAAGACTCCGGTAAAGAC	TCCGGTAAAGACGCCGTAAGA CTCCGGTAAAGACTCCGGTAAAGAC	gene ( <i>rpl20</i> )	218	LSC
11	GTTTTTTTTT	GTTTTTTTTT, GTTTTTTTTTT	noncoding region	94.5	LSC
12	GAAAAAAA	GAAAAAA	noncoding region	66.5	LSC
13	GAAAAAAA	GAAAAAAA, GAAAAAAA	noncoding region	90.5	LSC
14	CTTTT	CTTTT	noncoding region	214	LSC
15	ATTCTTATTTTTT	ATTATTTTTT	gene ( <i>rps19</i> )	203	LSC
16	TCCCCC	TCCCCC	noncoding region	185	SSC

Note: ALT = alteration; LSC = large single-copy; QUAL = Phred-scaled quality score; REF = reference; SSC = small single-copy.

A total of 125 potential SSRs motifs were identified, located mostly in the noncoding regions (Table 1); of these, the majority belonged to tetranucleotide (50%) and trinucleotide (26%) repeats. All other types of SSRs, such as di- and pentanucleotide motifs, were relatively low (25%). The majority of tetranucleotide SSRs had the AAAT/AATA/ATAA motif, followed by those with the ATAA/TAAA/AAAT motif; the TTTG/TTGT/TGTT, TCTT/CTTT/TTTC, and AATT/ATTA/TTAA motifs were found with similar frequency (7.2%). Two different repeats—those with the TTTTA/TTTAT/TTATT and TTATT/TATTT/ATTTT motifs—were identified among pentanucleotide SSRs. The TTC/TCT/CTT and TTA/TAT/ATT motifs were identified among the trinucleotide SSRs, but only the TA/AT motif was identified for the dinucleotide SSRs (Table 1). In total, 125 potential SSRs motifs were identified in the 156.8-kb sequence of the *Capsicum* chloroplast genome. Hence, the observed frequency of SSRs motifs was approximately one per 1250 bp of chloroplast genome.

Comparison of the *C. frutescens* chloroplast genome sequence with the reference chloroplast sequence of *C. annuum* revealed a total of 34 mutations (18 SNPs and 16 indels), with 15 of these variants involving more than one nucleotide (Table 2 and 3). Among the detected variants, six SNPs and two indels were observed in the coding region of the chloroplast genome. Among these SNPs and indels, there were 29 and five mutations located in the LSC and SSC regions, respectively. These molecular markers will facilitate studies of genetic diversity, population genetic structure, and sustainable conservation for *C. frutescens*.

The size of the *C. frutescens* chloroplast genome identified here is more closely related to that of *C. annuum* var. *glabriusculum* reported previously (Raveendar et al., 2015b). Moreover, the *C. frutescens* chloroplast genome has similar genome organization, gene order, gene sizes, and GC content, with only SNPs/indels variation. It has been reported that *C. annuum* var. *glabriusculum* is considered the wild parental species of the cultivated *C. annuum* (Votava et al., 2002; Aguilar-Meléndez et al., 2009; González-Jara et al., 2011).

## CONCLUSIONS

We provide here the complete chloroplast genome sequence of *C. frutescens*, a cultivated pepper in the United States. Availability of this sequence and the recently determined *C. annuum* chloroplast genome sequence (GenBank accession no. NC\_018552) enables us to assess genome-wide mutational dynamics within the genus *Capsicum*. The chloroplast genome

possesses similar genome organization, gene order, gene sizes, and GC content, with only SNPs/indels variation having been revealed. It is difficult to get accurate phylogenies and effective species discrimination using a small number of plastid genes in evolutionarily young lineages (Ruhsam et al., 2015). Therefore, complete plastid genome sequencing provides a solution to this problem. Availability of this sequence can enable researchers to design conserved primers to sequence new genomic regions that could provide useful phylogenetic information for closely related species. Moreover, the structural details of this *C. frutescens* chloroplast genome join the growing database of *Capsicum* species, which can facilitate investigations into gene expression and genetic variation of these crop species.

## LITERATURE CITED

- AGUILAR-MELÉNDEZ, A., P. L. MORRELL, M. L. ROOSE, AND S. C. KIM. 2009. Genetic diversity and structure in semiwild and domesticated chiles (*Capsicum annuum*; Solanaceae) from Mexico. *American Journal of Botany* 96: 1190–1202.
- BASU, S. K., A. K. DE, AND A. DE. 2003. *Capsicum*: Historical and botanical perspectives. In A. K. De [ed.], *Capsicum*: The genus *Capsicum*, 1–15. CRC Press, London, United Kingdom.
- BOSLAND, P. W. 1996. Capsicums: Innovative uses of an ancient crop. In J. Janick [ed.], *Progress in new crops*, 479–487. ASHS Press, Arlington, Virginia, USA.
- CARDLE, L., L. RAMSAY, D. MILBOURNE, M. MACAULAY, D. MARSHALL, AND R. WAUGH. 2000. Computational and experimental characterization of physically clustered simple sequence repeats in plants. *Genetics* 156: 847–854.
- CLEMENT, C. R., M. DE CRISTO-ARAÚJO, G. COPPENS D'EECKENBRUGGE, A. ALVES PEREIRA, AND D. PICAÑO-RODRIGUES. 2010. Origin and domestication of native Amazonian crops. *Diversity (Basel)* 2: 72–106.
- GONZÁLEZ-JARA, P., A. MORENO-LETELIER, A. FRAILE, D. PIÑERO, AND F. GARCÍA-ARENAL. 2011. Impact of human management on the genetic variation of wild pepper, *Capsicum annuum* var. *glabriusculum*. *PLoS ONE* 6: e28715.

- HEISER, C. B. 1979. Origins of some cultivated new world plants. *Annual Review of Ecology and Systematics* 10: 309–326.
- HEISER, C. B., AND P. G. SMITH. 1953. The cultivated *Capsicum* peppers. *Economic Botany* 7: 214–227.
- JO, Y. D., J. PARK, J. KIM, W. SONG, C. G. HUR, Y. H. LEE, AND B. C. KANG. 2011. Complete sequencing and comparative analyses of the pepper (*Capsicum annuum* L.) plastome revealed high frequency of tandem repeats and large insertion/deletions on pepper plastome. *Plant Cell Reports* 30: 217–229.
- LI, H., AND R. DURBIN. 2009. Fast and accurate short read alignment with Burrows–Wheeler transform. *Bioinformatics* 25: 1754–1760.
- LI, H., B. HANDSAKER, A. WYSOKER, T. FENNEL, J. RUAN, N. HOMER, G. MARTH, ET AL. 2009. The Sequence Alignment/Map format and SAMtools. *Bioinformatics* 25: 2078–2079.
- LIU, C., L. SHI, Y. ZHU, H. CHEN, J. ZHANG, X. LIN, AND X. GUAN. 2012. CpGAVAS, an integrated web server for the annotation, visualization, analysis, and GenBank submission of completely sequenced chloroplast genome sequences. *BMC Genomics* 13: 715.
- LOHSE, M., O. DRECHSEL, AND R. BOCK. 2007. OrganellarGenomeDRAW (OGDRAW): A tool for the easy generation of high-quality custom graphical maps of plastid and mitochondrial genomes. *Current Genetics* 52: 267–274.
- LOWE, T. M., AND S. R. EDDY. 1997. tRNAscan-SE: A program for improved detection of transfer RNA genes in genomic sequence. *Nucleic Acids Research* 25: 955–964.
- PALMER, J. D., AND W. F. THOMPSON. 1982. Chloroplast DNA rearrangements are more frequent when a large inverted repeat sequence is lost. *Cell* 29: 537–550.
- PURSEGLOVE, J. 1976. The origins and migrations of crops in tropical Africa. In J. R. Harlan [ed.], *Origins of African plant domestication*, 291–309. Mouton De Gruyter, The Hague, The Netherlands.
- RAVEENDAR, S., Y.-A. JEON, J.-R. LEE, G.-A. LEE, K. J. LEE, G.-T. CHO, K.-H. MAA, S.-Y. LEE, AND J.-W. CHUNG. 2015a. The complete chloroplast genome sequence of Korean landrace “Subicho” pepper (*Capsicum annuum* var. *annuum*). *Plant Breeding and Biotechnology* 3: 88–94.
- RAVEENDAR, S., Y. W. NA, J. R. LEE, D. SHIM, K. H. MA, S. Y. LEE, AND J. W. CHUNG. 2015b. The complete chloroplast genome of *Capsicum annuum* var. *glabriusculum* using Illumina sequencing. *Molecules (Basel, Switzerland)* 20: 13080–13088.
- ROBINSON, D. G., H. ARONSSON, AND A. S. SANDELIUS. 2009. The chloroplast: Interactions with the environment. Springer, Berlin, Germany.
- RUHSAM, M., H. S. RAI, S. MATHEWS, T. G. ROSS, S. W. GRAHAM, L. A. RAUBESON, W. MEI, ET AL. 2015. Does complete plastid genome sequencing improve species discrimination and phylogenetic resolution in *Araucaria*? *Molecular Ecology Resources* 15: 1067–1078.
- SHAW, J., E. B. LICKY, E. E. SCHILLING, AND R. L. SMALL. 2007. Comparison of whole chloroplast genome sequences to choose noncoding regions for phylogenetic studies in angiosperms: The tortoise and the hare III. *American Journal of Botany* 94: 275–288.
- SMITH, P. G., AND C. B. HEISER JR. 1951. Taxonomic and genetic studies on the cultivated peppers, *Capsicum annuum* L. and *C. frutescens* L. *American Journal of Botany* 38: 362–368.
- STUMMEL, J. R., AND P. BOSLAND. 2006. Ornamental pepper: *Capsicum annuum*. In N. O. Anderson [ed.], *Flower breeding and genetics: Issues, challenges, and opportunities for the 21st century*, 561–599. Springer, Dordrecht, The Netherlands.
- SUGIURA, M. 1992. The chloroplast genome. *Plant Molecular Biology* 19: 149–168.
- SUGIURA, M., T. HIROSE, AND M. SUGITA. 1998. Evolution and mechanism of translation in chloroplasts. *Annual Review of Genetics* 32: 437–459.
- TIMMIS, J. N., M. A. AYLIFFE, C. Y. HUANG, AND W. MARTIN. 2004. Endosymbiotic gene transfer: Organelle genomes forge eukaryotic chromosomes. *Nature Reviews. Genetics* 5: 123–135.
- VIJAYAN, K., AND C. H. TSOU. 2010. DNA barcoding in plants: Taxonomy in a new perspective. *Current Science (India)* 99: 1530–1541.
- VOTAVA, E., G. NABHAN, AND P. BOSLAND. 2002. Genetic diversity and similarity revealed via molecular analysis among and within an in situ population and ex situ accessions of chiltepin (*Capsicum annuum* var. *glabriusculum*). *Conservation Genetics* 3: 123–129.
- WYMAN, S. K., R. K. JANSEN, AND J. L. BOORE. 2004. Automatic annotation of organellar genomes with DOGMA. *Bioinformatics* 20: 3252–3255.
- ZENG, F. C., C. W. GAO, AND L. Z. GAO. 2014. The complete chloroplast genome sequence of American bird pepper (*Capsicum annuum* var. *glabriusculum*). *Mitochondrial DNA Part A* 27: 724–726.

APPENDIX 1. General features of the *Capsicum frutescens* chloroplast genome.

Chloroplast genome feature	Quantity
Genome size (bp)	156,817
GC content (%)	37.7
Total no. of genes	132
Protein-coding genes	87
rRNA genes	8
tRNA genes	37
Genes duplicated in IR regions	7
Total introns	12
Single intron (gene)	9
Double introns (gene)	3
Single intron (tRNA)	6

APPENDIX 2. Genes present in the *Capsicum frutescens* chloroplast genome.

Chloroplast genome feature	Gene products
Photosystem I	<i>psaA, psaB, psaC, psaI, psaJ, ycf3<sup>2</sup>, ycf4</i>
Photosystem II	<i>psbA, psbB, psbC, psbD, psbE, psbF, psbH, psbI, psbJ, psbK, psbL, psbM, psbN, psbT, psbZ</i>
Cytochrome <i>b<sub>6</sub>/f</i>	<i>petA, petB<sup>1</sup>, petD<sup>1</sup>, petG, petL, petN</i>
ATP synthase	<i>atpA, atpB, atpE, atpF<sup>1</sup>, atpH, atpI</i>
RuBisCO	<i>rbcL</i>
NADH oxidoreductase	<i>ndhA<sup>1</sup>, ndhB<sup>1,3</sup>, ndhC, ndhD, ndhE, ndhF, ndhG, ndhH, ndhI, ndhJ, ndhK</i>
Large subunit ribosomal proteins	<i>rpl2<sup>1,3</sup>, rpl14, rpl16<sup>1</sup>, rpl20, rpl22, rpl23<sup>3</sup>, rpl32, rpl33, rpl36</i>
Small subunit ribosomal proteins	<i>rps2, rps3, rps4, rps7<sup>3</sup>, rps8, rps11, rps12<sup>2,3,4</sup>, rps14, rps15, rps16<sup>1</sup>, rps18, rps19</i>
RNA polymerase	<i>rpoA, rpoB, rpoC1<sup>1</sup>, rpoC2</i>
Unknown function protein-coding gene	<i>ycf1<sup>3</sup>, ycf2<sup>3</sup>, ycf15<sup>3</sup></i>
Other genes	<i>accD, ccsA, cemaA, clpP<sup>2</sup>, matK</i>
Ribosomal RNAs	<i>rrn16<sup>3</sup>, rrn23<sup>3</sup>, rrn4.5<sup>3</sup>, rrn5<sup>3</sup></i>
Transfer RNAs	<i>trnA-UGC<sup>1,3</sup>, trnC-GCA, trnD-GUC, trnE-UUC, trnF-GAA, trnG-UCC<sup>1</sup>, trnG-GCC, trnH-GUG, trnI-CAU<sup>3</sup>, trnI-GAU<sup>1,3</sup>, trnK-UUU<sup>1</sup>, trnL-UAA<sup>1</sup>, trnL-UAG, trnL-CAA<sup>3</sup>, trnM-CAU, trnM-CAU, trnN-GUU<sup>3</sup>, trnP-UGG, trnQ-UUG, trnR-ACG<sup>3</sup>, trnR-UCU, trnS-GCU, trnS-GGA, trnS-UGA, trnT-GGU, trnT-UGU, trnV-UAC<sup>1</sup>, trnV-GAC<sup>3</sup>, trnW-CCA, trnY-GUA</i>

<sup>1</sup> Gene containing a single intron.

<sup>2</sup> Gene containing two introns.

<sup>3</sup> Two gene copies in IRs.

<sup>4</sup> Transsplicing gene.