

Comparative study of the effects of high- versus low-dose zinc oxide in the diet with or without probiotic supplementation on weaning pigs' growth performance, nutrient utilization, fecal microbes, noxious gas discharges, and fecal score

Authors: Biswas, Sarbani, Kim, Min Hee, Park, Jae Hong, Kim, Yangseon, and Kim, In Ho

Source: Canadian Journal of Animal Science, 103(1) : 33-43

Published By: Canadian Science Publishing

URL: <https://doi.org/10.1139/cjas-2022-0080>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Comparative study of the effects of high- versus low-dose zinc oxide in the diet with or without probiotic supplementation on weaning pigs' growth performance, nutrient utilization, fecal microbes, noxious gas discharges, and fecal score

Sarbani Biswas ^a, Min Hee Kim^a, Jae Hong Park^a, Yangseon Kim^b, and In Ho Kim ^a

^aDepartment of Animal Resource and Science, Dankook University, No. 29 Anseodong, Cheonan, Choongnam 330-714, South Korea;

^bDepartment of Research and Development, Center for Industrialization of Agriculture and Livestock Microorganisms, Jeongseup 56212, South Korea

Corresponding author: In Ho Kim (email: inhokim@dankook.ac.kr)

Abstract

This study was conducted to determine the effects of high- versus low-dose (3000 vs. 300) zinc oxide (ZnO) in combination with or without a probiotic complex (0.1%) on weaned piglet production efficiency, nutrient absorption, fecal bacterial counts, noxious gas emissions, and fecal score. A 42-day experiment included 180 crossbred weaned piglets [Duroc × (Yorkshire × Landrace); 28 days old; 6.61 ± 1.29 kg] and four dietary treatments. An HZ (high ZnO) diet increased body weight at week 6, average daily gain at week 3, week 6, and overall period, and gain-to-feed ratio (G:F) at week 3 compared with an LZ (low ZnO) diet. G:F tended to increase with the LZP (LZ with probiotic) diet compared with the HZP (HZ with probiotic) diet at week 1. *Escherichia coli* count decreased by HZ diet compared with the LZ diet. In addition, *E. coli* count decreased and *Lactobacillus* count increased with the HZP diet compared with the LZP diet. There was no effect of treatment on nutrient digestibility, noxious gas emission, and fecal score. No interactive effect was seen between ZnO and probiotic. Therefore, high-dose ZnO inclusion improved growth performance and probiotic addition improved fecal microbiota, but no synergistic effect was found from ZnO and probiotic complex interaction.

Key words: zinc oxide, probiotic complex, performance, weaning pig

Introduction

Weaning is a challenging stage for pigs, comprising nutritional, psychological, ecological, microbiological, and autoimmune challenges, which can lead to financial losses due to reduced growth performance, inefficiency of feed utilization, diarrhea, and impairment of gastrointestinal function and physiology (Lalles 2008). In previous decades, antibiotic growth enhancers were used clinically on farms to improve the productive efficiency and health welfare, and perhaps to prevent infections of weaned pigs. Antibiotics have various beneficial effects on such mammals, but their use must be restricted as it leads to the expansion of susceptibility by several different strains of bacteria (Pluske 2013; Liao and Nyachoti 2017). So, numerous alternative feed additives have been explored to substitute antibiotics. As a consequence, the usage of feed additives as an immune response and gut microbial modulator has become a routine practice, enhancing pig health and productivity during the postweaning phase (Lalles et al. 2007).

As a vital mineral ingredient, zinc (Zn) has a huge stabilizing influence on the body's improvement and evolution, immunological function, and gastrointestinal integrity. In weaned pigs, several Zn sources (e.g., montmorillonite zinc oxide (ZnO), nanoparticle ZnO, and coated ZnO) have been added to diets to promote intestinal immune function and structure and to modulate intestinal microbiota (Hu et al. 2012; Shen et al. 2014). For several years, therapeutic levels of ZnO have indeed been utilized to relieve weaning stress, enhance nutritional digestibility, and strengthen immune function as well as boost productive efficiency (Upadhyaya et al. 2018). Inclusion of a modified form of ZnO in weaned pig diets improved growth performance and average daily gain (ADG) during the study period (Cho et al. 2015). In addition, ZnO pharmacological dosages (2500–3500 mg/kg) have been shown to significantly reduce diarrhea, improve pig growth responses (Hill et al. 2001), and reduce noxious gas production (Shi et al. 2019). A ZnO dose below therapeutic levels (maximum of 150 mg/kg) is identified as a growth enhancer

for piglets (European Communities 2003). ZnO nanoparticles in animal diets have been demonstrated to boost body weight (BW), feed intake (FI), and feed conversion ratio (FCR) in poultry (Zhao et al. 2014), but had no effect on growth performance (Li et al. 2016) in piglets. In addition, it has been observed that an LZ diet (low ZnO; approximately 250 or 500 ppm) has no influence on the growth of newborn piglets (Hollis et al. 2005). The postulated mechanism for increased growth performance was related to improvements in gastrointestinal function and structure, antioxidant properties, mucosal immune strength, and the elimination of diarrhea in weaned piglets (Trckova et al. 2015; Zhu et al. 2017). However, the extensive use of ZnO in weanling pigs raised concerns about excretion of ZnO to the environment and its link to the development of antibiotic-resistant microorganisms (Slifierz et al. 2015; Jensen et al. 2016).

Probiotics are single or combined populations of living microbes that provide favorable health benefits to the animals when consumed. Antimicrobial substances produced by probiotics include amino substances, H₂O₂, antimicrobials, and natural polymers, which can all hinder the activity of pathogenic bacteria (Chaucheyras-Durand and Durand 2010). Organic acids (lactic acid, acetic acid) are the most prevalent compounds formed by probiotics, and they lower the pH in the gastrointestinal tract, creating an unfavorable environment for pathogenic organisms. Probiotics also boost susceptibility to enteric pathogens by competing for attachment sites and nutrients in the intestine (Boirivant and Strober 2007). Many studies have found that adding probiotics to weaned pigs' diets can enhance growth performance (Lan et al. 2017; Dumitru et al. 2021) and nutrient digestibility (Zhao and Kim 2015), and reduce diarrhea occurrence (Giang et al. 2012; Dumitru et al. 2021). Probiotics can improve immunity (Yu et al. 2008) and reduce fecal poisonous gas discharges (Zhao and Kim 2015), resulting in fewer pollutants in the atmosphere in weaned pigs (Ferket et al. 2002). For animals (Bedford et al. 2014), probiotics have been suggested as a potential antibacterial growth booster and antibiotic substitute. To improve the effect of adding ZnO to the diet, a probiotic compound was added to either low- or high-dose ZnO. In weanling pigs, Shi et al. (2019) discovered that low-dose ZnO mixed with probiotic compounds enhanced growth efficiency and had no negative effect on nutrient absorption, immune parameters, or the occurrence of gastroenteritis. However, dietary ZnO (1200 ppm) inclusion with probiotic compounds compares favorably with the ZnO pharmacological (3000 ppm) level regarding economic growth, dietary digestion, gastrointestinal microbiota, poisonous gas discharges, and fecal index in piglets (Wang et al. 2021). So, probiotic supplementation can help to reduce the dose of ZnO applied in weaning pigs to improve the production performance and gut health. Considering the above benefits, we hypothesized that the combination of ZnO and probiotic compound will increase productive efficiency, nutrient absorption, intestinal microbiota population, noxious gas emission, and fecal score of weaned pigs by acting synergistically.

Therefore, the objective of this study was to determine the effect of high and low doses of ZnO with or without a 0.1% probiotic complex (*Bacillus coagulans*, *B. licheniformis*, *B. subtilis*,

and *Clostridium butyricum*) inclusion on growth performance, fecal bacterial enumeration, noxious gas emissions, and fecal score of weaning pigs. A second objective was to estimate whether there would be any interactive effect between ZnO and probiotic.

Materials and methods

Ethical statement

All animal management and experimental procedures were approved by the Institutional Animal Care and Use Committee of Dankook University, Republic of Korea (Approval No. DK-1-1937).

Source of probiotics

The probiotic compound used in the current experiment was acquired from a commercial company (Synerbig Co. Ltd., Seoul, South Korea), named SynerZymeF10. This product is assured to comprise at least 1×10^{12} CFU/kg *B. coagulans*, 5×10^{11} CFU/kg *B. licheniformis*, 1×10^{12} CFU/kg *B. subtilis*, and 1×10^{11} CFU/kg *C. butyricum*, which is a combination of *B. coagulans*, *B. licheniformis*, *B. subtilis*, and *C. butyricum* spray-dried spores.

Experimental design and animal management

In a 42-day trial, 180 crossbred weanling piglets (Yorkshire \times Landrace \times Duroc; 6.61 ± 1.29 kg; 28 days of age) were used. According to starting BW and sex, total pigs were arbitrarily allocated to one of four nutritional treatments (nine duplicate pens, each including two gilts and three barrows). The nutritional treatment groups were TRT1, positive control: basal diet added with 3000 ppm ZnO (HZ, high ZnO); TRT2, HZ + 0.1% probiotic complex (HZP); TRT3, negative control (NC): basal diet added with 300 ppm ZnO (LZ); and TRT4, LZ + 0.1% probiotic complex (LZP). The trial period was allocated into three phases: days 0–7 of age (phase 1), days 8–21 of age (phase 2), and days 22–42 of age (phase 3). The diets (Tables 1–3) were designed to meet or surpass the National Research Council (NRC)'s nutritional requirements (NRC 2012). All pigs were kept in a biologically maintained room with a mechanical aeration system that provided a 0.26 m \times 0.53 m area for each pig. All over the experiment, each pen was supplied with a self-feeder along with a nipple drinker made of stainless steel to permit the piglets ad libitum access to feed and water. Twelve hours of artificial light for each day was automatically adjusted. The room's ambient temperature was maintained at around 30 °C during the first week, and afterward the temperature was reduced by 1 °C per week.

Growth performance and nutrient absorption

On initial, 7th, 21th, and 42nd days of the trial, BWs of individual pigs were measured and feed intake was documented on a pen basis during the research to examine ADG, average daily feed intake (ADFI), and gain-to-feed ratio (G:F). Chromium oxide (Cr₂O₃, 0.20%; Samchun Pure Chemical Co., Ltd., Gyeonggi, South Korea) was included in the diet as an indigestible indicator 1 week prior to fecal assembly. On day 42, fecal specimens were collected via rectal massage from

Table 1. Experimental diet composition on a fed basis.

Item	Phase 1 (days 0–7 of age)			
	TRT1	TRT2	TRT3	TRT4
Ingredients (%)				
Corn	39.32	39.32	40.04	40.04
Soya bean meal	16.22	16.22	16.10	16.10
Fermented soya bean meal	5.00	5.00	5.00	5.00
Spray-dried plasma protein	6.00	6.00	6.00	6.00
Tallow	2.82	2.82	2.56	2.56
Lactose	12.88	12.88	12.88	12.88
Sugar	3.00	3.00	3.00	3.00
Whey protein	11.00	11.00	11.00	11.00
Monocalcium phosphate	0.88	0.88	0.88	0.88
Limestone	1.18	1.18	1.18	1.18
Salt	0.20	0.20	0.20	0.20
Methionine (99%)	0.20	0.20	0.20	0.20
Lysine	0.49	0.49	0.49	0.49
Mineral mix ^a	0.20	0.20	0.20	0.20
Vitamin mix ^b	0.20	0.20	0.20	0.20
Choline (25%)	0.03	0.03	0.03	0.03
Zinc oxide (80%)	0.38	0.38	0.04	0.04
Calculated value				
Crude protein, %	20.00	20.00	20.00	20.00
Metabolizable energy, kcal/kg	3450	3450	3450	3450
Calcium, %	0.80	0.80	0.80	0.80
Phosphorus, %	0.60	0.60	0.60	0.60
Lysine, %	1.60	1.60	1.60	1.60
Methionine, %	0.48	0.48	0.48	0.48
Fat, %	4.52	4.52	4.28	4.28
Lactose, %	20.00	20.00	20.00	20.00
Zinc oxide, ppm	3053	3053	333	333

^aProvided per kg diet: Fe, 100 mg as ferrous sulfate; Cu, 17 mg as copper sulfate; Mn, 17 mg as manganese oxide; I, 0.5 mg as potassium iodide; Se, 0.3 mg as sodium selenite. ^bProvided per kg of diet: vitamin A, 10 800 IU; vitamin D₃, 4000 IU; vitamin E, 40 IU; vitamin K₃, 4 mg; vitamin B₁, 6 mg; vitamin B₂, 12 mg; vitamin B₆, 6 mg; vitamin B₁₂, 0.05 mg; biotin, 0.2 mg; folic acid, 2 mg; niacin, 50 mg; D-calcium pantothenate, 25 mg.

two pigs for each pen (one barrow and one gilt). On a pen basis, they were combined and pooled, and a random specimen was kept in a freezer at -20°C until analyzed. For chemical study, fecal specimens were dried out at 60°C for 72 h, and then finely crushed to a size that could go through a 1 mm screen. By utilizing the method developed by AOAC (2010), diet and fecal samples were examined to regulate dry matter (DM) and nitrogen (N). To determine energy (E) and excreta samples, specimens were taken and put in a calorimeter (Parr Instruments, Moline, IL, USA) to assess thermal combustion in the samples as well as chromium, which was evaluated by atomic absorption spectrophotometry (Shimadzu UV-1201; Shimadzu, Kyoto, Japan). Apparent total tract digestibility (ATTD) was then measured by the following formula:

Digestibility (%) = $[1 - (N_f \times C_d)/(N_d \times C_f)] \times 100$, where N_f represents the nutrient concentration in feces (% DM), N_d represents the nutrient concentration in diet (% DM), C_d represents the chromium concentration in diet (% DM), and C_f represents the chromium concentration in feces (% DM).

Fecal noxious gas emissions

At the end of the study, fresh fecal specimens (300 g) were amassed randomly from each pen (two pigs), mixed evenly, kept in a 2.6 L sealed plastic box with small holes on any side, fixed firmly with tape, and fermented for 7 days to determine ammonia (NH_3) at 25°C , hydrogen sulfide (H_2S), methyl mercaptans, CO_2 , and acetic acid, utilizing a GV-100 gas sampling pump (model GV100S; Gastec Corp., Tokyo, Japan). Various measurement tubes (No. 3L, No. 4LT, and No. 70L; Gastec) were used to detect total mercaptans. On the 8th day, a 100 mL sample was grabbed from the headspace (2 cm) for air passage, and it was sealed again. To understand the creation of a hard crust on the top, the specimen box was shaken manually for 30 s. Finally, the concentrations of NH_3 , H_2S , methyl mercaptan, CO_2 , and acetic acid were evaluated.

Fecal microbiota

Fecal specimens were taken from two pigs from each pen by massaging the rectums and kept on ice for transfer to the research laboratory, where they were instantly tested to de-

Table 2. Experimental diet composition on a fed basis.

Item	Phase 2 (days 8–21 of age)			
	TRT1	TRT2	TRT3	TRT4
Ingredients (%)				
Corn	51.67	51.67	52.39	52.39
Soya bean meal	16.74	16.74	16.62	16.62
Fermented soya bean meal	4.00	4.00	4.00	4.00
Spray-dried plasma protein	3.00	3.00	3.00	3.00
Tallow	2.82	2.82	2.56	2.56
Lactose	7.78	7.78	7.78	7.78
Sugar	3.00	3.00	3.00	3.00
Whey protein	7.00	7.00	7.00	7.00
Monocalcium phosphate	1.08	1.08	1.08	1.08
Limestone	1.20	1.20	1.20	1.20
Salt	0.10	0.10	0.10	0.10
Methionine (99%)	0.15	0.15	0.15	0.15
Lysine	0.65	0.65	0.65	0.65
Mineral mix ^a	0.20	0.20	0.20	0.20
Vitamin mix ^b	0.20	0.20	0.20	0.20
Choline (25%)	0.03	0.03	0.03	0.03
Zinc oxide (80%)	0.38	0.38	0.04	0.04
Calculated value				
Crude protein, %	18.00	18.00	18.00	18.00
Metabolizable energy, kcal/kg	3400	3400	3400	3400
Calcium, %	0.80	0.80	0.80	0.80
Phosphorus, %	0.60	0.60	0.60	0.60
Lysine, %	1.50	1.50	1.50	1.50
Methionine, %	0.40	0.40	0.40	0.40
Fat, %	4.91	4.91	4.67	4.67
Lactose, %	12.00	12.00	12.00	12.00
Zinc oxide, ppm	3054	3054	334	334

^aProvided per kg diet: Fe, 100 mg as ferrous sulfate; Cu, 17 mg as copper sulfate; Mn, 17 mg as manganese oxide; I, 0.5 mg as potassium iodide; Se, 0.3 mg as sodium selenite. ^bProvided per kg of diet: vitamin A, 10 800 IU; vitamin D₃, 4000 IU; vitamin E, 40 IU; vitamin K₃, 4 mg; vitamin B₁, 6 mg; vitamin B₂, 12 mg; vitamin B₆, 6 mg; vitamin B₁₂, 0.05 mg; biotin, 0.2 mg; folic acid, 2 mg; niacin, 50 mg; D-calcium pantothenate, 25 mg.

termine the gastrointestinal microbial flora. Fecal samples were obtained to study the microbiota in the feces. Every sample had 1 g of fecal droppings combined with 9 mL of 1% peptone broth mixed using a vortex stirrer for 1 min. Samples were serially blended from 10^{-1} to 10^{-6} , and were injected by 50 μ L in two selective agar media, MacConkey agar plates (Difco Laboratories, Detroit, MI, USA) and *Lactobacillus* medium III agar plates (Medium 638, DSMZ, Braunschweig, Germany), for the purpose of isolating *Escherichia coli* and *Lactobacillus*. For the *Lactobacillus* bacterial count, the agar plates in the media were incubated in anaerobic environments for 48 h at 39 °C, and incubated anaerobically for 24 h at 37 °C for the *E. coli* counts. Colony counts were then totaled and the outcomes were displayed as log₁₀ transformed data.

Fecal score

Fecal scores were estimated and documented on a pen source at 08:00 and 20:00 h on days 7, 14, 21, 28, 35, and 42. Using the above fecal scoring scheme, the fecal score was cal-

culated as the average of the five pigs in each pen: 1, hard, dry pellet; 2, firm, formed stools; 3, soft, moist stools that maintain shape; 4, soft, unformed stools that adapt to the form of the vessel; and 5, liquids that are watery and pourable.

Statistical analysis

The data were analyzed with the generalized model with a 2 × 2 factorial arrangement of SAS (SAS Institute Inc., Cary, NC, USA), using the pen as a trial unit. Data were log transformed prior to statistical analysis of microbial counts. The data variability was expressed as pooled SE, with $P < 0.05$ being significant and $P < 0.10$ indicating a trend.

Results

Growth performance

Table 4 summarizes the outcomes of the growth performance. When HZ was added to the piglets' diets, BW increased significantly ($P < 0.05$) at week 6 compared with

Table 3. Experimental diet composition on a fed basis.

Item	Phase 3 (days 22–42 of age)			
	TRT1	TRT2	TRT3	TRT4
Ingredients (%)				
Corn	58.48	58.48	59.18	59.18
Soya bean meal	22.60	22.60	22.48	22.48
Fermented soya bean meal	3.00	3.00	3.00	3.00
Tallow	2.77	2.77	2.53	2.53
Lactose	3.18	3.18	3.18	3.18
Sugar	3.00	3.00	3.00	3.00
Whey protein	3.00	3.00	3.00	3.00
Monocalcium phosphate	1.15	1.15	1.15	1.15
Limestone	1.22	1.22	1.22	1.22
Salt	0.10	0.10	0.10	0.10
Methionine (99%)	0.08	0.08	0.08	0.08
Lysine	0.61	0.61	0.61	0.61
Mineral mix ^a	0.20	0.20	0.20	0.20
Vitamin mix ^b	0.20	0.20	0.20	0.20
Choline (25%)	0.03	0.03	0.03	0.03
Zinc oxide (80%)	0.38	0.38	0.04	0.04
Calculated value				
Crude protein, %	18.00	18.00	18.00	18.00
Metabolizable energy, kcal/kg	3350	3350	3350	3350
Calcium, %	0.80	0.80	0.80	0.80
Phosphorus, %	0.60	0.60	0.60	0.60
Lysine, %	1.40	1.40	1.40	1.40
Methionine, %	0.35	0.35	0.35	0.35
Fat, %	5.14	5.14	4.93	4.93
Lactose, %	5.00	5.00	5.00	5.00
Zinc oxide, ppm	3057	3057	337	337

^aProvided per kg diet: Fe, 100 mg as ferrous sulfate; Cu, 17 mg as copper sulfate; Mn, 17 mg as manganese oxide; I, 0.5 mg as potassium iodide; Se, 0.3 mg as sodium selenite. ^bProvided per kg of diet: vitamin A, 10 800 IU; vitamin D₃, 4000 IU; vitamin E, 40 IU; vitamin K₃, 4 mg; vitamin B₁, 6 mg; vitamin B₂, 12 mg; vitamin B₆, 6 mg; vitamin B₁₂, 0.05 mg; biotin, 0.2 mg; folic acid, 2 mg; niacin, 50 mg; D-calcium pantothenate, 25 mg.

when LZ was added to the diet. Similarly, in week 3, week 6, and overall period, the ADG in pigs fed HZ diets increased ($P < 0.05$) compared with those fed LZ diets. Pigs fed HZP diets had a trend of higher G:F ($P < 0.05$) than pigs fed LZP diets. Furthermore, G:F was greater ($P < 0.05$) for the HZ diets at week 3 compared with the LZ diets. Throughout the experiment, the addition of ZnO (HZ vs. LZ) and the probiotic to the diet had no effect on ADFI ($P > 0.05$). In the growth performance measures, no interacting effects between ZnO and probiotic were found.

Nutrient digestibility

The ATTD values for DM, E, and N are summarized in [Table 5](#). When HZ was added to the diets of weaning pigs, there was no significant change in nutritional digestibility ($P > 0.05$) when compared with the LZ diet. Furthermore, there was no significant improvement in nutritional digestibility when probiotics were added to the diet. In terms of nutrient digestibility, no interaction between ZnO and probiotic was discovered.

Fecal noxious gas emissions

As shown in [Table 6](#), the supplementation of HZ into diet had no significant effect ($P > 0.05$) on fecal noxious gas emissions (NH₃, H₂S, and total mercaptans) when compared with LZ diet. Furthermore, the piglet administered a probiotic complex in its diet showed no influence on gas emission ($P > 0.05$). No interaction between ZnO and the probiotic was observed.

Fecal microbiota

[Table 7](#) shows the findings of the fecal microbiota analysis. HZ-supplemented diets had significantly lower ($P < 0.05$) *E. coli* counts as compared with LZ, while failed ($P > 0.05$) to affect *Lactobacillus* enumeration. However, piglets fed the HZP diet showed decreased *E. coli* ($P < 0.05$) and increased ($P < 0.05$) *Lactobacillus* counts compared with the LZP diet. For fecal microbiota, no interactions between probiotic and ZnO were observed.

Table 4. The effect of dietary supplementation of probiotic complex additive on growth performance in weaning pigs.

Items	High ZnO (3000 ppm)		Low ZnO (300 ppm)		SEM	P value		
	Probiotic– (HZ)	Probiotic+ (HZIP)	Probiotic– (LZ)	Probiotic+ (LZIP)		Probiotic effect	ZnO effect	ZnO × probiotic
Body weight, kg								
Initial	6.55	6.55	6.55	6.55	0.002	0.998	0.998	0.998
Week 1	7.94	8.00	8.02	7.95	0.03	0.970	0.887	0.954
Week 3	12.77	12.95	12.87	12.86	0.10	0.893	0.254	0.773
Week 6	23.53ab	24.23a	23.51b	23.95ab	0.24	0.411	0.034	0.853
1 week								
ADG	166	162	166	169	4	0.860	0.355	0.651
ADFI	200	202	203	205	6	0.493	0.942	0.890
G/F	0.911	0.891	0.893	0.903	0.011	0.062	0.185	0.624
3 weeks								
ADG	407ab	412a	396b	398ab	8	0.859	0.045	0.483
ADFI	473	490	484	479	8	0.955	0.455	0.174
G/F	0.841ab	0.842a	0.831b	0.838ab	0.007	0.689	0.015	0.432
6 weeks								
ADG	604ab	624a	592b	612ab	10	0.241	0.025	1.000
ADFI	840	849	843	854	9	0.653	0.229	0.910
G/F	0.719	0.735	0.703	0.717	0.010	0.218	0.184	0.939
Overall								
ADG	461ab	474a	459b	464ab	6	0.3 587	0.019	0.825
ADFI	611	619	616	621	6	0.6 499	0.222	0.649
G/F	0.756	0.763	0.745	0.748	0.008	0.1 886	0.071	0.911

Abbreviations: TRT1, positive control: basal diet added with 3000 ppm ZnO (HZ); TRT2, HZ + 0.1% probiotic complex (HZIP); TRT3, negative control: basal diet added with 300 ppm ZnO (LZ); TRT4, LZ + 0.1% probiotic complex (LZIP); ADG, average daily gain; ADFI, average daily feed intake; G/F, gain-to-feed ratio; SEM, standard error of means.

Table 5. The effect of dietary supplementation of probiotic complex additive on nutrient digestibility in weaning pigs.

Items	High ZnO (3000 ppm)		Low ZnO (300 ppm)		SEM	P value		
	Probiotic– (HZ)	Probiotic+ (HZIP)	Probiotic– (LZ)	Probiotic+ (LZIP)		Probiotic effect	ZnO effect	ZnO × probiotic
Dry matter	81.47	82.40	80.67	81.14	0.81	0.777	0.213	0.397
Nitrogen	78.87	79.74	78.56	78.74	0.66	0.500	0.403	0.654
Energy	79.69	80.66	79.27	79.43	0.65	0.396	0.220	0.542

Abbreviations: TRT1, positive control: basal diet added with 3000 ppm ZnO (HZ); TRT2, HZ + 0.1% probiotic complex (HZIP); TRT3, negative control: basal diet added with 300 ppm ZnO (LZ); TRT4, LZ + 0.1% probiotic complex (LZIP); SEM, standard error of means.

Table 6. The effect of dietary supplementation of probiotic complex additive on gas emission in weaning pigs.

Items	High ZnO (3000 ppm)		Low ZnO (300 ppm)		SEM	P value		
	Probiotic– (HZ)	Probiotic+ (HZIP)	Probiotic– (LZ)	Probiotic+ (LZIP)		Probiotic effect	ZnO effect	ZnO × probiotic
NH ₃	1.4	1.0	1.9	1.6	0.7	0.649	0.472	0.932
H ₂ S	1.9	1.6	2.3	3.0	0.7	0.779	0.221	0.445
Methyl mercaptans	2.3	1.6	4.9	3.5	1.1	0.351	0.058	0.757
CO ₂	700	450	875	725	258	0.424	0.370	0.839
Acetic acid	0.5	0.2	0.9	0.7	0.2	0.258	0.071	0.955

Abbreviations: TRT1, positive control: basal diet added with 3000 ppm ZnO (HZ); TRT2, HZ + 0.1% probiotic complex (HZIP); TRT3, negative control: basal diet added with 300 ppm ZnO (LZ); TRT4, LZ + 0.1% probiotic complex (LZIP); SEM, standard error of means.

Table 7. The effect of dietary supplementation of probiotic complex additive on fecal microbiota in weaning pigs.

Items	High ZnO (3000 ppm)		Low ZnO (300 ppm)		SEM	P value		
	Probiotic– (HZ)	Probiotic+ (HZP)	Probiotic– (LZ)	Probiotic+ (LZP)		Probiotic effect	ZnO effect	ZnO × probiotic
<i>E. coli</i>	6.18 ^b	6.09 ^b	6.26 ^a	6.20 ^a	0.08	0.007	0.002	0.495
<i>Lactobacillus</i>	9.15 ^b	9.23 ^a	9.16 ^b	9.22 ^a	0.05	0.012	0.902	0.638

Abbreviations: TRT1, positive control: basal diet added with 3000 ppm ZnO (HZ); TRT2, HZ + 0.1% probiotic complex (HZP); TRT3, negative control: basal diet added with 300 ppm ZnO (LZ); TRT4, LZ + 0.1% probiotic complex (LZP); SEM, standard error of means.

Fecal score

The results of the piglets' fecal scores are summarized in Table 8. In comparison to LZ, adding HZ to the diet had no significant influence on the fecal score of weaned pigs. Similarly, the addition of the probiotic complex to the piglets' diet had no effect on the piglets' fecal score ($P > 0.05$), and no interacting effects were detected.

Discussion

Feed rejection, malnutrition, lower growth, and poor health condition in piglets are all symptoms of weaning stress. In an intensified farming industry, animal producers employ a variety of ways to alleviate weaning distress in piglets. ZnO and probiotics have been used in animal farming for a long period of time, providing good results (Lan et al. 2017; Upadhaya et al. 2018).

Effect of high- versus low-dose ZnO diet

In the present research, pigs provided therapeutic levels of ZnO had significantly greater BW, ADG, and G:F than those provided 300 ppm ZnO. Weaning pigs fed therapeutic doses of ZnO increased ADG, ADFI, and G:F (Hill et al. 2001; Sales 2013). Moreover, pigs fed ZnO as a positive control (ZH) diet increased ADG and BW compared with ZnO as an NC diet (ZL) (Shi et al. 2019). Supplementing with 3000 ppm standard ZnO and 200 ppm customized ZnO enhanced BW (Cho et al. 2015), and 2500 mg/kg of Zn from ZnO increased ADG (Hollis et al. 2005), but had no effect on ADFI or G:F. In agreement with our study, the addition of protected ZnO resulted in significantly greater BW, ADG, and ADFI in weaning pigs (Upadhaya et al. 2018) compared with pigs fed a basal diet with the nutritional content of zinc. In contrast, research found no change in growth performance (BW, ADG, ADFI, and G:F) in weanling pigs fed HZ compared with LZ (Shen et al. 2014). The enhancements in BWG, ADG, and G:F found in this trial may be due to the antibacterial properties of ZnO administration and improved gut health.

Several studies have indicated enhanced growth rate and nutrient digestibility using a therapeutic level (3000–4000 ppm) of ZnO as a substitute for in-feed antibiotics (Cho et al. 2015; Milani et al. 2017). In line with present research, nutrient digestibility was not influenced by supplementing ZH into diet (Shi et al. 2019) compared with ZL. In contrast, adding protected ZnO resulted in higher ATTD of DM and E when compared with the NC diet in piglets (Upadhaya et al. 2018). As the concentration of ZnO reduces, the nutritional

digestibility also reduces quantitatively, where the ATTD of DM and N were found to be identical (Wang et al. 2021). Fecal noxious gas emissions such as NH_3 and H_2S have become one of the leading air pollutants in recent intensive pig production. There are few reports on the effects of ZnO feeding on noxious gas emissions in weaned piglets. In line with our finding, incorporating a probiotic complex into a diet with LZ and HZ did not affect fecal gas emissions in weaned piglets (Shi et al. 2019; Wang et al. 2021). Our findings found no differences in nutrient consumption; therefore, it was reasonable to presume that there would be no differences in noxious gas emissions as well, consistent with Hoque et al. (2021).

Escherichia coli concentrations in the feces of weaned pigs fed diets containing HZ were found to be significantly lower, and *Lactobacillus* was higher (Kaevska et al. 2016; Upadhaya et al. 2018), coherent with current research. In contradiction to this study, consuming ZnO at a level of 2500 mg/kg enhances the level of *E. coli* (Bednorz et al. 2013) in the intestine. With high dosages of ZnO, Hojberg et al. (2005) found that *Lactobacillus* was reduced, but there was no impact on *E. coli* (Broom et al. 2006). Carlson et al. (2008) and Shen et al. (2014) revealed that a greater level of ZnO lowered the diarrhea incidence of weaning pigs. The fecal score, which could indicate diarrhea status, was lower in piglets fed the customized ZnO (Cho et al. 2015) and ZnO nanoparticles (Milani et al. 2017) compared with control diet. The fecal count, which can indicate diarrhea severity, was similar across all treatments, confirming no diarrhea (Shen et al. 2014; Upadhaya et al. 2018), coherent with current research. Higher ZnO doses were thought to minimize the incidence of diarrhea by reducing harmful bacteria counts (Sales 2013). These findings indicated that the impact of ZnO in decreasing diarrhea could be unrelated to the direct inhibition of pathogenic *E. coli*, but rather is linked to an enhancement in gut microbiome, which promotes competition among enterobacteria (Amezcuza et al. 2008; Wang et al. 2021). In the current investigation, diarrhea was not observed in pigs given ZnO diet, which could be related to the decreased *E. coli* count and pigs being raised in a hygienic method.

Effect of probiotics

Probiotics have a favorable influence on the host's health and improve the functioning of the intestinal tract. Decades of research has indicated that the use of probiotics in farm animals is beneficial as it improves feed efficiency, weight gain, and immune response (Al-Shawi et al. 2020). In this study, probiotic inclusion showed a tendency to decrease G:F. Previous research from our laboratory revealed that includ-

Table 8. The effect of dietary supplementation of probiotic complex additive on fecal score in weaning pigs.

Items	High ZnO (3000 ppm)		Low ZnO (300 ppm)		SEM	P value		
	Probiotic– (HZ)	Probiotic+ (HZP)	Probiotic– (LZ)	Probiotic+ (LZP)		Probiotic effect	ZnO effect	ZnO × probiotic
Fecal score								
Week 1	3.61	3.59	3.66	3.61	0.06	0.543	0.515	0.760
Week 2	3.45	3.52	3.54	3.55	0.07	0.524	0.380	0.722
Week 3	3.43	3.39	3.45	3.46	0.11	0.922	0.704	0.819
Week 4	3.27	3.32	3.30	3.32	0.07	0.575	0.785	0.813
Week 5	3.29	3.29	3.34	3.29	0.07	0.765	0.743	0.765
Week 6	3.21	3.21	3.25	3.27	0.07	0.907	0.542	0.907

Abbreviations: TRT1, positive control: basal diet added with 3000 ppm ZnO (HZ); TRT2, HZ + 0.1% probiotic complex (HZP); TRT3, negative control: basal diet added with 300 ppm ZnO (LZ); TRT4, LZ + 0.1% probiotic complex (LZP); SEM, standard error of means.

ing multistrain probiotics (0.1%, *B. licheniformis* and *B. subtilis*) in the diets had a positive impact on ADG, ADFI, and G:F all through the experiment (Lan et al. 2016; Zhang et al. 2020). Indeed, adding a probiotic complex into diet increased growth performance in weaned pigs and broilers in our prior study (Balamuralikrishnan et al. 2017; Nguyen et al. 2019). Moreover, the diet inclusion of ZnO with probiotic complex decreased ADG, but had no effect on ADFI and G:F (Wang et al. 2021). The administration of *Bacillus*-based probiotics in the diet of finishing pigs had no effect on ADFI and G:F (Chen et al. 2006), ADG, or feed efficiency (Munoz et al. 2007), which is slightly in line with our research. Some recent studies found that feeding *B. subtilis* and *B. licheniformis* to basal diet improved BW and FCR for grower-finisher pigs (Alexopoulos et al. 2004), and enhanced ADG and ADFI for fattening goats (Lu et al. 2021). Probiotics would support stabilizing the microbiota by encouraging the beneficial bacteria, increasing gut health, decreasing the occurrence of diarrhea, and boosting growth performance (Hu et al. 2014).

Weaning pigs' feeds enriched with multimicrobe probiotic-containing products such as *B. subtilis* enhanced ATTD of DM, N, and E, according to Nguyen et al. (2019), which correlates with Lee et al. (2014). Multistrain probiotic (0.1%) administration can improve ATTD of DM, N, and E (Upadhaya et al. 2015; Lan et al. 2016), which is supported by Hu et al. (2021). However, diverse effects were accompanied by Chen et al. (2006), who showed that addition of *Bacillus*-based probiotics into diet had no effect on DM and N digestibility, in agreement with current study. In contrary, probiotics (*B. subtilis* and *C. butyricum* endospore) administration in growing-finishing pig diet enhanced ATTD of N and E, whereas interactive effects of probiotics were detected on the digestibility of N at week 10 and E at week 5 (Meng et al. 2010). These variable probiotic actions were linked to different types of probiotics and the level of feeding and management (Yu et al. 2020). Nutritive probiotic addition is assumed to have optimistic influence on the gut microbiome ecosystem, causing a shift in the intestinal flora that leads to increased nitrogen consumption and, in turn, reduced noxious gas emissions from the feces (Jeong et al. 2015). In comparison to control diet, Chen et al. (2006) and Hu et al. (2021) discovered that dietary addition with a *Bacillus*-based complex (0.2%) lowered NH₃ and amounts of total mer-

captan. As observed, probiotic treatment with *B. subtilis* and *B. licheniformis* considerably reduces slurry noxious NH₃ emission, but had no effect on H₂S and mercaptan discharge in growing pigs (Wang et al. 2009). In the present investigation, dietary inclusion of probiotic had no effect on noxious gas emission. If there is no change in digestibility, there will be no change in noxious gas discharges.

While the beneficial bacterium *Lactobacillus* is responsible for maintaining intestinal health and can help to prevent diarrhea, *E. coli* is one of the most common intestinal pathogens, and some strains are serious enough to cause diarrhea (Hu et al. 2014). In contrast with current research, *E. coli* increased, but *Lactobacillus* decreased in the intestinal flora of weaned piglets (Konstantinov et al. 2006). Lan et al. (2016) and Upadhaya et al. (2017) found that probiotic treatment resulted in higher intestinal *Lactobacillus* counts and decreased *E. coli* counts in weanling piglets, which is in agreement with our findings. Probiotics benefit the host animal by maintaining intestinal barrier and producing gut microecological environments that suppress infectious organisms while increasing helpful bacteria (Chen et al. 2013). In weaned piglets, Giang et al. (2012) discovered that *Bacillus*-based microbiota lowered diarrhea incidence and reduced fecal score. Furthermore, Alexopoulos et al. (2004) discovered that feeding probiotic spores significantly reduced occurrence of diarrhea in postweaning period. In this study, addition of dietary probiotic to the diet increased fecal *Lactobacillus* and reduced *E. coli* counts in piglets, resulting in no diarrhea incidence. Probiotics in the feed had a significant impact on intestinal bacterial populations, thus decreasing diarrhea scores (Dowarah et al. 2017).

Interactive effects between ZnO and probiotics

In all of the analyzed parameters (growth performance, nutrient utilization, fecal microbiome, noxious gas production, and fecal index), there were no interaction effects between ZnO and probiotics.

Conclusion

In conclusion, high-dose ZnO proved its beneficial effect on growth performance and gut condition of weaning pigs.

However, there was no individual or synergistic effect between probiotic addition and ZnO on weaning pigs' performance and nutrient digestibility. This study suggests to apply a higher dose or different combinations of probiotics to select a suitable lower dose of ZnO to facilitate weaning pigs' overall performance.

Acknowledgements

This work was supported by Korea Institute of Planning and Evaluation for Technology in Food, Agriculture and Forestry (IPET) through Useful Agricultural Life Resources Industry Technology Development Program, funded by Ministry of Agriculture, Food and Rural Affairs (MAFRA) (321094-2) and The Department of Animal Resource & Science was supported through the Research-Focused Department Promotion & Interdisciplinary Convergence Research Projects as a part of the University Innovation Support Program for Dankook University in 2022.

Article information

History dates

Received: 18 June 2022

Accepted: 22 August 2022

Accepted manuscript online: 22 October 2022

Version of record online: 18 November 2022

Copyright

© 2022 The Author(s). Permission for reuse (free in most cases) can be obtained from [copyright.com](https://creativecommons.org/licenses/by/4.0/).

Data availability

The data that aid this investigation will be shared upon reasonable request to the corresponding author.

Author information

Author ORCIDs

Sarbani Biswas <https://orcid.org/0000-0001-9762-807X>

In Ho Kim <https://orcid.org/0000-0001-6652-2504>

Author notes

In Ho Kim served as an Associate Editor at the time of manuscript review and acceptance; peer review and editorial decisions regarding this manuscript were handled by Lee-Anne Huber and Filippo Miglior.

Author contributions

Conceptualization, data curation, formal analysis, investigation, methodology, validation, writing—original draft: SB
Software, Writing—review and editing: MHK
Investigation, software, validation, visualization: JHP
Software, writing—review and editing: YK
Conceptualization, funding acquisition, investigation, methodology, project administration, resources, validation, writing—review and editing: IHK

Competing interests

There are no recognized conflicting interests for the authors.

References

- Alexopoulos, C., Georgoulakis, I.E., Tzivara, A., Kyriakis, C.S., Govaris, A., and Kyriakis, S.C. 2004. Field evaluation of the effect of a probiotic-containing *Bacillus licheniformis* and *Bacillus subtilis* spores on the health status, performance, and carcass quality of grower and finisher pigs. *J. Vet. Med. Ser. A*, **51**(6): 306–312. doi:[10.1111/j.1439-0442.2004.00637.x](https://doi.org/10.1111/j.1439-0442.2004.00637.x).
- Al-Shawi, S.G., Dang, D.S., Yousif, A.Y., Al-Younis, Z.K., Najm, T.A., and Matarnah, S.K. 2020. The potential use of probiotics to improve animal health, efficiency, and meat quality: a review. *Agriculture*, **10**(10): 452. doi:[10.3390/agriculture10100452](https://doi.org/10.3390/agriculture10100452).
- Amezcu, R., Friendship, R.M., and Dewey, C.E. 2008. An investigation of the presence of *Escherichia coli* O149:K91:F4 on pig farms in southern Ontario and the use of antimicrobials and risk factors associated with the presence of this serogroup. *Can. Vet. J.* **49**(1): 39. PMID: [18320976](https://pubmed.ncbi.nlm.nih.gov/18320976/).
- AOAC. 2010. Official methods of analysis. 17th ed. Association of Official Analytical Chemists, Washington, DC.
- Balamuralikrishnan, B., Lee, S.I., and Kim, I.H. 2017. Dietary inclusion of different multi-strain complex probiotics: effects on performance in broilers. *Br. Poult. Sci.* **58**(1): 83–86. doi:[10.1080/00071668.2016.1257112](https://doi.org/10.1080/00071668.2016.1257112). PMID: [27918205](https://pubmed.ncbi.nlm.nih.gov/27918205/).
- Bednorz, C., Oelgeschlager, K., Kinnemann, B., Hartmann, S., Neumann, K., Pieper, R., et al. 2013. The broader context of antibiotic resistance: zinc feed supplementation of piglets increases the proportion of multi-resistant *Escherichia coli* in vivo. *Int. J. Med. Microbiol.* **303**: 396–403. doi:[10.1016/j.ijmm.2013.06.004](https://doi.org/10.1016/j.ijmm.2013.06.004). PMID: [23856339](https://pubmed.ncbi.nlm.nih.gov/23856339/).
- Bedford, A., Huynh, E., Fu, M., Zhu, C., Wey, D., Lange, C.D. and Li, J. 2014. Growth performance of early-weaned pigs is enhanced by feeding epidermal growth factor-expressing *Lactococcus lactis* fermentation product. *J. Biotechnol.* **173**: 47–52. doi:[10.1016/j.jbiotec.2014.01.012](https://doi.org/10.1016/j.jbiotec.2014.01.012).
- Boirivant, M., and Strober, W. 2007. The mechanism of action of probiotics. *Curr. Opin. Gastroenterol.* **23**(6): 679–692. doi:[10.1097/MOG.0b013e3282f0cfc](https://doi.org/10.1097/MOG.0b013e3282f0cfc).
- Broom, L.J., Miller, H.M., Kerr, K.G., and Knapp, J.S. 2006. Effects of zinc oxide and *Enterococcus faecium* SF68 dietary supplementation on the performance, intestinal microbiota and immune status of weaned piglets. *Res. Vet. Sci.* **80**(1): 45–54. doi:[10.1016/j.rvsc.2005.04.004](https://doi.org/10.1016/j.rvsc.2005.04.004). PMID: [15946717](https://pubmed.ncbi.nlm.nih.gov/15946717/).
- Carlson, D., Sehested, J., Feng, Z., and Poulsen, H.D. 2008. Serosal zinc attenuate serotonin and vasoactive intestinal peptide induced secretion in piglet small intestinal epithelium in vitro. *Comp. Biochem. Physiol. A: Mol. Integr. Physiol.* **149**(1): 51–58. doi:[10.1016/j.cbpa.2007.10.005](https://doi.org/10.1016/j.cbpa.2007.10.005).
- Chaucheyras-Durand, F., and Durand, H. 2010. Probiotics in animal nutrition and health. *Benef. Microbes*, **1**(1): 3–9. doi:[10.3920/BM2008.1002](https://doi.org/10.3920/BM2008.1002).
- Chen, W., Wang, J.P., Yan, L., and Huang, Y.Q. 2013. Evaluation of probiotics in diets with different nutrient densities on growth performance, blood characteristics, relative organ weight and breast meat characteristics in broilers. *Br. Poult. Sci.* **54**: 635–641. doi:[10.1080/00071668.2013.825369](https://doi.org/10.1080/00071668.2013.825369). PMID: [24112072](https://pubmed.ncbi.nlm.nih.gov/24112072/).
- Chen, Y.J., Min, B.J., Cho, J.H., Kwon, O.S., Son, K.S., Kim, H.J., and Kim, I.H. 2006. Effects of dietary *Bacillus*-based probiotic on growth performance, nutrients digestibility, blood characteristics and fecal noxious gas content in finishing pigs. *Asian-Australas. J. Anim. Sci.* **19**(4): 587–592. doi:[10.5713/ajas.2006.587](https://doi.org/10.5713/ajas.2006.587).
- Cho, J.H., Upadhaya, S.D., and Kim, I.H. 2015. Effects of dietary supplementation of modified zinc oxide on growth performance, nutrient digestibility, blood profiles, fecal microbes shedding and fecal score in weanling pigs. *Anim. Sci. J.* **86**(6): 617–623. doi:[10.1111/asj.12329](https://doi.org/10.1111/asj.12329). PMID: [25473794](https://pubmed.ncbi.nlm.nih.gov/25473794/).
- Dowarah, R., Verma, A.K., Agarwal, N., Patel, B.H.M., and Singh, P. 2017. Effect of swine based probiotic on performance, diarrhoea, intestinal microbiota and gut health of grower–finisher crossbred pigs. *Livest. Sci.* **195**: 74–79. doi:[10.1016/j.livsci.2016.11.006](https://doi.org/10.1016/j.livsci.2016.11.006).
- Dumitru, M., Hăbeanu, M., Sorescu, I., and Tabuc, C. 2021. Effects of *Bacillus* spp. as a supplemental probiotic in diets for weaned piglets. *S. Afr. J. Anim. Sci.* **51**(5). doi:[10.4314/sajas.v51i5.4](https://doi.org/10.4314/sajas.v51i5.4).

- European Communities. 2003. Commission Regulation (EC) No. 1334/2003 of 25 July 2003. Amending the conditions for authorization of a number of additives in feeding stuffs belonging to the group of trace elements. Off. J. Eur. Union 6.
- FAO/WHO. 2002. Guidelines for the evaluation of probiotics in food. Report of a Joint FAO/WHO Working Group on Drafting Guidelines for the Evaluation of Probiotics in Food. Available from http://www.who.int/foodsafety/fs_management/en/probiotic_guidelines.pdf [accessed 9 September 2020].
- Ferket, P.R., Van Heugten, E., Van Kempen, T.A., and Angel, R. 2002. Nutritional strategies to reduce environmental emissions from non-ruminants. *J. Anim. Sci.* **80**: 168–182. doi:10.2527/animalsci2002.80E-Suppl_2E168x.
- Giang, H.H., Viet, T.Q., Ogle, B., and Lindberg, J.E. 2012. Growth performance, digestibility, gut environment and health status in weaned piglets fed a diet supplemented with a complex of lactic acid bacteria alone or in combination with *Bacillus subtilis* and *Saccharomyces boulardii*. *Livest. Sci.* **143**(2–3): 132–141. doi:10.1016/j.livsci.2011.09.003.
- Hill, G.M., Mahan, D.C., Carter, S.D., Cromwell, G.L., Ewan, R.C., Harrold, R.L., et al. 2001. Effect of pharmacological concentrations of zinc oxide with or without the inclusion of an antibacterial agent on nursery pig performance. *J. Anim. Sci.* **79**(4): 934–941. doi:10.2527/2001.794934x. PMID: 11325200.
- Højberg, O., Canibe, N., Poulsen, H.D., Hedemann, M.S., and Jensen, B.B. 2005. Influence of dietary zinc oxide and copper sulfate on the gastrointestinal ecosystem in newly weaned piglets. *Appl. Environ. Microbiol.* **71**(5): 2267–2277. doi:10.1128/AEM.71.5.2267-2277.2005. PMID: 15870311.
- Hollis, G.R., Carter, S.D., Cline, T.R., Crenshaw, T.D., Cromwell, G.L., Hill, G.M., and Veum, T.L. 2005. Effects of replacing pharmacological levels of dietary zinc oxide with lower dietary levels of various organic zinc sources for weanling pigs. *J. Anim. Sci.* **83**(9): 2123–2129. doi:10.2527/2005.8392123x. PMID: 16100067.
- Hoque, M.R., Jung, H.I., and Kim, I.H. 2021. Effect of yeast culture (*Saccharomyces cerevisiae*) supplementation on growth performance, excreta microbes, noxious gas, nutrient utilization, and meat quality of broiler chicken. *Poult. Sci. J.* **58**: 216–221. doi:10.2141/jpsa.0190144.
- Hu, J., Young, Y.H., and Kim, I.H. 2021. Effects of two *Bacillus* strains probiotic supplement on reproduction performance, nutrient digestibility, blood profile, fecal score, excreta odor contents and fecal microflora in lactation sows, and growth performance in sucking piglets. *Livest. Sci.* **244**: 104293. doi:10.1016/j.livsci.2020.104293.
- Hu, C.H., Gu, L.Y., Luan, Z.S., Song, J., and Zhu, K. 2012. Effects of montmorillonite–zinc oxide hybrid on performance, diarrhea, intestinal permeability and morphology of weanling pigs. *Anim. Feed Sci. Technol.* **177**(1–2): 108–115. doi:10.1016/j.anifeeds.2012.07.028.
- Hu, Y., Dun, Y., Li, S., Zhao, S., Peng, N., and Liang, Y. 2014. Effects of *Bacillus subtilis* KN-42 on growth performance, diarrhea and fecal bacterial flora of weaned piglets. *Asian-Australas. J. Anim. Sci.* **27**(8): 1131–1140. doi:10.5713/ajas.2013.13737. PMID: 25083107.
- Jensen, J., Larsen, M.M., and Bak, J. 2016. National monitoring study in Denmark finds increased and critical levels of copper and zinc in arable soils fertilized with pig slurry. *Environ. Pollut.* **214**: 334–340. doi:10.1016/j.envpol.2016.03.034. PMID: 27107257.
- Jeong, J., Kim, J., Lee, S., and Kim, I.H. 2015. Evaluation of *Bacillus subtilis* and *Lactobacillus acidophilus* probiotic supplementation on reproductive performance and noxious gas emission in sows. *Ann. Anim. Sci.* **15**(3): 699–710. doi:10.1515/aoas-2015-0018.
- Kaevska, M., Lorencova, A., Videnska, P., Sedlar, K., Provaznik, I., and Trckova, M. 2016. Effect of sodium humate and zinc oxide used in prophylaxis of post-weaning diarrhoea on faecal microbiota composition in weaned piglets. *Vet. Med.* **61**(6): 328–336. doi:10.17221/54/2016-VETMED.
- Konstantinov, S.R., Awati, A.A., Williams, B.A., Miller, B.G., Jones, P., Stokes, C.R., and de Vos, W.M. 2006. Post-natal development of the porcine microbiota composition and activities. *Environ. Microbiol.* **8**(7): 1191–1199. doi:10.1111/j.1462-2920.2006.01009.x. PMID: 16817927.
- Lalles, J.P. 2008. Nutrition and gut health of the young pig around weaning: what news? *Arch. Zootech.* **11**: 5–15.
- Lalles, J., Bosi, P., Smidt, H., and Stokes, C.R. 2007. Weaning—a challenge to gut physiologists. *Livest. Sci.* **108**(1–3): 82–93. doi:10.1016/j.livsci.2007.01.091.
- Lan, R.X., Lee, S.I., and Kim, I.H. 2016. Effects of multistrain probiotics on growth performance, nutrient digestibility, blood profiles, faecal microbial shedding, faecal score and noxious gas emission in weaning pigs. *J. Anim. Physiol. Anim. Nutr.* **100**(6): 1130–1138. doi:10.1111/jpn.12501.
- Lan, R., Tran, H., and Kim, I. 2017. Effects of probiotic supplementation in different nutrient density diets on growth performance, nutrient digestibility, blood profiles, fecal microflora and noxious gas emission in weaning pig. *J. Sci. Food Agric.* **97**(4): 1335–1341. doi:10.1002/jsfa.7871. PMID: 27342952.
- Lee, S.H., Ingale, S.L., Kim, J.S., Kim, K.H., Lokhande, A. Kim, E.K., et al. 2014. Effects of dietary supplementation with *Bacillus subtilis* LS 1–2 fermentation biomass on growth performance, nutrient digestibility, cecal microbiota and intestinal morphology of weanling pig. *Anim. Feed Sci. Technol.* **188**: 102–110. doi:10.1016/j.anifeeds.2013.12.001.
- Li, M.Z., Huang, J.T., Tsai, Y.H., Mao, S.Y., Fu, C.M., and Lien, T.F. 2016. Nanosize of zinc oxide and the effects on zinc digestibility, growth performances, immune response and serum parameters of weanling piglets. *Anim. Sci. J.* **87**(11): 1379–1385. doi:10.1111/asj.12579. PMID: 26805551.
- Liao, S., and Nyachoti, M. 2017. Using probiotics to improve swine gut health and nutrient utilization. *Anim. Nutr.* **3**: 331–343. doi:10.1016/j.aninu.2017.06.007. PMID: 29767089.
- Lu, J., Chen, Z., Gao, Q., Li, P., Wang, J., Cai, Y., et al. 2021. Combined supplementation of probiotics and enzymes improves performance and regulates rumen microbiota in fattening goats. *Research Square*, 1–26. doi:10.21203/rs.3.rs-1043297/v1
- Meng, Q.W., Yan, L., Ao, X., Zhou, T.X., Wang, J.P., Lee, J.H., and Kim, I.H. 2010. Influence of probiotics in different energy and nutrient density diets on growth performance, nutrient digestibility, meat quality, and blood characteristics in growing-finishing pigs. *J. Anim. Sci.* **88**(10): 3320–3326. doi:10.2527/jas.2009-2308. PMID: 20562363.
- Milani, N.C., Sbardella, M., Ikeda, N.Y., Arno, A., Mascarenhas, B.C., and Miyada, V.S. 2017. Dietary zinc oxide nanoparticles as growth promoter for weanling pigs. *Anim. Feed Sci. Technol.* **227**: 13–23. doi:10.1016/j.anifeeds.2017.03.001.
- Munoz, D., Lanz, A., Lucero, P., Soria, F., Renteria, F., Cuaron, J.A., et al. 2007. Strategies for enhancing microbiological gut's barrier: BMD y BioPlus 2B. *J. Anim. Sci.* **85**: 150.
- National Research Council. 2012. Nutrient requirement of swine. 11th rev. ed. National Academies Press, Washington, DC.
- Nguyen, D.H., Nyachoti, C.M., and Kim, I.H. 2019. Evaluation of effect of probiotics mixture supplementation on growth performance, nutrient digestibility, fecal bacterial enumeration, and noxious gas emission in weaning pigs. *Ital. J. Anim. Sci.* **18**(1): 466–473. doi:10.1080/1828051X.2018.1537726.
- Pluske, J.R. 2013. Feed- and feed additives-related aspects of gut health and development in weanling pigs. *J. Anim. Sci. Biotechnol.* **4**: 1–6. doi:10.1186/2049-1891-4-1. PMID: 23289727.
- Sales, J. 2013. Effects of pharmacological concentrations of dietary zinc oxide on growth of post-weaning pigs: a meta-analysis. *Biol. Trace Elem. Res.* **152**(3): 343–349. doi:10.1007/s12011-013-9638-3. PMID: 23463368.
- Shen, J., Chen, Y., Wang, Z., Zhou, A., He, M., Mao, L., et al. 2014. Coated zinc oxide improves intestinal immunity function and regulates microbiota composition in weaned piglets. *Br. J. Nutr.* **111**(12): 2123–2134. doi:10.1017/S0007114514000300. PMID: 24606984.
- Shi, H., Upadhaya, S.D., and Kim, I.H. 2019. Comparative effects of low ZnO dose with or without probiotics relative to high ZnO dose on the performance, nutrient digestibility, blood metabolites, and noxious gas emissions in weaned piglets. *Can. J. Anim. Sci.* **99**: 881–889. doi:10.1139/cjas-2019-0059.
- Slifierz, M.J., Friendship, R., and Weese, J.S. 2015. Zinc oxide therapy increases prevalence and persistence of methicillin-resistant *Staphylococcus aureus* in pigs: a randomized controlled trial. *Zoonoses Public Health.* **62**: 301–308. doi:10.1111/zph.12150. PMID: 25209545.
- Trckova, M., Lorencova, A., Hazova, K., and Sramkova Zajacova, Z., 2015. Prophylaxis of post-weaning diarrhoea in piglets by zinc

- oxide and sodium humate. *Vet. Med.* **60**: 351–360. doi:[10.17221/8182-VETMED](https://doi.org/10.17221/8182-VETMED).
- Upadhaya, S.D., Kim, S.C., Valientes, R.A., and Kim, I.H. 2015. The effect of *Bacillus*-based feed additive on growth performance, nutrient digestibility, fecal gas emission, and pen cleanup characteristics of growing–finishing pigs. *Asian-Australas. J. Anim. Sci.* **28**(7): 999–1005. doi:[10.5713/ajas.15.0066](https://doi.org/10.5713/ajas.15.0066). PMID: [26104405](https://pubmed.ncbi.nlm.nih.gov/26104405/).
- Upadhaya, S.D., Shanmugam, S.K., Kang, D.K., and Kim, I.H. 2017. Preliminary assessment on potentials of probiotic *B. subtilis* RX7 and *B. methylotrophicus* C14 strains as an immune modulator in *Salmonella*-challenged weaned pigs. *Trop. Anim. Health Prod.* **49**(5): 1065–1070. doi:[10.1007/s11250-017-1278-8](https://doi.org/10.1007/s11250-017-1278-8). PMID: [28326485](https://pubmed.ncbi.nlm.nih.gov/28326485/).
- Upadhaya, S.D., Kim, Y.M., Lee, K.Y., and Kim, I.H. 2018. Use of protected zinc oxide in lower doses in weaned pigs in substitution for the conventional high dose zinc oxide. *Anim. Feed Sci. Technol.* **240**: 1–10. doi:[10.1016/j.anifeedsci.2018.03.012](https://doi.org/10.1016/j.anifeedsci.2018.03.012).
- Wang, H., Kim, K.P., and Kim, I.H. 2021. Evaluation of the combined effects of different dose levels of zinc oxide with probiotics complex supplementation on the growth performance, nutrient digestibility, faecal microbiota, noxious gas emissions and faecal score of weaning pigs. *J. Anim. Physiol. Anim. Nutr.* **105**: 286–293. doi:[10.1111/jpn.13493](https://doi.org/10.1111/jpn.13493).
- Wang, Y., Cho, J.H., Chen, Y.J., Yoo, J.S., Huang, Y., Kim, H.J., and Kim, I.H. 2009. The effect of probiotic BioPlus 2B® on growth performance, dry matter and nitrogen digestibility and slurry noxious gas emission in growing pigs. *Livest. Sci.* **120**(1–2): 35–42. doi:[10.1016/j.livsci.2008.04.018](https://doi.org/10.1016/j.livsci.2008.04.018).
- Yu, H.F., Wang, A.N., Li, X.J., and Qiao, S.Y. 2008. Effect of viable *Lactobacillus fermentum* on the growth performance, nutrient digestibility and immunity of weaned pigs. *J. Anim. Feed Sci.* **17**(1): 61–69. doi:[10.22358/jafs/66470/2008](https://doi.org/10.22358/jafs/66470/2008).
- Yu, W., Hao, X., Zhiyue, W., Haiming, Y., and Lei, X. 2020. Evaluation of the effect of *Bacillus subtilis* and *Pediococcus acidilactici* mix on serum biochemistry, growth promotion of body and visceral organs in Lohmann brown chicks. *Braz. J. Poult. Sci.* **22**: 1274. doi:[10.1590/1806-9061-2020-1274](https://doi.org/10.1590/1806-9061-2020-1274).
- Zhang, S., Yoo, D.H., Ao, X., and Kim, I.H. 2020. Effects of dietary probiotic, liquid feed and nutritional concentration on the growth performance, nutrient digestibility and fecal score of weaning piglets. *Asian-Australas. J. Anim. Sci.* **33**(10): 1617–1623. doi:[10.5713/ajas.19.0473](https://doi.org/10.5713/ajas.19.0473). PMID: [32054199](https://pubmed.ncbi.nlm.nih.gov/32054199/).
- Zhao, P.Y., and Kim, I.H. 2015. Effect of direct-fed microbial on growth performance, nutrient digestibility, fecal noxious gas emission, fecal microbial flora and diarrhea score in weanling pigs. *Anim. Feed Sci. Technol.* **200**: 86–92. doi:[10.1016/j.anifeedsci.2014.12.010](https://doi.org/10.1016/j.anifeedsci.2014.12.010).
- Zhao, C.Y., Tan, S.X., Xiao, X.Y., Qiu, X.S., Pan, J.Q., and Tang, Z.X. 2014. Effects of dietary zinc oxide nanoparticles on growth performance and antioxidative status in broilers. *Biol. Trace Elem. Res.* **160**: 361–367. doi:[10.1007/s12011-014-0052-2](https://doi.org/10.1007/s12011-014-0052-2). PMID: [24973873](https://pubmed.ncbi.nlm.nih.gov/24973873/).
- Zhu, C., Lv, H., Chen, Z., Wang, L., Wu, X. Zhang, W., et al. 2017. Dietary zinc oxide modulates antioxidant capacity, small intestine development, and jejunal gene expression in weaned piglets. *Biol. Trace Elem. Res.* **175**: 331–338. doi:[10.1007/s12011-016-0767-3](https://doi.org/10.1007/s12011-016-0767-3). PMID: [27339255](https://pubmed.ncbi.nlm.nih.gov/27339255/).