



## **Disentangling the taxonomy of the *Cleome africana* species complex (Cleomaceae) in North Africa**

Authors: Lemmel, Claude, and Chatelain, Cyrille

Source: *Candollea*, 79(2) : 277-281

Published By: The Conservatory and Botanical Garden of the City of Geneva (CJBG)

URL: <https://doi.org/10.15553/c2024v792a4>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# Disentangling the taxonomy of the *Cleome africana* species complex (Cleomaceae) in North Africa

Claude Lemmel & Cyrille Chatelain

## Abstract

LEMMELE, C. & C. CHATELAIN (2024). Disentangling the taxonomy of the *Cleome africana* species complex (Cleomaceae) in North Africa. *Candollea* 79: 277–281. In English, English and French abstracts. DOI: <http://dx.doi.org/10.15553/c2024v792a4>

We propose to revisit the taxonomy of *Cleome africana* Botsch. (Cleomaceae) in North Africa. The taxonomic circumscription of this species has been problematic over time leading to numerous confusions and misidentifications in most floras from this region. In North Africa, we propose to accept a single species, i.e., *C. amblyocarpa* Barratte & Murb. with two subspecies. *Cleome africana* Botsch., described as a replacement name for *Siliquaria glandulosa* Forssk., is accepted at subspecific rank under *C. amblyocarpa* as *C. amblyocarpa* var. *glandulosa* (Forssk.) Botsch. We consider that *C. arabica* L. only occurs in the Middle East. An identification key is provided for the *C. africana* complex.

## Résumé

LEMMELE, C. & C. CHATELAIN (2024). Éclaircir la taxonomie du genre *Cleome* (Cleomaceae) en Afrique du Nord. *Candollea* 79: 277–281. En anglais, résumés anglais et français. DOI: <http://dx.doi.org/10.15553/c2024v792a4>

Nous proposons de revisiter la taxonomie de *Cleome africana* Botsch. (Cleomaceae) en Afrique du Nord. La circonscription taxonomique de cette espèce a été problématique au fil du temps, ce qui a conduit à de nombreuses confusions et erreurs d'identification dans la plupart des flores de cette région. En Afrique du Nord, nous proposons d'accepter une seule espèce, à savoir *C. amblyocarpa* Barratte & Murb. avec deux sous-espèces. *Cleome africana* Botsch., désigné comme nom de remplacement pour *Siliquaria glandulosa* Forssk., est accepté au rang de sous-espèce de *C. amblyocarpa* en tant que *C. amblyocarpa* var. *glandulosa* (Forssk.) Botsch. Nous considérons que *C. arabica* L. n'est présent qu'au Moyen-Orient. Une clé d'identification est fournie pour le complexe *C. africana*.

## Keywords

CLEOMACEAE – *Cleome africana* – *Cleome arabica* – Africa – synonymy – taxonomy

---

Adresses des auteurs:

CL: 52150 Boudnib – Errachidia, Morocco.

CC: Conservatoire et Jardin botaniques de Genève, ch. Impératrice 1, 1292 Chambésy, Switzerland.

E-Mail: [cyrille.chatelain@geneve.ch](mailto:cyrille.chatelain@geneve.ch)

Submitted on April 29, 2024 – Accepted on August 15, 2024 – First published online on October 15, 2024

ISSN: 0373-2967 (print), 2235-3658 (online) – Published by the Conservatoire et Jardin botaniques de Genève

Open access article under Creative Commons Attribution Licence (CC BY 4.0)

## Introduction

The name *Cleome arabica*, originally described by LINNAEUS (1755) on the basis of a Hasselquist specimen, was subsequently ascribed to two different taxonomic entities in several publications by LINNAEUS himself (1755, 1759, 1763, 1767) and LINNAEUS f. (1767), which rendered numerous misinterpretations by posterior botanists. The original material used by LINNAEUS (1755: 20) to describe *C. arabica* in *Centuria I. Plantarum* was collected by Fredrik Hasselquist (1722–1752) in Egypt in 1750–1752. This specimen corresponds to a taxon with unifoliolate leaves. The same diagnosis was also included in the *Amoenitates academicae* (1759: 281). In the second edition of *Species plantarum*, LINNAEUS (1763: 937–940) listed 15 species of *Cleome*, including *C. arabica* based on Hasselquist specimen.

The confusion arose when young Linnaeus redescribed *C. arabica* in *Plantarum rariorum horti upsaliensis fasciculus primus* (LINNAEUS f., 1767). He referred to the description of his father based on the specimen of Hasselquist as “foliis simplicibus subrotundo-ovatis” (LINNAEUS f., 1767: 15), but provided a copper engraving based on a cultivated plant in the Upsala Botanical Garden originating from seeds collected by Peter Forsskål (1732–1763) in Egypt in 1760–1762 (LINNAEUS f., 1767: tab. VIII). This engraving depicted a plant with trifoliolate leaves. The same year, LINNAEUS (1767: 448) in the twelfth edition of his *Systema naturae* also referred to the latter engraving when describing his *C. arabica* with unifoliolate leaves.

Carten Niebuhr (1733–1815) posthumously published the work of Forsskål as *Flora Aegyptiaco-Arabica: sive descriptiones plantarum* (FORSSKÅL, 1775) with an additional volume of engravings entitled *Icones rerum naturalium* (FORSSKÅL, 1776). *Siliquaria glandulosa* Forssk. was described in the *Flora Aegyptiaco-Arabica* (FORSSKÅL, 1775: 78) and illustrated it in the *Icones* (FORSSKÅL, 1776: tab. XVI). Furthermore, FORSSKÅL (1776: 6) included the text “*Cleome arabica*. L. S. N. Tom. II. pag. 448. n. 11”. Forsskål’s *Siliquaria glandulosa* clearly represents the species with trifoliolate leaves depicted in LINNAEUS f. (1767: tab. VIII) and cited by his father the same year. Since then, Forsskål’s name was inaccurately included in the synonymy of *Cleome arabica*.

In the beginning of the XX<sup>th</sup> century, MURBECK (1905) described a new species of *Cleome* from Algeria, *C. amblyocarpa* Barratte & Murb., and compared it with *C. arabica* in the sense of the trifoliolate species [hereafter as “auct.”]. Among other characters, *C. amblyocarpa* differs from *C. arabica* auct. by having seeds with shorter trichomes. BOTSCHANTZEV (1964) published *C. africana* Botsch. as a replacement name for *Siliquaria glandulosa* [non *C. glandulosa* Ruiz & Pav. ex DC., 1824] for a broad concept species including *C. arabica* auct. At that time, Botschantzev overlooked *C. amblyocarpa*. In 1968, when he realised about its existence (BOTSCHANTZEV,

1968), accepted it as a distinct taxon and proposed the variety *glandulosa* (Forssk.) Botsch. to keep the concept of his *C. africana*. At the same time, he redefined the “genuine” *C. arabica* L. [in the sense of the unifoliolate species] retrieving the name *C. trinervia* Fresen., which has no priority over Linnaeus’s name.

HEDGE & LAMOND (1970) appeared to be well-aware of all this confusion when publishing the *Cleome* treatment for *Flora iranica*. On one hand, they retained the name *C. amblyocarpa* (with *C. arabica* auct. in the synonymy) for Africa and the Middle East region. On the other hand, restricted the “genuine” *C. arabica* to the Middle East region.

More recently, ROALSON (2021) recognised only one taxon for North Africa, *C. amblyocarpa*, placing *Siliquaria glandulosa* and Botschantzev’s replacement name *C. africana* in the synonymy. This taxonomic criterion is not accepted in the present treatment, where two taxa around *C. amblyocarpa* are recognised. Concerning the “genuine” *C. arabica*, Roalson restricted it to Middle East.

In such a situation, in part originating by the misinterpretation of the original concept of *C. arabica* but also the ambiguous concept behind the replacement name *C. africana*, the treatments proposed around those names in North Africa have been disparate over time (e.g. MAIRE, 1965; ZOHARY, 1966; BOULOS, 1999; FENNANE et al. 1999; POWO, 2024). Facing that, we propose (1) to accept *C. amblyocarpa* var. *amblyocarpa* to circumscribe the taxon with seeds covered by short trichomes; (2) to retain the original concept of *Siliquaria glandulosa* under the name *C. amblyocarpa* var. *glandulosa* (Forssk.) Botsch. as proposed by BOTSCHANTZEV (1968) for the taxon with seeds covered by long trichomes; and (3) to keep the name *C. arabica* for the species originally described by Linnaeus and characterized by having unifoliolate leaves.

## Key to the *Cleome africana* complex in North Africa

1. Leaves simple, trinerved, lanate at maturity ... *C. arabica*
  - 1a. Leaves trifoliolate, linear to elliptic, covered with glandular hairs sticking to sand, but sometimes with glandular hairs only on stem ..... 2
  2. Petals clawed, entirely purple (also when dried) a little longer than wide, with hairs on seeds (< 2/3 diameter of seed), inner part of pod without reticulate veins ..... *C. amblyocarpa* var. *amblyocarpa*
  - 2a. Petals clawed, purple at tip and white at base (also when dried), longer than wide, with very long hairs on seeds (> diameter of seed) ..... *C. amblyocarpa* var. *glandulosa*

## Taxonomy

*Cleome amblyocarpa* Barratte & Murb. in Acta Univ. Lund., ser. 2, 1(4): 25. 1905.

**Lectotypus** (first step designated by BOTSCHANTZEV, 1968; second step designated by ROALSON, 2021: 58): **MOROCCO**: Oasis d'Akka, 1873, *Mardochee* s.n. (P [P05328491]!); isolecto-: LE [LE00013126] image!, P [P05328488, P05328490, P05328493]!).

- *Cleome arabica* subsp. *amblyocarpa* (Barratte & Murb.) Ozenda, Fl. Sahara Sept. Centr.: 247. 1958 [comb. inval.].

*Cleome amblyocarpa* var. *amblyocarpa* (Fig. 1A, C, D).

**Distribution.** – Morocco to Algeria (Fig. 2).

**Additional specimens examined.** – **MOROCCO**: Oasis d'Akka, 1873, *Mardochee* (G [G00628015]); Taroudant, 12 km ESE of Talouine, 17.IV.1986, *Podlech 40979* (G [G00628010]); Drâa-Tafilalet, col du Itiz-n-Tinfift, 14.IV.1986, *Podlech 41037* (G [G00628009]); Drâa-Tafilalet, Ayachi, cirque de Jaffar, 10.VI.1992, *Achbal et al. 6/272* (G [G00433761]).

**ALGERIA**: Biskra, Sahara env. Biskra, 15.IV.1903, *Murbeck s.n.* (LD [LD1221985]); Biskra, col de Sfa, au N de Biskra, 4.V.1906, *Romieux s.n.* (G [G0043377]).

*Cleome amblyocarpa* var. *glandulosa* (Forssk.) Botsch. in Novosti Sist. Vyssh. Rast. 5: 237. 1968 (Fig. 1B–D).

- *Siliquaria glandulosa* Forssk., Fl. Aegypt.-Arab.: 78. 1775 [non *C. glandulosa* Ruiz & Pav. ex DC., 1824]. = *Cleome africana* Botsch. in Novosti Sist. Vyssh. Rast. 1: 130. 1964 [nom. nov., excl. syn.].
- *Cleome siliquaria* R. Br. in Salt, Voy. Abyss., App.: lxv. 1814 [nom. illeg. superfl.].

**Lectotypus**: (designated by ROALSON, 2021: 58): **EGYPT**: “ad Birket el hâdgi prope Kâhîram”, 1762, *Forsskål 640* (C [C10003078] image!; isolecto-: C [C10003079] image!).

- *Cleome arabica* sensu LINNAEUS (1767).
- *Cleome arabica* sensu LINNAEUS f. (1767).
- *Cleome arabica* subsp. *arabica* sensu OZENDA (1958).
- *Cleome arabica* sensu MAIRE (1965).
- *Cleome arabica* sensu ZOHARY (1966).
- *Cleome amblyocarpa* sensu HEDGE & LAMOND (1970).
- *Cleome amblyocarpa* sensu ALI & JAFRI (1976).
- *Cleome africana* sensu FENNANE et al. (1999).
- *Cleome amblyocarpa* sensu BOULOS (1999).
- *Cleome africana* sensu ROALSON (2021).
- *Siliquaria glandulosa* sensu ROALSON (2021).

**Distribution.** – From Morocco to Egypt, Sinai, Palestine to Iraq (Fig. 2).

**Notes.** – Herbarium specimens filed under *Cleome arabica* in most herbaria correspond to *C. amblyocarpa* var. *glandulosa* (Forssk.) Botsch., with a few exceptions.

**Additional specimens examined.** – **ALGERIA**: Biskra, 17.IV.1853, *Balansa 823* (G [G00628020]); M'Sila, Bou Saada, 1.IV.1890, *Battandier & Trabut 512* (G [G00417548]); Tamanrasset, Pic Laperrine, 23.X.2019, *Chatelain et al. CM807* (ENSA, G); Tamanrasset, Tifokraoui, au SW de Idles, Tefedest, 25.X.2019, *Chatelain et al. CM936* (ENSA, G); Tamanrasset, Irharar Kecherouet, 29.IX.2022, *Chatelain et al. CM2220* (G); Tamanrasset, Ouest Tazerouk, 2.IX.2022, *Chatelain et al. CM2340* (G); Naâma, Djebel Aissa, 20.III.2016, *Chatelain CC3790* (G); Biskra, 7.IV.1896, *Chevallier 13* (G [G00417546]); M'Sila, Biskra, IV.1888, *Girod s.n.* (G [G00628014]); Naâma, Ain Sefra, 14.V.1901, *Hochbreutiner 192* (G [G00628017]); Biskra, 27.IX.1851, *Jamin 258* (G [G00628019]); M'Sila, Baniou, près de Bou Saada, 1.III.1869, *Meyer 53* (G [G00417547]).

**EGYPT**: Arabia Petraea, III.1864, *Boissier s.n.* (G-BOIS [G00797584]); Sinâi, 1839, *Bové s.n.* (G-BOIS [G00797513]); Sinâi, désert et l'isthme, Wadi el Arys, 31.III.1891, *Defflers 165* (MPU); sine loco, 1799, *Delile s.n.* (MPU, herb. Delile); “Inter Cairo et Suez, in arvensis deserti”, 8.III.1855, *Kotschy s.n.* (G-BOIS [G00797514]); “Aegyptische Küste zwischen Kasser und Ras-Benass”, 25.II.1865, *Schweinfurth 1213* (G-BOIS [G00797502, G00797504]); “Flora des Ssoturba-Gebirges an der Nubischen Küste”, 5.III.1865, *Schweinfurth 1215* (G-BOIS [G00797583]).

**MOROCCO**: Laâyoune, Smara, 23.III.2015, *Chatelain CC3470* (G); Ouad Eddahab, Gare Ouchfget, 27.III.2017, *Chatelain CC4180b* (G); Boujdour, Zemmour, Agalmim Al Maghdar, 2.IV.2017, *Chatelain CC4384* (G); Drâa-Tafilalet, Djebel Saghro, 3.VI.2019, *Chatelain & Mombrial CM208* (G); Errachidia, N de Tazzarine, 2.VI.2019, *Chatelain & Mombrial CM220* (G); Dakhla, Djebel Negjyr (Nagjir), 29.XII.2017, *Garcin s.n.* (G); Laâyoune, Zemmour, 9.I.2018, *Garcin s.n.* (G [G00394332]).

**TUNISIA**: Kebili, Douz, Kebili, 4.IV.2014, *Chatelain CC2360* (G); Gafsa, Moularès, VI.1944, *Cuénod s.n.* (G [G00433763]); Gafsa, Délégation de Gafsa, 13.IV.1896, *Murbeck s.n.* (G [G00628021]); Sfax, IV.1907, *Pitard 40* (G [G00628013]); Gafsa, 12.IV.1904, *Romieux 22* (G [G00628022]); “in Regno Tunetano”, s.d., *Vahl s.n.* (G).

*Cleome arabica* L., Cent. Pl. 1: 20. 1755.

**Lectotypus** (first step designated by HEDGE & LAMOND, 1970: 27; second step designated by ROALSON, 2021: 58): **EGYPT**: “Arabia”, s.d., *Hasselquist s.n.* (SBT [2.2.3.2] image!; isolecto-: SBT [2.2.3.1] image!, S-LINN [S09-24947] image!).

= *Cleome trinervia* Fresen. in Mus. Senckenberg. 1: 177. 1834. **Lectotypus** (designated by ROALSON 2021: 58): [icon] Fresenius in Mus. Senckenberg. 1: tab. XI.1834.

= *Cleome aschersoniana* Pfund in Flora 57: 413. 1874. **Typus**: “in monte Atakka prope Suez” (not found).

**Distribution.** – Egypt (Sinai), Israel, Palestine, Saudi Arabia to S Iran (Fig. 2).

**Additional specimens examined.** – **JORDAN**: Aqaba, 13.III.1974, *Boulos 137* (G [G0043374]).

**EGYPT**: Assaka, region de Suez, 10.III.1904, *Kneucker 2497* (G).





**Fig. 1.** – **A.** Flowers of *Cleome amblyocarpa* Barratte & Murb. var. *amblyocarpa*; **B.** Flowers of *C. amblyocarpa* var. *glandulosa* (Forssk.) Botsch.; **C–D.** Fruits of *C. amblyocarpa* var. *amblyocarpa* (up) and *C. var. glandulosa* (Forssk.) Botsch. (down).  
[Photos: C. Lemmel]




Fig. 2. – Distribution map of *Cleome arabica* L. (red), *C. amblyocarpa* Barratte & Murb. var. *amblyocarpa* (black), and *C. amblyocarpa* var. *glandulosa* (Forssk.) Botsch. (orange) in North Africa and neighbouring countries. All data based on specimens examined. [available on <https://efloramaghreb.org> (except for Libya)]

## Acknowledgements

We thank Ulf Arup, Zahora Attioui, Mathieu Chambouleyron, Alain Dobignard, Ori Fragman-Sapir, Annie Garcin, Abdelmonaim Homrani-Bakali, Jean-Paul Peltier and Errol Vela. Thanks are also due to Caroline Loup in MPU for her help in the Delile herbarium, Patrik Frödén for the photograph of Murbeck's type specimen, Eric Roalson for his collaboration, Martin Callmander, Joel Calvo, Adelaide Stork, and Nicolas Fumeaux for their helpful review.

## References

- ALI, S.I. & S.M.H. JAFRI (1976). *Flora of Libya*, vol. 11. Al Faateh University, Tripoli.
- BOTSCHANTZEV, V.P. (1964). Quid est *Cleome arabica* Jusl.? *Novosti Sist. Vyssh. Rast.* 1: 129–131.
- BOTSCHANTZEV, V.P. (1968). De nonnullis plantis florum algeriensis notae. *Novosti Sist. Vyssh. Rast.* 5: 234–242.
- BOULOS, L. (1999). *Flora of Egypt*, vol. 1. Al-Hadara Publishing, Cairo.
- FENNANE, M., M. IBN TATTOU, J. MATHEZ, A. OUYAHYA & J. EL OUALIDI (ed.) (1999). Flore pratique du Maroc. Manuel de détermination des plantes vasculaires, vol. 1. *Trav. Inst. Sci. Univ. Mohammed V, Sér. Bot.* 36.
- FORSKÅL, P. (1775). *Flora aegyptiaco-arabica*. Officina Mölleri, Hauniae. [<https://bibdigital.rjb.csic.es/idurl/1/10914>]
- FORSKÅL, P. (1776). *Icones rerum naturalium*. Officina Mölleri, Hauniae. DOI: <https://doi.org/10.5962/bhl.title.63481>
- HEDGE, I. & J. LAMOND (1970). Capparidaceae. In: RECHINGER K.H. (ed.), *Flora Iranica* 68: 1–39.
- LINNAEUS, C. (1755). *Centuria I. Plantarum*. L.M. Höjer, Upsaliae. DOI: <https://doi.org/10.5962/bhl.title.51985>
- LINNAEUS, C. (1759). *Amoenitates academicae*, vol. 4. Laurentii Salvii, Holmiae.
- LINNAEUS, C. (1763). *Species plantarum*, ed. 2, vol. 2. Laurentii Salvii, Holmiae. DOI: <https://doi.org/10.5962/bhl.title.11179>
- LINNAEUS, C. (1767). *Systema naturae*, ed. 12, vol. 2. Laurentii Salvii, Holmiae. DOI: <https://doi.org/10.5962/bhl.title.68927>
- LINNAEUS f., C. (1767). *Plantarum rariorum horti upsaliensis*. Siegfried Lebrecht Crusii, Lipsiae.
- MAIRE, R. (1965). Capparidaceae. In: *Flore de l'Afrique du Nord*, vol. 12: 115–139. Paul Lechevalier, Paris.
- MURBECK, S. (1905). Contributions à la connaissance de la flore du nord-ouest de l'Afrique et plus spécialement de la Tunisie. *Acta Univ. Lund.*, 2 1(4): 1–40. [<https://www.biodiversitylibrary.org/page/45448740>]
- OZENDA, P. (1958). *Flore du Sahara septentrional et central*. Centre National de la Recherche Scientifique, Paris.
- POWO (2024). *Plants of the World Online*. Royal Botanic Gardens, Kew. [<http://www.plantsoftheworldonline.org>]
- ROALSON, E.H. (2021). A revised synonymy, typification, and key to species of *Cleome* sensu stricto (Cleomaceae). *Phytotaxa* 496: 54–68. DOI: <https://doi.org/10.11646/phytotaxa.496.1.2>
- ZOHARY, M. (1966). *Flora Palaestina*, vol. 1. The Israel Academy of Sciences and Humanities, Jerusalem.