



Effect of Erythritol and Sucralose Formulation on the Survivorship of the Mosquito *Aedes aegypti*

Authors: Maestas, Lauren, Lee, Jana C., and Choi, Man-yeon

Source: Florida Entomologist, 106(2) : 129-132

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.106.0210>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Effect of erythritol and sucralose formulation on the survivorship of the mosquito *Aedes aegypti*

Lauren Maestas^{1,2}, Jana C. Lee³, and Man-yeon Choi^{3,*}

Erythritol (MW 122.1) is a 4-carboned polyol, zero-calorie artificial sweetener, with the active ingredient of Truvia®. Recently it was demonstrated to have insecticidal properties on multiple insect species: *Drosophila melanogaster* (Meigen) (Diptera: Drosophilidae) (Baudier et al. 2014); *Bactocera dorsalis* (Hendel) (Diptera: Tephritidae) (Zheng et al. 2016); *Drosophila suzukii* (Matsumura) (Diptera: Drosophilidae) (Choi et al. 2017; Sampson et al. 2017a); *Musca domestica* (L.) (Diptera: Muscidae) (Burgess & King 2017; Fisher et al. 2017); *Stomoxys calcitrans* (L.) (Diptera: Muscidae) (Burgess & Geden 2019); *Aedes aegypti* (L.) (Diptera: Culicidae) (Gilkey et al. 2018); and *Cacopsylla pyricola* (Förster) (Hemiptera: Psyllidae) (Wentz et al. 2020). Erythritol had minimal or no detectable non-target impacts on adult honey bees, *Apis mellifera* (L.) (Hymenoptera: Apidae) (Choi et al. 2019), or a predatory spider mite, *Tetranychus urticae* (Koch) (Trombidiformes: Tetranychidae) (Schmidt-Jeffris et al. 2021).

Sucrose (MW 342.3) is a commonly used phagostimulant, enhancing attractiveness and consumption of baits to insects. When combined with baits, sucrose can increase the efficacy of insecticides or toxic baits because insect pests are stimulated to feed more frequently (Vander Meer et al. 1995; Allan 2011; Cowles et al. 2015; Tochen et al. 2016; Roubos et al. 2019). However, adding sucrose provides a nutritional carbohydrate that will be metabolized in the same pests, if they are not killed immediately. Non-nutritive erythritol has been suggested as a potential alternative to sucrose as a phagostimulant. Further, it has been used to infect house flies with entomopathogenic fungi (Burgess et al. 2018), and in insecticides used in *D. suzukii* management (Gullickson et al. 2019). Although erythritol is being used as a phagostimulant, the sweetness of erythritol is about 30% less than sucrose (Perko & DeCock 2008). Sucrose-mixed diets were fed upon more than erythritol exclusive diets, which resulted in higher mortality to *D. suzukii* and mosquitoes (Choi et al. 2017; Gilkey et al. 2018). A similar result also was observed in the fire ant *Solenopsis invicta* Buren (Hymenoptera: Formicidae) (Vander Meer et al. 1995). Given the lower sweetness of erythritol, adding sucrose into the formulation enhances insecticidal activity. However, previous formulations contained 0.5 M sucrose, or a 35.4% sucrose to erythritol concentration, which is very sticky (Tang et al. 2017; Price et al. 2021). Also, if insects ingest a non-lethal dose, the sucrose may sustain them. Thus, it is desirable to find a non-nutritive, less sticky phagostimulant that is sweeter than sucrose, requiring a lower concentration.

Sucralose (MW 397.6) is the main ingredient of Splenda, a sucrose substitute in foods (Binns 2003). Because sucralose is about 600 times sweeter than sucrose, adding 0.1 M sucralose to 1.5 M erythritol (0.1 M sucralose + 1.5 M erythritol formulation) is sweeter than erythritol alone and is similar in sweetness to the 1.5 M erythritol + 0.5 M sucrose formulation. *Drosophila suzukii* flies fed more on an erythritol + sucralose formulation than erythritol alone, and died faster (Price et al. 2021).

The yellow fever mosquito, *Aedes aegypti*, is a vector of human pathogens leading to morbidity and mortality (Pridgeon et al. 2008). One disease in particular, Dengue virus, is one of the fastest growing global diseases (Brady & Hay 2020). A long history of chemical use has led to broad spectrum resistance to insecticides (Estep et al. 2017), and alternate natural and biological control methods for disease vectors are needed (Goolsby et al. 2022). Here, we tested erythritol formulations combined with sucrose or sucralose to identify insecticidal activities on adult *A. aegypti*. Our results showed that the erythritol formulation with sucralose reduced the survivorship of adult mosquitoes, and this less sticky erythritol formulation may have merit as an alternative method for mosquito control.

Adult yellow fever mosquitoes (*A. aegypti*) strain (ORL) 1952 were reared as described in Pridgeon et al. (2008). Mosquitos were subjected to 4 treatments: (1) water control, (2) 0.5 M sucrose, (3) 1.5 M erythritol (99% purity, Oakwood Products, Estill, South Carolina, USA) and 0.5 M sucrose (Fischer Scientific, Hampton, New Hampshire, USA), and (4) 1.5 M erythritol and 0.1 M sucralose (Fischer Scientific). We ran 3 reps per treatment across 3 eight-d sessions (9 reps per treatment total), with 20 adult males and 20 adult females per cup. The solution was delivered in a cotton-soaked wick and changed as needed. The cumulative proportion dead in each replicate cup was recorded each d, and male and female data were analyzed separately. The proportion of dead mosquitoes were analyzed by treatment, d, and interaction (fixed effects), trial as a random effect, and each cup as a random subject effect with repeated measures in a generalized linear mixed model in Proc Glimmix (SAS Institute 2016) with the best fit distribution based on residuals. A Tukey HSD test was used to compared treatment means.

The mortality assay tested insecticidal effects when *A. aegypti* adults consumed erythritol, sucrose, and sucralose formulations. Mosquito survivorship significantly decreased for 7 d (Fig. 1) for both fe-

¹USDA-ARS, Center for Medical, Agricultural, and Veterinary Entomology, Mosquito & Fly Research Unit, 1600-1700 S.W. 23rd Drive, Gainesville, Florida 32608; E-mail: lauren.maestas@usda.gov (L. M.)

²USDA-ARS, Cattle Fever Tick Research Unit, 22675 North Moorefield Road, Edinburg, Texas 78028, USA; E-mail: lauren.maestas@usda.gov (L. M.)

³USDA-ARS, Horticultural Crops Research Unit, 3420 N.W. Orchard Avenue, Corvallis, Oregon 97330, USA; E-mail: man-yeon.choi@usda.gov (M. Y. C.); jana.lee@usda.gov (J. C. L.)

*Corresponding author; E-mail: man-yeon.choi@usda.gov




Fig. 1. Survivorship of *Aedes aegypti* after consumption of erythritol formulations: E + S (1.5 M erythritol + 0.5 M sucrose), E + Sul (1.5 M erythritol + 0.1 M sucralose), Sucrose (0.5 M sucrose) as a positive control, and Water as a negative control for 7 d. Different letters denote ($*P < 0.05$) differences by Tukey HSD test.

males (treatment $F_{3,44} = 700.6$; $P < 0.001$; $d F_{6,264} = 461.0$; $P < 0.001$; treatment* $d F_{18,264} = 69.7$; $P < 0.001$) and males (treatment $F_{3,44} = 32.6$; $P < 0.001$; $d F_{6,264} = 40.8$; $P < 0.001$; treatment* $d F_{18,264} = 6.75$; $P < 0.001$). The fastest and highest mortality occurred in the 1.5 M erythritol + 0.1 M sucralose (E + Sul), followed by 1.5 M erythritol + 0.5 M sucrose (E + S) in both males and females, with mortality faster than the water only (a negative control) in females. Mortality was not statistically different between E + Sul and E + S treatments. In general, the females were more robust, and a greater proportion of females survived than males (Fig. 1). In this study, we did not test consumption of erythritol only because 1 M erythritol killed all *A. aegypti* mosquitoes within 5 d (Gilkey et al. 2018).

In our previous studies on *D. sukukii*, the most effective erythritol formulations were 1.5 M erythritol + 0.5 M sucrose (35.4% sugar solution) (Tang et al. 2017) and 1.5 M erythritol + 0.1 M sucralose (22.3% sugar solution) (Price et al. 2021). In *A. aegypti*, the insecticidal effect of 1M erythritol + 1 M sucrose (46.4% sugar solution) on the mosquito was greater than the mortality of erythritol alone (Gilkey et al. 2018). Although adding sucrose in erythritol formulations enhanced the insecticidal effect, it creates a sticky solution with more than a 30% sugar concentration (Tang et al. 2017). If the sucrose amount is reduced to less than 0.5 M, the erythritol consumption by the fly will decrease due to the low sweetness of the formulation, resulting in a lower insecticidal effect (Choi et al. 2017). We have observed that mosquitoes avoid a solution that is unpalatable, even to the point of death from desiccation (L. M., unpublished data). Therefore, a 1.5 M erythritol (> 18.3%) and 0.5 M sucrose (> 17.1%) (= 35.4% sugar concentration total) was used in our study, though it is still a sticky solution.

As an alternative to sucrose, sucralose will sweeten the formulation with a minimal concentration (i.e., 0.1 M), will not provide nutritional carbohydrates, and is less sticky in erythritol formulations. In fact, the 1.5 M erythritol + 0.1 M sucralose formulation has a 22.3% sugar concentration, and was the most effective formulation to kill *D. sukukii* in various sized arena tests (Price et al. 2021). A variety of artificial sweeteners were toxic causing mortality, decreased fecundity, and negative physiological impacts such as osmotic imbalance on different life stages from various insects (reviewed by O'Donnell et al. 2016, 2018; Sampson et al. 2017b, 2019; Tang et al. 2017; Lee et al. 2021). We hypothesized the mode of action of the zero-calorie sweeteners is that insects ingest non-metabolizable carbohydrates, starve, and experience hyperosmotic pressure in the body, causing a decrease in fly fitness (Price et al., 2022). In mosquitoes, erythritol and sucralose likely are non-metabolizable and phagostimulative to *A. aegypti* adults, and the formulation could be used as a human-safe insecticide to control mosquitoes.

We thank Neil Sanscrainte and Alden Estep for technical support and discussion on the mosquito experiment.

Summary

Erythritol, a non-nutritive 4-carboned polyol, was demonstrated to have insecticidal properties for multiple arthropod pests. The erythritol formulations combined with sucrose, as a phagostimulant, enhanced consumption and insecticidal efficiency on the pests. However, adding sucrose contributes nutritional carbohydrate resources in the same pests, can lead to the development of microbes in the field, and creates a sticky residue on plants sprayed with the solution. In this study, we tested and compared erythritol formulations combined with sucrose or sucralose, and identified insecticidal activities on adults of the mosquito *Aedes aegypti*. The

erythritol and sucralose formulation is a less sticky solution, and significantly reduced the survivorship of adult mosquitoes, it may have merit as an alternative method for mosquito control and warrants further evaluation.

Key Words: artificial sweeteners; zero-calorie sugar; human-safe insecticide; mosquito control

Sumario

Se demostró que el eritritol, un poliol de 4 carbonos no nutritivo, tiene propiedades insecticidas para múltiples plagas de artrópodos. Las formulaciones de eritritol combinadas con sacarosa, como fagoestimulante, mejoraron el consumo y la eficacia insecticida sobre las plagas. Sin embargo, el agregar sacarosa aporta recursos nutricionales de carbohidratos en las mismas plagas, que puede conducir al desarrollo de microbios en el campo y crear un residuo pegajoso en las plantas rociadas con la solución. En este estudio, probamos y comparamos formulaciones de eritritol combinadas con sacarosa o sucralosa, e identificamos actividades insecticidas en adultos del mosquito *Aedes aegypti*. La formulación de eritritol y sucralosa es una solución menos pegajosa y redujo significativamente la sobrevivencia de los mosquitos adultos, puede tener mérito como método alternativo para el control de mosquitos y justifica una evaluación adicional.

Palabras Clave: azúcar artificial; azúcar sin calorías; insecticida seguro para humanos; control de mosquitos

References Cited

- Allan SA. 2011. Susceptibility of adult mosquitoes to insecticides in aqueous sucrose baits. *Journal of Vector Ecology* 36: 59–67.
- Baudier KM, Kaschock-Marenda SD, Patel N, Diangelus KL, O'Donnell S, Marenda DR. 2014. Erythritol, a non-nutritive sugar alcohol sweetener and the main component of Truvia®, is a palatable ingested insecticide. *PLoS One* 9: e98949. doi: 10.1371/journal.pone.0098949
- Binns NM. 2003. Sucralose – all sweetness and light. *British Nutrition Foundation Nutrition Bulletin* 28: 53–58.
- Brady OJ, Hay SI. 2020. The global expansion of Dengue: how *Aedes aegypti* mosquitoes enabled the first pandemic Arbovirus. *Annual Review of Entomology* 65: 191–208.
- Burgess IV ER, Geden CJ. 2019. Larvicidal potential of the polyol sweeteners erythritol and xylitol in two filth fly species. *Journal of Vector Ecology* 44: 11–17.
- Burgess IV ER, King BH. 2017. Insecticidal potential of two sugar alcohols to *Musca domestica* (Diptera: Muscidae). *Journal of Economic Entomology* 110: 2252–2258.
- Burgess IV ER, Johnson DM, Geden CJ. 2018. Mortality of the house fly (Diptera: Muscidae) after exposure to combinations of *Beauveria bassiana* (Hypocreales: Clavicipitaceae) with the polyol sweeteners erythritol and xylitol. *Journal of Medical Entomology* 55: 1237–1244.
- Choi MY, Lucas H, Sagili R, Cha DH, Lee JC. 2019. Effect of Erythritol on *Drosophila sukukii* (Diptera: Drosophilidae) in the presence of naturally-occurring sugar sources, and on the survival of *Apis mellifera* (Hymenoptera: Apidae). *Journal of Economic Entomology* 112: 981–985.
- Choi MY, Tang SB, Ahn SJ, Amarasekare KG, Shearer P, Lee JC. 2017. Effect of non-nutritive sugars to decrease the survivorship of spotted wing drosophila, *Drosophila sukukii*. *Journal of Insect Physiology* 99: 86–94.
- Cowles RS, Rodriguez-Saona C, Holdcraft R, Loeb GM, Elsensohn JE, Hesler SP. 2015. Sucrose improves insecticide activity against *Drosophila sukukii* (Diptera: Drosophilidae). *Journal of Economic Entomology* 108: 640–653.
- Estep AS, Sanscrainte ND, Waits CM, Louton JE, Becnel JJ. 2017. Resistance status and resistance mechanisms in a strain of *Aedes aegypti* (Diptera: Culicidae) from Puerto Rico. *Journal of Medical Entomology* 54: 1643–1648.
- Fisher ML, Fowler FE, Denning SS, Watson DW. 2017. Survival of the house fly (Diptera: Muscidae) on truvia and other sweeteners. *Journal of Medical Entomology* 54: 999–1005.
- Gilkey PL, Bolshakov DT, Kowala JG, Taylor LA, O'Donnell S, Marenda DR, Sirot LK. 2018. Lethal effects of erythritol on the mosquito *Aedes aegypti* Linnaeus (Diptera: Culicidae). *Journal of Applied Entomology* 142: 873–881.

- Goolsby JA, Maestas L, Saelao P, Lohmeyer KH. 2022. Evaluation of repellency of Stop the Bites® botanical pesticide to white-tailed deer at corn feeders. *Southwestern Entomologist* 47: 277–284.
- Gullickson MG, Rogers MA, Burkness EC, Hutchison WD. 2019. Efficacy of organic and conventional insecticides for *Drosophila suzukii* when combined with erythritol, a non-nutritive feeding stimulant. *Crop Protection* 125: 104878. doi.org/10.1016/j.cropro.2019.104878
- Lee SH, Choe DH, Lee CY. 2021. The impact of artificial sweeteners on insects. *Journal of Economic Entomology* 114: 1–13.
- O'Donnell S, Baudier K, Marenda DR. 2016. Non-nutritive polyol sweeteners differ in insecticidal activity when ingested by adult *Drosophila melanogaster* (Diptera: Drosophilidae). *Journal of Insect Science* 16: 47. doi: 10.1093/jisesa/iew031
- O'Donnell S, Baudier K, Fiocca K, Marenda DR. 2018. Erythritol ingestion impairs adult reproduction and causes larval mortality in *Drosophila melanogaster* fruit flies (Diptera: Drosophilidae). *Journal of Applied Entomology* 142: 37–42.
- Perko R, DeCock P. 2008. Erythritol, p. 157. In Mitchell H [Ed.], *Sweeteners and Sugar Alternatives in Food Technology*. Wiley, Chichester, United Kingdom.
- Price BE, Lee JC, Choi MY. 2021. Erythritol combined with non-nutritive sucralose increases feeding by *Drosophila suzukii*, quickens mortality and reduces oviposition. *Crop Protection* 150: e105812. doi.org/10.1016/j.cropro.2021.105812
- Price BE, Yoon JS, Choi MY, Lee JC. 2022. Effects of nonnutritional sugars on lipid and carbohydrate content, physiological uptake, and excretion in *Drosophila suzukii*. *Archives of Insect Biochemistry and Physiology* 109: e21860. doi: 10.1002/arch.21860
- Pridgeon JW, Pereira RM, Becnel JJ, Allan SA, Clark GG, Linthicum KJ. 2008. Susceptibility of *Aedes aegypti*, *Culex quinquefasciatus* Say, and *Anopheles quadrimaculatus* Say to 19 pesticides with different modes of action. *Journal of Medical Entomology* 45: 82–87.
- Roubos CR, Gautam BK, Fanning PD, Van Timmeren S, Spies J, Liburd OE, Isaacs R, Curry S, Little BA, Sial AA. 2019. Impact of phagostimulants on effectiveness of OMRI-listed insecticides used for control of spotted-wing drosophila (*Drosophila suzukii* Matsumura). *Journal of Applied Entomology* 143: 609–625.
- Sampson BJ, Werle CT, Stringer SJ, Adamczyk JJ. 2017a. Ingestible insecticides for spotted wing *Drosophila* control: a polyol, erythritol, and an insect growth regulator, lufenuron. *Journal of Applied Entomology* 141: 8–18.
- Sampson BJ, Easson MW, Stringer SJ, Werle CT, Magee D, Adamczyk JJ. 2019. Laboratory and field assessments of erythritol derivatives on the survival, reproductive rate, and control of *Drosophila suzukii* (Diptera: Drosophilidae). *Journal of Economic Entomology* 112: 173–180.
- Sampson BJ, Marshall DA, Smith BJ, Stringer SJ, Werle CT, Magee DJ, Adamczyk JJ. 2017b. Erythritol and Lufenuron detrimentally alter age structure of wild *Drosophila suzukii* (Diptera: Drosophilidae) populations in blueberry and blackberry. *Journal of Economic Entomology* 110: 530–534.
- SAS Institute. 2016. SAS vers. 9.4. SAS Institute, Cary, North Carolina, USA.
- Schmidt-Jeffris RA, Beers EH, Smytheman P, Rehfield-Ray L. 2021. Erythritol, an artificial sweetener, is acaricidal against pest mites and minimally harmful to a predatory mite. *Journal of Economic Entomology* 114: 1701–1708.
- Tang SB, Lee JC, Jung JK, Choi MY. 2017. Effect of erythritol formulation on the mortality, fecundity and physiological excretion in *Drosophila suzukii*. *Journal of Insect Physiology* 101: 178–184.
- Tochen S, Walton VM, Lee JC. 2016. Impact of floral feeding on adult *Drosophila suzukii* survival and nutrient status. *Journal of Pest Science* 89: 793–802.
- Vander Meer RK, Lofgren CS, Seawright JA. 1995. Specificity of the red imported fire ant (Hymenoptera: Formicidae) phagostimulant response to carbohydrates. *Florida Entomologist* 78: 144–154.
- Wentz K, Cooper WR, Horton DR, Kao R, Nottingham LB. 2020. The artificial sweetener, erythritol, has insecticidal properties against pear psylla (Hemiptera: Psyllidae). *Journal of Economic Entomology* 113: 2293–2299.
- Zheng C, Zeng L, Xu Y. 2016. Effect of sweeteners on the survival and behaviour of *Bactrocera dorsalis* (Hendel) (Diptera: Tephritidae). *Pest Management Science* 72: 990–996.