

Genetic Diversity and Dispersal Potential of the Stonefly *Dinocras cephalotes* in a Central European Low Mountain Range

Authors: Elbrecht, Vasco, Feld, Christian K., Gies, Maria, Hering, Daniel, Sondermann, Martin, et al.

Source: Freshwater Science, 33(1) : 181-192

Published By: Society for Freshwater Science

URL: <https://doi.org/10.1086/674536>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Genetic diversity and dispersal potential of the stonefly *Dinocras cephalotes* in a central European low mountain range

Vasco Elbrecht^{1,3}, Christian K. Feld^{2,4}, Maria Gies^{2,5}, Daniel Hering^{2,6}, Martin Sondermann^{2,7}, Ralph Tollrian^{1,8}, and Florian Leese^{1,9}

¹Department of Animal Ecology, Evolution and Biodiversity, Faculty of Biology and Biotechnology, Ruhr University Bochum, Universitätsstrasse 150, 44801 Bochum, Germany

²Department of Aquatic Ecology, Faculty of Biology, University Duisburg-Essen, Universitätsstrasse 5, 45141 Essen, Germany

Abstract: Aquatic insects are widely used as indicator taxa to assess the ecological state of streams and to evaluate the success of stream restoration projects. Information on intraspecific genetic diversity and population connectivity is often lacking for such indicator taxa. However, these parameters are of critical importance for restoration plans and conservation management because: 1) species sometimes consist of several cryptic species and 2) species can recolonize only those restored habitats within a reachable distance from their source populations. Gene flow generally cannot be observed directly, and molecular markers provide a reasonable alternative to assess the dispersal potential and evaluate species' genetic diversity. We investigated the genetic diversity and dispersal potential of the predatory stonefly *Dinocras cephalotes* using 323 specimens from 29 populations in the Sauerland, a low mountain range in Germany. We used a 658 base pair (bp) fragment of the mitochondrial cytochrome *c* oxidase subunit I gene (COI) and found 2 distinct and diverse haplotype groups, which were shared across most populations. The groups were separated by a minimum intraspecific *p*-distance of 4.3%, suggesting historic isolation and possible presence of cryptic species. However, complementing analyses of the nuclear Wingless gene and 3 newly developed microsatellite markers clearly showed that individuals from both COI haplotype groups are interbreeding, and therefore, *D. cephalotes* is considered a single valid species. Population comparisons indicated high connectivity among all populations, with only a few individual populations showing signatures of isolation. Based on the molecular data, we conclude that dispersal is primarily achieved by the adult females of *D. cephalotes*.

Key words: population genetics, COI, historic isolation, cryptic species, gene flow, landscape genetics, restoration ecology, wingless, microsatellites, historic isolation

Human activity has dramatically altered and degraded stream ecosystems (Poff et al. 2007) and poses a threat to global freshwater biodiversity (Vörösmarty et al. 2010). The loss of biodiversity threatens ecosystem functioning (Vaughn 2010), with potential direct negative consequences for the provision of ecosystem services (Cardinale 2011). The Water Framework Direc-

tive (WFD) of the European Union explicitly obliges its member states to counteract degradation of freshwater ecosystems and demands good ecological and chemical status of surface waters by 2015 (Directive 2000/60/EC, Annex V; European Union 2000). In a recent report on the status of Europe's waters, the European Environment Agency concluded that most

E-mail addresses: ³vasco.elbrecht@rub.de; ⁴christian.feld@uni-due.de; ⁵maria.gies@uni-due.de; ⁶daniel.hering@uni-due.de; ⁷martin.sondermann@uni-due.de; ⁸tollrian@rub.de; ⁹To whom correspondence should be addressed, florian.leese@rub.de

DOI: 10.1086/674536. Received 17 February 2013; Accepted 28 June 2013; Published online 22 January 2014.
Freshwater Science. 2014. 33(1):181–192. © 2014 by The Society for Freshwater Science.

surface-water bodies in Europe are unlikely to meet this target by 2015 (EEA 2012), and many management and restoration activities have been launched to implement this ambitious Directive.

The criteria for attaining good ecological status have not yet been fully developed, but the recovery of intact freshwater communities is the key to improve ecological conditions in lotic systems (Palmer et al. 1997, 2010, Jähnig et al. 2009, Feld et al. 2011). A primary assessment tool for quantifying the ecological status of streams is analysis of the biodiversity and abundance of biological indicator species, in particular macroinvertebrates (Hellawell 1986, Metcalfe 1989, Rosenberg and Resh 1993, Hering et al. 2006). Once restoration projects are completed, the native bioindicator organisms must be capable of reaching the restored ecosystems from source populations (Palmer et al. 1997, Lake et al. 2007, Smith et al. 2009). Successful recolonization depends on several additional variables, such as the individual species' life cycle (holo- vs merolimnic) and its duration, physical dispersal traits (flying, crawling, active, and passive), behavioral patterns, the spatial distribution of source populations in the stream network, and the presence of barriers that hinder dispersal (Smith et al. 2009). Macroinvertebrate species often may be unable to recolonize restored habitats successfully, despite favorable habitat conditions at the restored site (Lake et al. 2007). In such cases, the absence of macroinvertebrate indicator taxa is not automatically indicative of poor habitat quality and misleads stream-quality assessment.

Empirical or experimental data on the dispersal abilities of aquatic insects are scarce and, therefore, often neglected in restoration plans (Palmer et al. 1997, Smith et al. 2009). Studies of aquatic insects with stable isotopes (Coutant 1982, Briers et al. 2004), mark-and-recapture experiments (Stettmer 1996, Hassall and Thompson 2011), and light traps (e.g., Kovats et al. 1996) indicate that long-distance dispersal or passive drift >1 km from the stream channel is possible for winged insects. Although a strong preference of insects to stay in the vicinity of the stream channel can be observed (90% of adult stoneflies stayed within the stream channel; Briers et al. 2002), rare but successful long-distance flights may connect populations. In the absence of direct observations, molecular tools allow assessment of the connectivity of populations with comparisons of allele diversity

and frequencies among populations (Hughes et al. 2008). These markers also help identify overlooked or cryptic species, which show no or only subtle morphological differences (Pfrender et al. 2010, Zhou et al. 2010).

We analyzed partitioning of genetic diversity in the predatory stonefly *Dinocras cephalotes* (Curtis 1827) in the Sauerland region, a low mountain range in western Germany. The larvae of *D. cephalotes* inhabit cold and fast-flowing streams and are reliable indicators of good water quality (Eiseler and Enting 2012). *Dinocras cephalotes* has a life cycle of ~3 y (Iannilli et al. 2002), and only the female imagines have fully developed wings (Tierno de Figueroa et al. 2006). The dispersal potential of *D. cephalotes* at local and regional scales has been questioned because they were described as clumsy flyers (Ketmaier et al. 2001) and population subdivisions were reported even on very small geographical scales (Ketmaier et al. 2001).

The Sauerland region has been anthropogenically influenced since medieval times by agriculture, forestry, ore mining, and metal production, which in concert, led to severely impacted stream networks. However, many headwater streams in the Sauerland region are chemically and ecologically classified as being in good condition. Isolated and genetically depauperate headwater populations of *D. cephalotes* might be expected because of the heavy influence of several anthropogenic stressors (e.g., pollution and fragmentation by weirs, dams, and hydromorphological alterations) on higher-order streams in the network. This isolation may be especially strong for *D. cephalotes* because flight capability is restricted to females.

The aim of our study was to use mitochondrial and nuclear deoxyribonucleic acid (DNA) markers to test whether populations of *D. cephalotes* in different headwater streams are isolated and genetically depauperate, which in turn, would increase demographic stochasticity and the risk of local extinctions. The regional dispersal potential, which is the prerequisite to recolonize restored stream habitats, of *D. cephalotes* and its value as an indicator taxon for assessing restoration measures were assessed.

METHODS

Sample collection

Dinocras cephalotes populations in headwater streams were sampled, mainly in the Ruhr river ba-

sin of North Rhine-Westphalia, Germany (rivers Ruhr and Lenne; Fig. 1, Table S1) from May to June 2010–2012 and stored in 80% ethanol at -20°C . A total of 323 specimens from 29 populations were analyzed with molecular methods.

Microsatellite development

Microsatellites were developed from an unenriched sequence library that was sequenced on 454 GS Junior sequencer (Roche, Basel, Switzerland). One microgram of high-quality DNA was used for library creation according to the manufacturer's protocol. Resulting reads were quality-controlled (FastQC, version 0.10.1; <http://www.bioinformatics.babraham.ac.uk/projects/fastqc/>), analyzed for potential contamination (Blast+, version 2.2.26; Camacho et al. 2009), and subsequently assembled using MIRA (version 3.4.0.1; Chevreaux et al. 1999) to exclude potential multicopy sequences (settings as in Leese et al. 2012 except `-AL:mrs=70:mo=20-CL:pec=off`). Microsatellites were identified with Phobos (version 1.0.6; Mayer 2006) and a custom R script (version 2.15.1; R Project for Statistical Computing, Vienna, Austria) was used to select the best candidate sequences based on flanking region

length, repeat motif, and number. Sequences were inspected in Geneious Pro (version 6.0.5; Kearse et al. 2012). Primers were developed with the Primer3 plugin (Rozen and Skaletsky 2000).

Laboratory methods

All specimens were identified and photographed prior to DNA extraction. DNA was extracted following a modified salt-extraction protocol (Sunnucks and Hales 1996). Negative controls were included for DNA extractions and polymerase chain reactions (PCR). A fragment of the mitochondrial barcoding gene cytochrome *c* oxidase subunit I (COI) was amplified with standard invertebrate primers HCO2198 and LCO1490 (Folmer et al. 1994) in a reaction consisting of $1\times$ PCR buffer, 2.5 mM MgCl_2 , 0.2 mM deoxyribonucleotide triphosphates (dNTPs), 0.5 μM of each primer, 0.02 U/ μL Euro Taq (EuroClone, Milano, Italy), 1 μL DNA, filled up to a total volume of 25 μL with high-performance liquid chromatography (HPLC) H_2O (PCR program: $94^{\circ}\text{C}/120\text{ s}$, 36 cycles of [$94^{\circ}\text{C}/20\text{ s}$, $46^{\circ}\text{C}/30\text{ s}$, $72^{\circ}\text{C}/60\text{ s}$], $72^{\circ}\text{C}/7\text{ min}$). PCR success was confirmed with agarose gel electrophoresis, and samples that could not be amplified were repeated using HotMaster




Figure 1. Haplotype map of 307 *Dinocras cephalotes* specimens analyzed for the marker cytochrome *c* oxidase subunit I (COI). Each population is represented by a circle, with the population size indicated by an outer white circle. Samples that carry a haplotype from group A are indicated in black, other samples belong to haplotype group B. In the haplotype group B, the 2 most common haplotypes are indicated by lines (h1) and dots (h5). Populations that were investigated with the Wingless marker are highlighted with an asterisk. The dashed line indicates the east/west grouping used in the analysis of molecular variance. The dotted lines indicate the divide of the Ruhr and Lenne drainage basins. Small inset shows location of study area in Germany. Detailed information about the populations is available in Table S1.

Taq (5Prime; Gaithersburg, Maryland; parameters identical as for Euro Taq, but with 65°C extension temperature). A fragment of the nuclear Wingless gene was amplified from 68 samples of 10 populations (illustrated with an asterisk in the haplotype map; Fig. 1) according to the protocol described by Pauls et al. (2008). Populations for additional Wingless analysis were selected to cover a wide geographic range and, if possible, to include individuals belonging to both COI haplotype groups. Ten µL of PCR product were purified enzymatically with 0.5 µL Exonuclease I (20 U/µL; Thermo Fisher Scientific, Waltham, Massachusetts) and 1 µL FastAP (1 U/µL, Thermo Fisher Scientific) by incubating in a thermocycler at 37°C for 15 min followed by 96°C for 15 min prior to sequencing (Werle et al. 1994). Bidirectional Sanger sequencing was carried out on an ABI 3730 sequencer (Applied Biosystems, Carlsbad, California) by GATC Biotech (Konstanz, Germany). Ambiguous and short sequences were resequenced.

Microsatellite markers were optimized with a gradient PCR (annealing temperature range: 45–69°C) and PCR enhancers (dimethylsulfoxide [DMSO]: Carl Roth, Karlsruhe, Germany; Betaine: Sigma–Aldrich, Steinheim, Germany). The PCR settings were as described above for COI, but with a concentration of 0.2 µM for each forward and reverse primer, 0.05 µM of the tailed M13 primer (5'-CACGACGTTGTAAAA CGAC-3'), and 0.02 U/µL Euro Taq (PCR program: 94°C/120 s, 36 cycles of [94°C/20 s, 51–63°C/30 s, 72°C/60 s], 72°C/45 min). Primers that amplified reliably for a subset of samples were used for all samples. Allele sizes were determined by acrylamide gel electrophoresis on a Li-Cor analyzer 4300 with the software Saga2 GT (Li-Cor Biosciences, Lincoln, Nebraska). Alleles that could not be identified reliably were rerun or scored as missing data.

Sequence data analysis

Sequences with sufficient length and quality were assembled and an alignment was constructed using the MAFFT plugin (version 1.3; Katoh et al. 2002) for Geneious. Uncorrected genetic *p*-distances between haplotypes were calculated with MEGA (version 5.05; Tamura et al. 2011). To test for genetic fixation between populations, pairwise F_{ST} and Φ_{ST} estimators were calculated for the COI data set using Arlequin (version 3.5.1.3; Excoffier et al. 2005).

Statistical significance was assessed with 1000 iterations and significance level was Bonferroni corrected. Isolation by distance was tested with a Mantel test (10,000 replications; R package *ade4*; Dray and Dufour 2007) using the genetic differentiation measures F_{ST} and Φ_{ST} with 3 distance measures: direct distance between populations, shortest distance following the streams, and elevation differences between populations (calculated with QGIS, version 1.8; Quantum GIS Development Team; <http://qgis.osgeo.org>). Population partitioning with east/west and grouping by catchments was tested with an analysis of molecular variance (AMOVA) as implemented in Arlequin. Grouping was used to test for local adaptations resulting from effects related to altitude and isolation between catchments in case of poor dispersal. A minimum spanning network was calculated with Arlequin and visualized with HapStar (version 0.6; Teacher and Griffiths 2010).

Wingless sequences were assembled like the COI sequences, and a minimum spanning network generated as described above. Wingless haplotypes were compared to the respective COI haplotypes of the 63 tested specimens. The Wingless marker is a nuclear gene and was sequenced to validate the patterns found with the mitochondrial marker COI. A relatively small sample size for the Wingless gene was sufficient to test whether the patterns of both markers were similar.

Microsatellite data analysis

The microsatellite data were checked for scoring errors with MicroChecker (version 2.2.3; Van Oosterhout et al. 2004) and for deviations from Hardy–Weinberg and linkage-equilibrium with Arlequin. To measure genetic differences between populations F_{ST} values were calculated with GenoDive (version 1.0b23; Meirmans and Van Tienderen 2004). D_{est} was calculated with the R package *DEMEtics* (version 0.8-5; Jueterbock et al. 2012) with Bonferroni-corrected *p*-values (populations with a sample size = 1 were excluded from the analysis). F_{ST} between the 2 main haplotype groups for the COI marker was calculated by creating 2 artificial populations, each containing the microsatellite data for all individuals of haplotype group A or group B (GenoDive, 1000 iterations). The Mantel test and AMOVA were calculated as described for the COI data. In addition, the micro-

satellite data were analyzed for population clustering using STRUCTURE (version 3.2.4; Pritchard et al. 2000, Falush et al. 2003; default settings, burn-in = 10,000 followed by 50,000 Markov Chain Monte Carlo steps) and the most likely number of clusters was determined with the Evanno method (Evanno et al. 2005) as implemented in STRUCTURE HARVESTER (version 0.6.93; Earl and vonHoldt 2011; <http://taylor0.biology.ucla.edu/structureHarvester/>). Last, a principal component analysis (PCA) was carried out in the R package *Adegenet* (Jombart 2008) for data conversion (function *scaleGen* and *dudi.pca*) as implemented in the package *ade4*.

RESULTS

Haplotype groups (COI and Wingless)

Reliable COI sequences of 658 bp length were obtained for 307 specimens (GenBank accession numbers KF410897–KF410943). The other sequences were excluded because of poor read quality, short read length, or double peaks. Some sequences that still showed double peaks after PCR and sequencing reactions were repeated and may hint at the presence of pseudogenes (numts) or heteroplasmy. Reliable sequences of 400 bp length were obtained for all of the 68 samples analyzed for the Wingless gene (GenBank accession numbers KF442621–KF442625). However, 2 single-nucleotide polymorphisms (SNPs) were observed in 5 sequences. The individual peaks in the chromatograms showed very exact overlap, so phasing was not possible, and the 5 sequences were discarded from the data set.

A total of 47 unique COI haplotypes were found. These clustered into 2 diverse groups separated by

p -distances of 4.3 to 5.2% (Fig. 2A). The Wingless gene showed 1 major haplotype group that was less diverse (only 5 alleles) than the COI marker (Fig. 2B). Most Wingless alleles were shared by individuals from both COI haplotype groups (Fig. 2B). The COI haplotypes had a relatively homogeneous distribution across the study area (Fig. 1). A very weak but significant differentiation existed between the east/west groups (AMOVA, $\Phi_{CT} = 0.065$, $p < 0.001$; Table 1). Only a few pairwise population comparisons had significant F_{ST} values (uncorrected mean $F_{ST} = 0.1858$, $\sigma = 0.0751$, $p = 0.05$, $n = 34$ of 406; Fig. S1) or significant Φ_{ST} values (uncorrected mean $\Phi_{ST} = 0.2960$, $\sigma = 0.1605$, $p = 0.05$, $n = 32$ of 406; Fig. S2). Only 1 pair (F_{ST} values for GR and E03) remained significant after Bonferroni correction. The populations E13, E19, GR, NH showed slightly higher differentiation estimates than other populations (Figs. S1, S2). The differentiation values were unreliable for the 3 populations represented by single specimens (RU2, NH, BB) and were not considered further. A weak but significant positive correlation was found between Φ_{ST} and differences in altitude between populations (Mantel test, $r = 0.1692$, $p = 0.0097$; Fig. S3).

Microsatellite data analysis

Four of 15 developed microsatellite markers were used for all samples (Table 2). Possible stuttering problems were identified for markers C1 and L1 and possible null alleles for the markers C1, C2, and L1 (MicroChecker), but bands of all markers were clearly identifiable despite slight stuttering. All markers deviated slightly from Hardy–Weinberg equilibrium, but




Figure 2. Minimum Spanning Network of the mitochondrial cytochrome *c* oxidase subunit I (COI) haplotypes and the nuclear gene *Wingless* (numbers indicate the number of specimens with the respective haplotype). A.—Network of 307 COI sequences (white circles indicate singletons, black dots represent hypothetical haplotypes). Two main haplotype groups were identified as indicated by dashed circles (groups A and B, 4.3% minimum uncorrected sequence difference between groups). B.—*Wingless* network for individual alleles from 63 sequences. The numbers in the circles refer to the number of alleles that show a COI haplotype belonging to group A or B. The circle size corresponds to the frequency of the respective allele. The sequences were obtained from populations E03, E08, E09, E12, E20, E21, E23, E25, ME, and VA (indicated by asterisks in Fig. 1), encompassing individuals from both COI haplotype groups A and B.

Table 1. Results of 2 analyses of molecular variance (AMOVA) for the cytochrome *c* oxidase subunit I (COI) marker with grouping by catchments and grouping according to geographical position (east/west).

Source of variation	East/west			Ruhr/Lenne		
	Φ	p	% variance	Φ	p	% variance
Among groups	$\Phi_{CT} = 0.065$	<0.001	6.49	$\Phi_{CT} = -0.006$	0.937	-0.62
Among populations within groups	$\Phi_{SC} = -0.016$	0.700	-1.48	$\Phi_{SC} = 0.023$	0.026	2.26
Within populations	$\Phi_{ST} = 0.050$	0.180	94.99	$\Phi_{ST} = 0.016$	0.053	98.36

only markers C1 and L1 showed very strong deviations (Table 2). L1 was excluded from the data set because of strong heterozygote deficiency and possible linkage to markers C1 and C2.

With microsatellite markers, only a few pairwise population comparisons showed significant F_{ST} (uncorrected mean $F_{ST} = 0.009$, $\sigma = 0.039$, $p = 0.05$, $n = 36$ of 325; Fig. S4) or significant D_{est} values (uncorrected mean $D_{est} = 0.042$, $\sigma = 0.091$, $p = 0.05$, $n = 50$ of 325; Fig. S5). The populations E05, E20 and SO showed slightly higher differentiation values than other populations (Figs S4, S5) and did not match the populations differentiated in COI. Six comparisons had significant D_{est} values after Bonferroni correction. Most variance was found within populations (AMOVA; Table 3). River distance and D_{est} values were weakly but significantly correlated (Mantel test, $r = 0.1563$, $p = 0.0259$; Fig. S6). The structure analysis for all populations did not indicate distinct clusters (Fig. 3), and the data analysis with the Evanno method confirmed that one cluster is most likely. The PCA analysis with the microsatellite data clustered all populations together (overlapping ellipsoids; Fig. 4A). Some

populations showed deviations from the cluster. However these populations did not show a strong differentiation when analyzed for F_{ST} , Φ_{ST} , or D_{est} (COI and microsatellite data). Results of the same PCA analysis also were shown to group by COI haplotypes A and B instead of populations, and both haplotype groups showed a clear overlap (Fig. 4B). The microsatellite-based F_{ST} between the 2 haplotype groups was not significant ($F_{ST} = -0.006$, $p = 0.967$) indicating panmixia among members of the 2 groups at the nuclear level.

DISCUSSION

Evidence for cryptic species

An implicit requirement when analyzing the dispersal potential of a target species is that the candidate species does not consist of a complex of cryptic species. An obvious result of the COI analysis of *D. cephalotes* is the presence of great intraspecific distances (4.3–5.2%) and a prominent barcoding gap between members of 2 haplotype groups (groups A and B). Such genetic signatures often indicate the presence of cryptic or unrecognized species (Hebert et al.

Table 2. Overview of developed primer sequences used on 323 *Dinocras cephalotes* samples. Primers with an M13 extension at the 5' end are indicated by an asterisk. Hardy–Weinberg equilibrium proportions of expected (H_e) and observed (H_o) heterozygosity were calculated. Significant deviations are indicated by asterisks. * = $p < 0.05$, *** = $p < 0.001$. Temp = temperature, F = forward, R = reverse, A = adenine, T = thymine, G = guanine, C = cytosine.

Locus	Primer sequences (5'–3')	Temp (°C)	Repeat motif	No. of alleles detected	Size range	H_e	H_o	GenBank accession no.
C1	F: *GCTAAGATGAGAGCGGCTCCAGTG R: AAGTCGCCACCGTCCGTGAGA	62	(CA) ₈	10	163–211	69.8	36.1***	KF410944
C2	F: AACGCGCTGGTCGAGAACGTG R: *ATGGGCTGACGGCAGAAACC	54	(CA) ₈	21	254–326	81.6	77.0***	KF410945
L1	F: AGTCGTCGCTGCTGGTTCTG R: *CAACGCCTCGACGAGAGTGCC	61.5	(AT) ₁₀	4	126–138	69.2	12.2***	KF410946
L11	F: ACGTGTGAATCTCTCACTTC R: *GAAGGTGTAGTTGGGAAGC	51	(ATCA) ₇	4	164–176	51.1	52.4*	KF410947

Table 3. Results of 2 analyses of molecular variance (AMOVA) for 3 microsatellite markers with grouping by catchments and grouping according to geographical position (east/west).

Source of variation	East/west			Ruhr/Lenne		
	Φ	p	% variance	Φ	p	% variance
Among groups	$\Phi_{CT} = -0.003$	0.892	-0.26	$\Phi_{CT} = -0.000$	0.479	-0.01
Among populations within groups	$\Phi_{SC} = 0.012$	0.001	1.23	$\Phi_{SC} = 0.009$	0.003	0.94
Within populations	$\Phi_{ST} = 0.010$	0.001	99.04	$\Phi_{ST} = 0.009$	0.002	99.07

2004). Other plecopteran taxa analyzed so far have shown intraspecific COI distances below and above the typical barcoding gap threshold of 2 to 3% distance (Sweeney et al. 2011, Zhou et al. 2009, 2010), with intraspecific distances of up to 5.8% (Mynott et al. 2011). These studies relied only on COI data, so the possibility that they were dealing with cryptic species could not be ruled out. The variability of *D. cephalotes* is at the upper margin of values reported as intraspecific distances in the literature (intraspecific uncorrected distance up to 5.2%). A barcode gap of 4.3% between the 2 diverse COI haplotype groups A and B suggests the presence of cryptic species.

However, the barcode gap for the *D. cephalotes* populations also could be the result of historic isolation and independent lineage sorting in glacial refugia without subsequent reproductive isolation. Therefore, the absence of recombination within mitochondrial genes would lead to the persistence of these historically accumulated differences in secondary contact even under panmixia. Historic isolation has been discussed as a primary force underlying contemporary genetic variation in other aquatic insects

(Pauls et al. 2006, Lehrian et al. 2010, Bálint et al. 2011, Alp et al. 2012, Theissinger et al. 2012). The central question to be addressed with regard to such prominent differences in mitochondrial DNA is whether members of these groups still interbreed successfully in secondary contact. If a reproductive barrier had evolved (either because of, e.g., pleiotropic effects in small refugial populations or a specific differential selection regime), the genetic signatures of isolation also should be found in nuclear DNA markers. If interbreeding were still possible, recombination should homogenize the accumulated nuclear differences and lead to differing patterns between mitochondrial and nuclear DNA. To test both hypotheses, the nuclear Wingless marker was sequenced for a subset of individuals from both COI haplotype groups and compared to the COI data. In addition, COI data were compared to allele frequencies of 3 microsatellite markers. Individuals from both haplotype groups shared the same Wingless haplotype and the 3 microsatellites revealed no differences in allele frequencies between members of groups A and B. Thus, the hypothesis that haplotype groups A and B represent




Figure 3. Results of the Bayesian cluster analysis with STRUCTURE with 3 microsatellite loci for $K = 2$ to 4 (burn-in = 10,000, Markov Chain Monte Carlo [MCMC] steps = 50,000, 20 independent runs/ K). Individuals per population are sorted according to their membership coefficients to the black cluster (populations are separated by white lines). The longer the bar for an individual color, the higher the probability that an individual actually belongs to the respective cluster. No distinct substructures can be seen for any of the populations.




Figure 4. Principal Component Analysis (PCA) of 316 samples for the microsatellite markers C1, C2, and L11. A.—Ellipses encompass populations. B.—Identical PCA, but with distinction of common (group A) and rare (group B) cytochrome *c* oxidase subunit I (COI) haplotypes of the individuals. PC axis 1 and PC axis 2 explain 6.04% and 4.82% of the total variance.

genetically distinct cryptic or unrecognized species was rejected.

Connectivity of *D. cephalotes* populations

The COI haplotypes show a relatively homogeneous distribution across all populations, with a very weak differentiation between the eastern (higher altitude) and western (lower altitude) populations as revealed by significant F_{CT} in the AMOVA. However, only a few pairwise F_{ST} and Φ_{ST} comparisons showed significant differences even across larger geographical distances. These results indicate connectivity between populations. The small differences between eastern and western populations also could be explained by systematic differences in genotypes related to altitude as shown by the Mantel test.

The results for the 3 microsatellite loci are largely consistent with the results of the COI analysis, indicating stable population connectivity with only a few significant F_{ST} and D_{est} values. No patterns of isolation were revealed by the Bayesian clustering analysis in STRUCTURE or the PCA. However, detailed predictions about the number of migrating individu-

als per generation cannot be made because of the limited number of microsatellite loci investigated.

A very weak but significant correlation between D_{est} and river distance was found with the Mantel test. In addition, the differences between eastern and western COI populations found with the marker COI with the AMOVA were not found in the microsatellite data. These results contradict the COI-based results because gene flow among all groups, as indicated by the AMOVA, would affect the distribution of mitochondrial markers. The observed pattern of small but significant mitochondrial differentiation between lower-altitude western and higher-altitude eastern populations might be caused by directional selection acting on the mitochondrial genome (Ballard and Whitlock 2004) or the limited number of microsatellite loci investigated.

Dispersal strategy of *D. cephalotes*

Different dispersal strategies are described for aquatic insects (Hughes et al. 2009). Dispersal of *D. cephalotes* by larval migration within connected streams seems unlikely because a stronger correla-

tion between population differentiation and river distance would then be expected (Hughes et al. 2008). Furthermore, earlier studies based on genetic data showed that most aquatic insects in which both sexes are winged have sufficient dispersal abilities to maintain population connectivity across rivers (Hughes 2007, Hughes et al. 2008).

A study on *D. cephalotes* with isoenzyme markers found patchy gene flow with some gene flow between the populations of the rivers Aniene, Nera, and Velino in Italy (Ketmaier et al. 2001). Both the Nera and Aniene meet the Tiber River below 100 m elevation, so population connectivity by rivers is unlikely. Therefore, it seems plausible that dispersal is accomplished primarily by flying female imagines. Patchiness in population connectivity might be caused by habitat heterogeneity and land use between river catchments (Smith et al. 2009), as could be the case for *D. cephalotes* populations in the Sauerland region. Isotope labeling showed that individuals in another stonefly species flew to adjacent catchments at a distance >500 m (Briers et al. 2004).

Dinocras cephalotes has been described as a poor flyer (Ketmaier et al. 2001), but successful migration of only a few ovigerous females per generation may be sufficient to maintain gene flow. COI haplotypes have a relatively homogeneous distribution across the study area and the haplotype groups show a high diversity (47 different haplotypes), so effective population size probably is high because of recurrent exchanges. Thus *D. cephalotes* cannot be considered as genetically depauperate or endangered in this region. *Dinocras cephalotes* is a predatory stonefly (Bo et al. 2007), so its density in a habitat is controlled by prey availability. If populations were indeed small and isolated, they would quickly lose genetic diversity by genetic drift, which, in turn, would be detected quickly with the markers applied in our study.

Conclusions

Two genetically distinct COI haplotype groups were found for *D. cephalotes*, but patterns of genetic subdivision were not detected in the nuclear markers. Thus, historic isolation of refugial populations has led to prominent mitochondrial lineages but not to cryptic species because populations are interbreeding.

The lack of obvious pairwise differentiation among individual populations suggests that *D. cephalotes* is capable of dispersing between different headwater streams, presumably by flying female imagines. Our data suggest that, within a few generations, *D. cephalotes* should be capable of reaching and recolonizing restored habitats at a regional scale. Therefore, *D. cephalotes* is likely to be a valuable and reliable indicator species for monitoring the success of stream restorations if found in restored freshwater sections from which it was absent prior to restoration. Dispersal of *D. cephalotes* is a valuable case study, but restoration success per se must be estimated based on a wider range of taxa.

ACKNOWLEDGEMENTS

We thank Uwe John (AWI Bremerhaven) for carrying out the 454 sequencing. Andrey Rozenberg and Philipp Brand (Ruhr University Bochum) kindly helped with bioinformatic analyses. We also thank the members of the EvoEco Journal Club and 2 anonymous referees for helpful suggestions that improved this manuscript. This work was supported in part by the *GeneStream* project (<http://GeneStream.de>), which is funded by the Kurt Eberhard Bode Foundation within the Deutsches Stiftungszentrum, and by DFG grant no. HE 2764/2-1.

LITERATURE CITED

- Alp, M., I. Keller, A. M. Westram, and C. T. Robinson. 2012. How river structure and biological traits influence gene flow: a population genetic study of two stream invertebrates with differing dispersal abilities. *Freshwater Biology* 57: 969–981.
- Bálint, M., S. Domisch, C. H. M. Engelhardt, P. Haase, S. Lehrian, J. Sauer, K. Theissing, S. U. Pauls, and C. Nowak. 2011. Cryptic biodiversity loss linked to global climate change. *Nature Climate Change* 1:1–6.
- Ballard, J. W. O., and M. C. Whitlock. 2004. The incomplete natural history of mitochondria. *Molecular Ecology* 13:729–744.
- Bo, T., S. Fenoglio, and G. Malacarne. 2007. Diet of *Dinocras cephalotes* and *Perla marginata* (Plecoptera: Perlidae) in an Apennine stream (northwestern Italy). *Canadian Entomologist* 139:358–364.
- Briers, R. A., H. M. Cariss, and J. H. Gee. 2002. Dispersal of adult stoneflies (Plecoptera) from upland streams draining catchments with contrasting land-use. *Archiv für Hydrobiologie* 155:627–644.
- Briers, R. A., J. H. Gee, H. M. Cariss, and R. Geoghegan. 2004. Inter-population dispersal by adult stoneflies detected by stable isotope enrichment. *Freshwater Biology* 49:425–431.

- Camacho, C., G. Coulouris, V. Avagyan, N. Ma, J. Papadopoulos, K. Bealer, and T. L. Madden. 2009. BLAST+: architecture and applications. *BMC Bioinformatics* 10:421–430.
- Cardinale, B. J. 2011. Biodiversity improves water quality through niche partitioning. *Nature* 472:86–89.
- Chevreaux, B., T. Wetter, and S. Suhai. 1999. Genome sequence assembly using trace signals and additional sequence information. *Computer Science and Biology: Proceedings of the German Conference on Bioinformatics (GCB)* 99:45–56.
- Coutant, C. 1982. Evidence for upstream dispersion of adult caddisflies (Trichoptera: Hydropsychidae) in the Columbia River. *Aquatic Insects* 4:61–66.
- Curtis, J. 1827. *British entomology: being illustrations and descriptions of the genera of insects found in Great Britain and Ireland* 4:189–191.
- Dray, S., and A.-B. Dufour. 2007. The ade4 package: implementing the duality diagram for ecologists. *Journal of Statistical Software* 22:1–20.
- Earl, D. A., and B. M. vonHoldt. 2011. STRUCTURE HARVESTER: a website and program for visualizing STRUCTURE output and implementing the Evanno method. *Conservation Genetics Resources* 4:359–361.
- EEA (European Environment Agency) 2012. European waters — assessment of status and pressures. EEA Report No. 8/2012. European Environment Agency, Copenhagen, Denmark.
- Eiseler, B., and K. Enting. 2012. Verbreitungsatlas der Steinfliegen (Plecoptera) in Nordrhein Westfalen. LANUV-Fachbericht 23. Landesamt für Natur, Umwelt und Verbraucherschutz Nordrhein-Westfalen, Düsseldorf, Germany.
- European Union. 2000. Directive 2000/60/EC of the European Parliament and of the Council of 23 October 2000 establishing a framework for the Community action in the field of water policy. *Official Journal L327*:1–73.
- Evanno, G., S. Regnaut, and J. Goudet. 2005. Detecting the number of clusters of individuals using the software STRUCTURE: a simulation study. *Molecular Ecology* 14:2611–2620.
- Excoffier, L., G. Laval, and S. Schneider. 2005. Arlequin (version 3.0): an integrated software package for population genetics data analysis. *Evolutionary Bioinformatics Online* 1:47–50.
- Falush, D., M. Stephens, and J. K. Pritchard. 2003. Inference of population structure using multilocus genotype data: linked loci and correlated allele frequencies. *Genetics* 164:1567–1587.
- Feld, C. K., S. Birk, D. C. Bradley, D. Hering, J. Kail, A. Marzin, A. Melcher, D. Nemitz, M. L. Pedersen, F. Pletterbauer, D. Pont, P. F. M. Verdonshot, and N. Friberg. 2011. From natural to degraded rivers and back again: a test of restoration ecology theory and practice. *Advances in Ecological Research* 44:119–209.
- Folmer, O., M. Black, W. Hoeh, R. Lutz, and R. Vrijenhoek. 1994. DNA primers for amplification of mitochondrial cytochrome *c* oxidase subunit I from diverse metazoan invertebrates. *Molecular Marine Biology and Biotechnology* 3:294–299.
- Hassall, C., and D. J. Thompson. 2011. Study design and mark-recapture estimates of dispersal: a case study with the endangered damselfly *Coenagrion mercuriale*. *Journal of Insect Conservation* 16:111–120.
- Hebert, P. D. N., M. Y. Stoeckle, T. S. Zemlak, and C. M. Francis. 2004. Identification of birds through DNA barcodes. *PLoS Biology* 2:e312.
- Hellawell, J. M. 1986. *Biological indicators of freshwater pollution and environmental management*. Elsevier Applied Science Publishers, London, UK.
- Hering, D., R. K. Johnson, S. Kramm, S. Schmutz, K. Szoszkiewicz, and P. F. M. Verdonshot. 2006. Assessment of European streams with diatoms, macrophytes, macroinvertebrates and fish: a comparative metric-based analysis of organism response to stress. *Freshwater Biology* 51:1757–1785.
- Hughes, J. M. 2007. Constraints on recovery: using molecular methods to study connectivity of aquatic biota in rivers and streams. *Freshwater Biology* 52:616–631.
- Hughes, J. M., D. J. Schmidt, and D. S. Finn. 2009. Genes in streams: using DNA to understand the movement of freshwater fauna and their riverine habitat. *BioScience* 59:573–583.
- Hughes, J. M., D. J. Schmidt, A. McLean, and A. Wheatley. 2008. Population genetic structure in stream insects: what have we learned? Pages 268–288 *in* J. Lancaster and R. A. Robert (editors). *Aquatic insects: challenges to populations*. CABI International, Wallingford, UK.
- Iannilli, V., R. Fochetti, and J. Tierno de Figueroa. 2002. Life cycle of *Dinocras cephalotes* (Curtis, 1827) in central Italy (Plecoptera, Perlidae). *Boletín de la Sociedad Entomológica Aragonesa* 31:177–179.
- Jähnig, S. C., A. W. Lorenz, and D. Hering. 2009. Restoration effort, habitat mosaics, and macroinvertebrates—does channel form determine community composition? *Aquatic Conservation: Marine and Freshwater Ecosystems* 19:157–169.
- Jombart, T. 2008. adegenet: a R package for the multivariate analysis of genetic markers. *Bioinformatics* 24:1403–1405.
- Jueterbock, A., P. Kraemer, G. Gerlach, and J. Deppermann. 2012. DEMETics: evaluating the genetic differentiation between populations based on G_{ST} and D values. R Project for Statistical Computing, Vienna, Austria. (Available from: <http://cran.r-project.org/web/packages/DEMETics/index.html>)
- Katoh, K., K. Misawa, K.-I. Kuma, and T. Miyata. 2002. MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Research* 30:3059–3066.
- Kearse, M., R. Moir, A. Wilson, S. Stones-Havas, M. Cheung, S. Sturrock, S. Buxton, A. Cooper, S. Markowitz, C. Duran, T. Thierer, B. Ashton, P. Meintjes, and A. Drummond. 2012. Geneious basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* 28:1647–1649.
- Ketmaier, V., R. Fochetti, V. Iannilli, and E. De Matthea. 2001. Patterns of genetic differentiation and gene flow in Central

- Italian populations of *Dinocras cephalotes* (Curtis, 1827) (Insecta, Plecoptera). *Archiv für Hydrobiologie* 150:457–472.
- Kovats, Z., J. Ciborowski, and L. Corkum. 1996. Inland dispersal of adult aquatic insects. *Freshwater Biology* 36:265–276.
- Lake, P. S., N. Bond, and P. Reich. 2007. Linking ecological theory with stream restoration. *Freshwater Biology* 52:597–615.
- Leese, F., P. Brand, A. Rozenberg, C. Mayer, S. Agrawal, J. Dambach, L. Dietz, J. S. Doemel, W. P. Goodall-Copstake, C. Held, J. A. Jackson, K. P. Lampert, K. Linse, J. N. Macher, J. Nolzen, M. J. Raupach, N. T. Rivera, C. D. Schubart, S. Striewski, R. Tollrian, and C. J. Sands. 2012. Exploring Pandora's box: potential and pitfalls of low coverage genome surveys for evolutionary biology. *PLoS ONE* 7:e49202.
- Lehrian, S., M. Balint, P. Haase, and S. U. Pauls. 2010. Genetic population structure of an autumn-emerging caddisfly with inherently low dispersal capacity and insights into its phylogeography. *Journal of the North American Benthological Society* 29:1100–1118.
- Mayer, C. 2006. Phobos 3.3.11. Ruhr Universität Bochum, Bochum, Germany. (Available from: http://www.rub.de/spezzoo/cm/cm_phobos.htm)
- Meirmans, P. G., and P. H. Van Tienderen. 2004. GENOTYPE and GENODIVE: two programs for the analysis of genetic diversity of asexual organisms. *Molecular Ecology Notes* 4:792–794.
- Metcalf, J. L. 1989. Biological water quality assessment of running waters based on macroinvertebrate communities: history and present status in Europe. *Environmental Pollution* 60:101–139.
- Mynott, J. H., J. M. Webb, and P. J. Suter. 2011. Adult and larval associations of the alpine stonefly genus *Riekoperla* McLellan (Plecoptera: Gripopterygidae) using mitochondrial DNA. *Invertebrate Systematics* 25:11–12.
- Palmer, M. A., R. F. Ambrose, and N. L. Poff. 1997. Ecological theory and community restoration ecology. *Restoration Ecology* 5:291–300.
- Palmer, M. A., H. L. Menninger, and E. Bernhardt. 2010. River restoration, habitat heterogeneity and biodiversity: a failure of theory or practice? *Freshwater Biology* 55:205–222.
- Pauls, S. U., W. Graf, P. Haase, H. T. Lumbsch, and J. Waringer. 2008. Grazers, shredders and filtering carnivores—the evolution of feeding ecology in Drusinae (Trichoptera: Limnephilidae): insights from a molecular phylogeny. *Molecular Phylogenetics and Evolution* 46:776–791.
- Pauls, S. U., H. T. Lumbsch, and P. Haase. 2006. Phylogeography of the montane caddisfly *Drusus discolor*: evidence for multiple refugia and periglacial survival. *Molecular Ecology* 15:2153–2169.
- Pfrender, M. E., C. P. Hawkins, M. Bagley, G. W. Courtney, B. R. Creutzburg, J. H. Epler, S. Fend, D. Schindel, L. C. Ferrington, P. L. Hartzell, S. Jackson, D. P. Larsen, A. Lévesque, J. C. Morse, M. J. Petersen, D. Ruitter, and M. Whiting. 2010. Assessing macroinvertebrate biodiversity in freshwater ecosystems: advances and challenges in DNA-based approaches. *Quarterly Review of Biology* 85:319–340.
- Poff, N. L., J. D. Olden, D. Merritt, and D. Pepin. 2007. Homogenization of regional river dynamics by dams and global biodiversity implications. *Proceedings of the National Academy of Sciences of the United States of America* 104:5732–5737.
- Pritchard, J. K., M. Stephens, and P. Donnelly. 2000. Inference of population structure using multilocus genotype data. *Genetics* 155:945–959.
- Rosenberg, D. M., and V. H. Resh. 1993. *Freshwater bio-monitoring and benthic macroinvertebrates*. Chapman and Hall, New York.
- Rozen, S., and H. Skaletsky. 2000. Primer3 on the WWW for general users and for biologist programmers. *Methods in Molecular Biology* 132:365–386.
- Smith, R. F., L. C. Alexander, and W. O. Lamp. 2009. Dispersal by terrestrial stages of stream insects in urban watersheds: a synthesis of current knowledge. *Journal of the North American Benthological Society* 28:1022–1037.
- Stettmer, C. 1996. Colonisation and dispersal patterns of banded (*Calopteryx splendens*) and beautiful demoiselles (*C. virgo*) (Odonata: Calopterygidae) in south-east German streams. *European Journal of Entomology* 93:579–593.
- Sunnucks, P., and D. F. Hales. 1996. Numerous transposed sequences of mitochondrial cytochrome oxidase I–II in aphids of the genus *Sitobion* (Hemiptera: Aphididae). *Molecular Biology and Evolution* 13:510–524.
- Sweeney, B. W., J. M. Battle, J. K. Jackson, and T. Dapkey. 2011. Can DNA barcodes of stream macroinvertebrates improve descriptions of community structure and water quality? *Journal of the North American Benthological Society* 30:195–216.
- Tamura, K., D. Peterson, N. Peterson, G. Stecher, M. Nei, and S. Kumar. 2011. MEGA5: Molecular Evolutionary Genetics Analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Molecular Biology and Evolution* 28:2731–2739.
- Teacher, A. G. F., and D. J. Griffiths. 2010. HapStar: automated haplotype network layout and visualization. *Molecular Ecology Resources* 11:151–153.
- Theissing, K., M. Balint, K. A. Feldheim, P. Haase, J. Johannesen, I. Laube, and S. U. Pauls. 2012. Glacial survival and post-glacial recolonization of an arctic–alpine freshwater insect (*Arcynopteryx dichroa*, Plecoptera, Perlodidae) in Europe. *Journal of Biogeography* 40:236–248.
- Tierno de Figueroa, J. M., J. M. Luzón-Ortega, and M. J. López-Rodríguez. 2006. Mating balls in stoneflies (Insecta, Plecoptera) = Las bolas de apareamiento en plecópteros (Insecta, Plecoptera). *Zoologica Baetica* 17:93–96.
- Van Oosterhout, C., W. F. Hutchinson, D. P. M. Wills, and P. Shipley. 2004. Micro-checker: software for identifying and correcting genotyping errors in microsatellite data. *Molecular Ecology Notes* 4:535–538.

- Vaughn, C. C. 2010. Biodiversity losses and ecosystem function in freshwaters: emerging conclusions and research directions. *BioScience* 60:25–35.
- Vörösmarty, C. J., P. B. McIntyre, M. O. Gessner, D. Dudgeon, A. Prusevich, P. Green, S. Glidden, S. E. Bunn, C. A. Sullivan, C. R. Liermann, and P. M. Davies. 2010. Global threats to human water security and river biodiversity. *Nature* 467:555–561.
- Werle, E., C. Schneider, M. Renner, M. Völker, and W. Fiehn. 1994. Convenient single-step, one tube purification of PCR products for direct sequencing. *Nucleic Acids Research* 22:4354.
- Zhou, X., S. J. Adamowicz, L. M. Jacobus, R. E. DeWalt, and P. D. N. Hebert. 2009. Toward a comprehensive barcode library for arctic life - Ephemeroptera, Plecoptera, and Trichoptera of Churchill, Manitoba, Canada. *Frontiers in Zoology* 6:30.
- Zhou, X., L. M. Jacobus, R. E. DeWalt, S. J. Adamowicz, and P. D. N. Hebert. 2010. Ephemeroptera, Plecoptera, and Trichoptera fauna of Churchill (Manitoba, Canada): insights into biodiversity patterns from DNA barcoding. *Journal of the North American Benthological Society* 29:814–837.