



Decline of Raptors over a Three-Year Period in Laikipia, Central Kenya

Authors: Ogada, Darcy L., and Keesing, Felicia

Source: Journal of Raptor Research, 44(2) : 129-135

Published By: Raptor Research Foundation

URL: <https://doi.org/10.3356/JRR-09-49.1>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

DECLINE OF RAPTORS OVER A THREE-YEAR PERIOD IN LAIKIPIA, CENTRAL KENYA

DARCY L. OGADA¹

Mpala Research Centre, P.O. Box 555, Nanyuki, Kenya

FELICIA KEESING

Bard College, Box 5000, Annandale, NY 12504 U.S.A.

ABSTRACT.—Raptors were monitored monthly over a three-year period in a protected area in central Kenya. The number of raptors declined more than 40% per year. Scavenging birds accounted for most of the decline; sightings decreased by 70% during our surveys, although these declines were not statistically significant. During the time of the study, the overall populations of large wild herbivores showed little change, whereas domestic herbivores, particularly sheep and goats, increased markedly, suggesting that food limitation was not the cause of the vulture declines at the study site. Possible causes of raptor decline include the consumption of poisoned baits, which are placed by pastoralists to kill large predators that attack livestock. Scavenging birds provide one of the most important yet underappreciated ecosystem services of any avian group. The rapid decline of scavenging birds, especially vultures, in central Kenya warrants additional population monitoring to understand whether declines are local or regional, and to elucidate causes of population decreases.

KEY WORDS: *Africa, decline, Furadan, livestock, poisoning, raptors, vulture.*

DISMINUCIÓN DE AVES RAPACES DURANTE UN PERIODO DE TRES AÑOS EN LAIKIPIA, KENIA CENTRAL

RESUMEN.—Las aves rapaces fueron monitoreadas mensualmente por un periodo de tres años en un área protegida en el centro de Kenia. El número de aves rapaces disminuyó en más de un 40% por año. Las aves carroñeras fueron el grupo que dio cuenta de la mayor parte de esta disminución ya que sus avistamientos disminuyeron en un 70% durante nuestros muestreos, pero estas disminuciones no fueron estadísticamente significativas. Durante el periodo de nuestro estudio, la población de grandes herbívoros silvestres no varió mucho, mientras que la población de herbívoros domésticos, principalmente ovejas y cabras, aumentó considerablemente, lo que sugiere que la limitación de alimento no fue la causa de la disminución de los buitres en el sitio de estudio. Las posibles causas de la disminución de aves rapaces incluyen el consumo de cebos envenenados que son colocados por los pastores locales para matar depredadores de gran tamaño que atacan a su ganado. Entre todos los grupos de aves, las carroñeras proveen uno de los servicios ecosistémicos más importantes y más subestimados. La rápida disminución de aves carroñeras en Kenia, especialmente de los buitres, justifica monitoreos poblacionales adicionales para entender si estas disminuciones son locales o regionales y para elucidar sus posibles causas.

[Traducción del equipo editorial]

National parks and other protected areas are playing an increasingly important role in the conservation of raptors (Liversidge 1984, Thiollay 2006). Not surprisingly, protected areas are critical refuges for most raptor species and they harbor the bulk of large raptor populations in Africa (Sorley and An-

dersen 1994, Herremans and Herremans-Tonnoeyr 2000, Thiollay 2006). Outside protected areas, raptors face increased threats associated with human activities including deforestation, overgrazing, increase in pesticide use, poisoning of predators, overhunting, and direct persecution (Thiollay 2006).

Raptors living within protected areas are assumed to be relatively free from these disturbances. However, substantial increases in human population have occurred in recent decades and protected areas are

¹ Present address: National Museums of Kenya, Ornithology Section, P.O. Box 40658, 00100 Nairobi, Kenya; email address: darcycogada@yahoo.com




Figure 1. Location of the Mpala Research Centre within Laikipia District in central Kenya.

not immune to the pressures this has created on limited natural resources. A recent study showed that raptor numbers within 30 km of protected area boundaries were more than 40% lower than in the core, indicating a large edge effect on populations inside protected areas (Herremans and Herremans-Tonnoeyr 2000). In addition, some raptors, most notably vultures and large eagles, travel large distances in search of food and require extensive territories for breeding, making them more likely to range outside of protected areas to locations where they may become vulnerable to harmful human activities (Houston 1974, Pennycuik 1976, Herholdt et al. 1996).

The core of central Kenya's Laikipia District functions largely as a protected area despite being divided and privately owned. Almost half of the district's 966 600 ha consists of large-scale ranches and conservancies that are hospitable to wildlife and enable large-ranging birds like vultures and eagles to exist relatively free from habitat destruction and human interference. However, areas mostly outside of the core have seen considerable recent settlement by small-scale crop farmers and traditionally nomadic

pastoralists. Approximately 35% of the district consists of settlements or small-scale urbanization (Frank et al. 2005).

To determine whether the abundance of raptors in central Kenya was changing, we monitored raptors every month for three years (2001–2003) in central Laikipia District.

METHODS

Study Site. The study was conducted from January 2001–December 2003 at Mpala Research Centre (MRC), located in central Laikipia District in the highlands of central Kenya (Fig. 1). The habitat consists of wooded savanna, and the dominant tree species is *Acacia drepanolobium*. Large wildlife species include elephants (*Loxodonta africana*), reticulated giraffes (*Giraffa camelopardalis*), Grevy's zebras (*Equus grevyi*), elands (*Taurotragus oryx*), cape buffalos (*Syncerus caffer*), lions (*Panthera leo*), leopards (*P. pardus*), cheetahs (*Acinonyx jubatus*) and spotted hyenas (*Crocuta crocuta*).

The more common avian fauna in the study area comprises approximately 75 species (Ogada et al.

2008), of which nine are seasonal migrants. Common raptors include Rueppell's Vulture (*Cyps rueppellii*) and African White-backed Vulture (*G. africanus*), Black-winged Kites (*Elanus caeruleus*) and Eurasian Kestrels (*Falco tinnunculus*), which are seasonal migrants.

Study Design. This research was conducted within a long-term herbivore exclusion experiment (Kenya Long-term Exclusion Experiment, or KLEE; Young et al. 1998) that was established at MRC in 1995. KLEE consists of a randomized block design, with three replicates of six treatments. The six treatments selectively exclude different combinations of wild and domestic herbivore species and each treatment is 200 m × 200 m, or 4 ha. The total area surveyed was 1.12 km².

Raptor Surveys. We sampled raptors as part of a larger experiment that examined the effects of herbivores on small mammals (Keesing 2000). We conducted surveys from within six of the eighteen plots every month—three that excluded all large herbivores and three that allowed access to all large herbivores (Fig. 1). We sampled the same plots every month. We did not survey during November 2001, February 2002, and November 2003 due to excessive rainfall or other logistical complications, so we excluded the months of November and February from our analyses for all three years. Each survey was conducted between 10:00–12:30 H and was completed within three days. Surveys were conducted in late morning in order to include large vultures, which typically do not begin soaring until thermals are produced by solar heat (Brown 1971).

During surveys, observers walked the perimeter of each 200 m × 200 m plot, then walked diagonals through the center of each plot, scanning the horizon, trees, and ground for any perched, ground-roosting, walking, or flying raptors. Total walking time was approximately 30 min per plot. Plots were walked to ensure that all ground-roosting raptors (e.g., owls and harriers) were counted, because the grass height could reach 0.5 m in plots that excluded all large herbivores. We recorded all raptors observed using binoculars, regardless of whether the birds were inside or outside the plots. Raptors were identified to species level, when possible. Given the difficulty of identifying raptors, especially those flying at a distance, many were recorded only by group (i.e., vulture, falcon).

Large Herbivore Surveys. Estimates of large herbivore abundances were from two aerial censuses conducted January–March 2001 and February 2003

by the Laikipia Wildlife Forum and provide a sample count of large herbivores species in Laikipia District including the study area (Georgiadis and Ojwang 2001, Georgiadis et al. 2003).

Species were grouped for analyses. "Scavenging birds" included Rueppell's Vultures, African White-backed Vultures, Lappet-faced Vultures (*Torgos tracheliotus*), Bateleurs (*Terathopius ecaudatus*), and Tawny Eagles (*Aquila rapax*). "Non-scavenging eagles" were Wahlberg's Eagles (*A. wahlbergi*), Black-chested Snake-Eagles (*Circaetus pectoralis*), African Hawk-Eagles (*Hieraaetus spilogaster*), and Martial Eagles (*Polemaetus bellicosus*). "Hawks" included Augur Buzzards (*Buteo augur*), Eurasian Buzzards (*Buteo buteo*), and Eastern Chanting-Goshawk (*Melierax poliopterus*). We omitted harrier sightings, which included three unknown harriers (*Circus* sp.) and two Pallid Harriers (*C. macrourus*) because their numbers were so few. "Falcons" were Eurasian Kestrels and Pygmy Falcons (*Polihierax semitorquatus*). The only kite was the Black-winged Kite. We included "unknown" and harrier sightings for estimates of raptor abundance by year, but not for group-level analyses.

Because of their large ranges, raptors observed during the surveys of each KLEE treatment were not necessarily independent sightings. Therefore, we pooled data from all blocks and treatments. We conducted a correlation analysis between year (2001–2003) and the mean number of monthly sightings of all raptors for the first analysis, and of just scavenging birds for the second. We also performed a linear regression using the number of scavenging birds as the dependent variable and rainfall as the independent variable (rainfall was measured daily using a standard rainfall gauge as part of a weather station based at Mpala Research Centre).

RESULTS

Raptor Surveys. From January 2001–December 2003 we had 535 raptor sightings comprising 16 species, three of which were palearctic migrants. Of the species that were identified, the most common were *Cyps* vultures (69%) followed by Black-winged Kites (5%) and Tawny Eagles (4%).

The mean number of raptors observed each month declined significantly over time ($P = 0.05$). Raptors declined 40% in the first year and 48% in the second, for an overall decline of 68% between 2001 and 2003. Scavenging birds accounted for most of the decline, although the correlation be-




Figure 2. Annual monthly mean (+SE) number of raptor sightings during 2001–03, by raptor group. Species in each group are described in the text.

tween year and mean monthly sightings of scavenging birds was not significant ($P = 0.40$; Fig. 2). Scavenging birds declined approximately 50% per year and a total of 70% over the study period. Sightings for other raptor groups were too infrequent to permit analysis. The number of scavenging birds was not related to rainfall ($P = 0.28$).

DISCUSSION

In our surveys, raptors declined by 68% over three years. Scavenging birds were primarily responsible for this decline (Fig. 2). Scavenging birds, es-

pecially vultures, are declining worldwide due to poisoning, persecution, food shortages, and land-use changes (Pain et al. 2003, Thiollay 2006). Precipitous declines have occurred in West Africa (Thiollay 2006) and recent catastrophic population crashes have garnered attention in South Asia (Oaks et al. 2004, Prakash et al. 2003, Green et al. 2004). Our results suggest that this pattern is occurring in East Africa as well.

Because we sampled only a small area, the declines we observed may have resulted from a change in space use by scavenging birds rather than a re-

gion-wide population decline. Dispersion in subtropical African raptors is usually related to rainfall (Brown 1971, Herremans and Herremans-Tonnoeyr 2000). Rainfall at the study site during the period of our observations was not unusual in magnitude or distribution, and we detected no significant effect of rainfall on scavenging bird abundance. However, we cannot rule out the possibility that scavenging birds in our study area relocated to other sites. Nevertheless, abundance at our site declined significantly, whether these declines represent local or region-wide declines.

One hypothesis to explain the rapid decline we observed at the study site is a lack of available food. Large vultures in particular feed mainly on carrion of large- and medium-sized mammals and are obligate scavengers (Ruxton and Houston 2004). However, during the study period, wild herbivores in Laikipia District increased by 9% and domestic herbivores, especially sheep and goats, increased by 37% (Georgiadis and Ojwang 2001, Georgiadis et al. 2003). These data suggest that low food availability was not the primary cause of the declines in scavenging birds.

Another potential explanation for raptor decline is mortality from infectious disease. Although this hypothesis could not be ruled out, it seemed unlikely in our study region, as there were no reports of scavenging birds, e.g., vultures, dying of natural causes at a number of colonies where monitoring is ongoing (M. Virani pers. comm.). Infectious disease does not appear to be a major cause of mortality in *Gyps* vulture populations in southern Africa (Benson 2000).

A third hypothesis is that scavenging birds were poisoned at carcasses. Vultures are particularly vulnerable to poisoning, because they can rapidly congregate in large numbers at a carcass. Thus, one poisoned carcass can affect a large number of birds (Newton 1979). Indeed, reports of mass vulture poisonings within Africa are numerous (Borello 1985, Allen 1989, Anderson 1994, Simmons 1995). Two groups of neurotoxic pesticides, the organophosphorous and carbamate compounds, tend to be very acutely toxic to birds and cases of mortality are frequently reported (e.g., Mineau et al. 1999, Mineau 2005). Because of their acute toxicity, carbofurans are frequently implicated in abuse cases where raptors or other vertebrates are targeted directly (Mineau et al. 1999). Carbofuran, sold under the trade name Furadan, is an agricultural pesticide that is legally sold throughout Kenya, and its use as a cheap, effective poison is well known among

farming communities in Kenya (Ogada and Kibuthu 2008). In Laikipia District soil, water and plant samples have shown high environmental contamination with concentrations of carbofuran and its two toxic metabolites (Otieno et al. 2010). Recent studies have implicated carbofuran in the mortality of African White-backed Vultures near our study site (Otieno et al. in press a, Otieno et al. in press b). These reports also confirmed widespread involvement of carbofuran in vulture poisoning incidents throughout Kenya (Mijeje 2009, Otieno et al. in press a, Otieno et al. in press b). Vultures, and perhaps other scavenging birds, may have become unintended victims of poisoning of predators that occurs following livestock depredation events (Otieno 2009). Whether poisoning explains the rapid decline in abundance of scavenging birds at our site is currently under investigation.

Scavenging birds provide some of the most important yet underappreciated ecosystem services of any avian group (Houston and Cooper 1975, DeVault et al. 2003, Sekercioglu 2006). Yet among birds, nearly 40% of vultures are extinction-prone, more than any other functional group (Sekercioglu et al. 2004). All the vulture species observed during our study are on the IUCN Red List of Threatened Species (IUCN 2009). The apparent decline of scavenging birds that we observed in central Kenya warrants population monitoring to understand whether declines are local or regional and to elucidate causes of population decreases.

ACKNOWLEDGMENTS

We gratefully acknowledge the support and cooperation of the Ministry of Education of the Republic of Kenya and Kenyatta University. This research was supported by a National Science Foundation grant (DEB CAREER 0196177) to FK. The KLEE plots were built and maintained with grants from the James Smithson Fund of the Smithsonian Institution (to Alan Smith), the National Geographic Society (4691-91), the National Science Foundation (BSR-97-07477 and BSR-03-16402), and the African Elephant Program of the U.S. Fish and Wildlife Service (98210-0G563; to T.P. Young). Logistical support was provided by Mpala Research Centre, Truman Young, Mordecai Ogada, Kerry Outram, Nick Georgiadis, Heather Wallington, and Bard College. In 2002, surveys were led by Brian Allan. Douglas McCauley led some field surveys in 2003. Field assistance was provided by Abdikadir Ali Hassan, James Ponoto, and Kadir Age. Discussions with the Bird Committee of Nature Kenya, Paula Kahumbu, Laurence Frank, Martin Odino, Munir Virani, Simon Thomsett, and Joseph Lalah were instrumental in the preparation of this manuscript. The manuscript was improved by comments from Simon Thomsett and two anonymous reviewers.

LITERATURE CITED

- ALLEN, D.G. 1989. Strychnine poison and the conservation of avian scavengers in the Karoo, South Africa. *South African Journal of Wildlife Research* 19:102–106.
- ANDERSON, M.D. 1994. Mass African Whitebacked Vulture poisoning in the northern Cape. *Vulture News* 29:31–32.
- BENSON, P.C. 2000. Causes of Cape Vulture *Gyps coprotheres* mortality at the Kransberg colony: a 17-year update. Pages 77–86 in R.D. Chancellor and B.-U. Meyburg [EDS.], Raptors at risk; proceedings of the V World Conference on Birds of Prey and Owls. Hancock House Publishers, Blaine, WA U.S.A. and World Working Group on Birds of Prey and Owls, Berlin, Germany.
- BORELLO, W.D. 1985. Poisoned vultures in Botswana: known facts. *Babbler* 9:22–23.
- BROWN, L. 1971. African birds of prey. Houghton Mifflin Company, Boston, MA U.S.A.
- DEVAULT, T.L., O.E. RHODES, AND J.A. SHIVIK. 2003. Scavenging by vertebrates: behavioural, ecological, and evolutionary perspectives on an important energy transfer pathway in terrestrial ecosystems. *Oikos* 102:225–234.
- FRANK, L.G., R. WOODROFFE, AND M.O. OGADA. 2005. People and predators in Laikipia District, Kenya. Pages 286–304 in R. Woodroffe, S. Thirgood, and A. Rabinowitz [EDS.], People and wildlife: conflict or coexistence? Cambridge University Press, Cambridge, U.K.
- GEORGIADIS, N. AND G. OJWANG. 2001. Numbers and distributions of large herbivores in Laikipia, Samburu, and Isiolo districts sample count, January 22 to March 7, 2001 for The Laikipia Wildlife Forum. Laikipia Wildlife Forum Newsletter, Nov. 2001. Nanyuki, Kenya.
- , N. OLWERO, AND G. OJWANG. 2003. Numbers and distributions of large herbivores in Laikipia District, Leroghi, and Lewa Conservancy. Mpala Research Centre, Nanyuki, Kenya.
- GREEN, R.E., I. NEWTON, S. SHULTZ, A.A. CUNNINGHAM, M. GILBERT, D.J. PAIN, AND V. PRAKASH. 2004. Diclofenac poisoning as a cause of vulture population declines across the Indian subcontinent. *Journal of Applied Ecology* 41:793–800.
- HERHOLDT, J.J., A.C. KEMP, AND D. DU PLESSIS. 1996. Aspects of the breeding status and ecology of the Bateleur and Tawny Eagle in the Kalahari Gemsbok National Park, South Africa. *Ostrich* 67:126–137.
- HERREMANS, M. AND D. HERREMANS-TONNOEYR. 2000. Land use and the conservation status of raptors in Botswana. *Biological Conservation* 94:31–41.
- HOUSTON, D.C. 1974. Food searching behaviour in Griffon Vultures. *East African Wildlife Journal* 12:63–77.
- AND J.E. COOPER. 1975. The digestive tract of the Whiteback Griffon Vulture and its role in disease transmission among wild ungulates. *Journal of Wildlife Diseases* 11:306–313.
- IUCN. 2009. 2009 IUCN Red List of Threatened Species. Version 2009.1. www.iucnredlist.org. (last accessed on 09 February 2009).
- KEESING, F. 2000. Cryptic consumers and the ecology of an African savanna. *BioScience* 50:205–215.
- LIVERSIDGE, R. 1984. The importance of national parks for raptor survival. Pages 589–600 in J.A. Ledger [ED.], Proceedings of the Fifth Pan-African Ornithological Congress. Southern African Ornithological Society, Johannesburg, South Africa.
- MIJELE, D. 2009. Incidences of poisoning of vultures and lions in the Masai Mara National Reserve. Kenya Wildlife Service Masai Mara Veterinary Report. Nairobi, Kenya.
- MINEAU, P. 2005. Direct losses of birds to pesticides—beginnings of a quantification. Pages 1066–1070 in C.J. Ralph and T.D. Rich [EDS.], Bird conservation implementation and integration in the Americas: proceedings of the Third International Partners in Flight Conference 2002. USDA Forest Service, General Technical Report PSW-GTR-191, Albany, CA U.S.A.
- , M.R. FLETCHER, L.C. GLASER, N.J. THOMAS, C. BRASARD, L.K. WILSON, J.E. ELLIOTT, L.A. LYON, C.J. HENNY, T. BOLLINGER, AND S.L. PORTER. 1999. Poisoning of raptors with organophosphorus and carbamate pesticides with emphasis on Canada, the United States and the United Kingdom. *Journal of Raptor Research* 33:1–37.
- NEWTON, I. 1979. Population ecology of raptors. T. and A.D. Poyser, London, U.K.
- OAKS, J.L., M. GILBERT, M.Z. VIRANI, R.T. WATSON, C.U. METEYER, B.A. RIDEOUT, H.L. SHIVAPRASAD, S. AHMED, M.J.I. CHAUDHRY, M. ARSHAD, S. MAHMOOD, A. ALI, AND A.A. KHAN. 2004. Diclofenac residues as the cause of vulture population decline in Pakistan. *Nature* 427:630–633.
- OGADA, D.L. AND P.M. KIBUTHU. 2008. Conserving Mackinder's Eagle Owls in farmlands of Kenya: assessing the influence of pesticide use, tourism and local knowledge of owl habits in protecting a culturally loathed species. *Environmental Conservation* 35:252–260.
- , M.E. GADD, R.S. OSTFELD, T.P. YOUNG, AND F. KEESING. 2008. Impacts of large herbivorous mammals on bird diversity and abundance in an African savanna. *Oecologia* 156:387–397.
- OTIENO, P.O. 2009. Monitoring carbofuran residues in Laikipia and Isiolo districts in Kenya for ecological risk assessment. M.S. thesis, Department of Chemistry, Maseno University, Maseno, Kenya.
- , J.O. LALAH, M. VIRANI, I.O. JONDIKO, AND K.W. SCHRAMM. 2010. Soil and water contamination with carbofuran residues in agricultural farmlands in Kenya following Furadan application. *Journal of Environmental Science and Health Part B* 45:137–144.
- , ———, ———, ———, AND ———. In press a. Carbofuran and its toxic metabolites provide forensic evidence for Furadan exposure in vultures (*Gyps africanus*) in Kenya. *Bulletin of Environmental Contamination and Toxicology*.

- , ———, ———, ———, AND ———. In press b. Assessment of carbofuran residues for evidence of Furan poisoning of the African White-backed Vulture (*Gyps africanus*) in Kenya. *Bulletin of Environmental Contamination and Toxicology*.
- PAIN, D.J., A.A. CUNNINGHAM, P.F. DONALD, J.W. DUCKWORTH, D.C. HOUSTON, T. KATZNER, J. PARRY-JONES, C. POOLE, V. PRAKASH, P. ROUND, AND R. TIMMINS. 2003. Causes and effects of temporospatial declines of *Gyps* vultures in Asia. *Conservation Biology* 17:661–671.
- PRAKASH, V., D.J. PAIN, A.A. CUNNINGHAM, P.F. DONALD, N. PRAKASH, A. VERMA, R. GARGI, S. SIVAKUMAR, AND A.R. RAHMANI. 2003. Catastrophic collapse of Indian White-backed *Gyps bengalensis* and Long-billed *Gyps indicus* vulture populations. *Biological Conservation* 109:381–390.
- PENNYCUICK, C.J. 1976. Breeding of the Lappet-faced and White-headed vultures (*Torgos tracheliotus* Forster and *Trigonoceps occipitalis* Burchell) on the Serengeti Plains, Tanzania. *East African Wildlife Journal* 14:67–84.
- RUXTON, G.D. AND D.C. HOUSTON. 2004. Obligate vertebrate scavengers must be large soaring fliers. *Journal of Theoretical Biology* 228:431–436.
- SEKERCIOGLU, C.H. 2006. Increasing awareness of avian ecological function. *Trends in Ecology and Evolution* 21:464–471.
- , G.C. DAILY, AND P.R. EHRLICH. 2004. Ecosystem consequences of bird declines. *Proceedings of the National Academy of Sciences of the United States of America* 101:18042–18047.
- SIMMONS, R.E. 1995. Mass poisoning of Lappet-faced Vultures in the Namib Desert. *Journal of African Raptor Biology* 10:3.
- SORLEY, C.S. AND D.E. ANDERSEN. 1994. Raptor abundance in south-central Kenya in relation to land-use patterns. *African Journal of Ecology* 32:30–38.
- THIOLLAY, J.M. 2006. The decline of raptors in West Africa: long-term assessment and the role of protected areas. *Ibis* 148:240–254.
- YOUNG, T.P., B.D. OKELLO, D. KINYUA, AND T.M. PALMER. 1998. KLEE: a long-term multi-species herbivore exclusion experiment in Laikipia, Kenya. *African Journal of Range Forage Science* 14:94–102.

Received 24 June 2009; accepted 15 January 2010