

Spring Revival in the Mid-Hills of the Himalaya: A Socioeconomic Assessment Using Benefit–Cost Analysis

Authors: Butte, Giacomo, Khadka, Karishma, Aldinucci, Alice, Macaulay, Bobby, and Bhuchar, Sanjeev

Source: Mountain Research and Development, 44(4)

Published By: International Mountain Society

URL: <https://doi.org/10.1659/mrd.2024.00020>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Spring Revival in the Mid-Hills of the Himalaya: A Socioeconomic Assessment Using Benefit–Cost Analysis

Giacomo Butte^{1*}, Karishma Khadka^{2,3}, Alice Aldinucci⁴, Bobby Macaulay¹, and Sanjeev Bhuchar²

* Corresponding author: giacomo.butte.2@gmail.com

¹ Centre for Mountain Studies, University of Highlands and Islands (UHI), Crieff Road, Perth PH1 2NX, UK

² International Centre for Integrated Mountain Development (ICIMOD), GPO Box 3226, Kathmandu, Nepal

³ Center for Applied Earth Science and Engineering Research (CAESER), Wilder Tower, Suite 900, 3675 Alumni Drive, Memphis, TN 38152, USA

⁴ Independent researcher

© 2024 Butte et al. This open access article is licensed under a Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>). Please credit the authors and the full source.



Natural springs are the main water source for more than 100 million people living in the Hindu Kush Himalaya. In Nepal, 10 million people in the mid-hills and mountains depend on them. These sources are under stress

due to factors such as climate change, infrastructural development, and socioeconomic changes. To combat this trend, spring revival activities have been carried out across the Hindu Kush Himalaya in the last few years. Considering 2 study sites in Nepal, this work attempts the first benefit–cost analysis for spring

revival in rural settings. First, using literature and community consultations, a cause–effect map was drafted. Second, the benefits and costs were estimated quantitatively using 4 scenarios: 2 based on the study sites and 2 on more generic situations. Positive (>1) benefit–cost ratios were found in 3 scenarios, showing that spring revival has potential but local conditions (households served, presence of other water sources, usage) are important factors to be considered.

Keywords: cost–benefit analysis; Nepal; spring revival; water; Hindu Kush Himalaya (HKH); environmental economics; resilience.

Received: 7 May 2024 **Accepted:** 4 November 2024

Introduction

The hydrological system within the Himalayas is facing considerable strain for a variety of reasons. On the one hand, population increase, increased reliability of water supply (Chen et al 2023), and lifestyle and economic changes (Bharti et al 2020; Faulon and Sacareau 2020) are augmenting water demand. On the other, scarce governance capacity (Ojha et al 2020), urbanization and development (Wester et al 2019; Daniel et al 2021), and climate change are contributing to a decline in annual precipitation, glacier volume (Shea et al 2015), runoff (Nie et al 2021), and an increase in extreme weather events, such as floods and glacial lake outburst floods (Akhtar et al 2008; Shrestha and Aryal 2011; Nie et al 2021). Until recently, the main focus of studies on mountain water availability was on glaciers and snowpacks, mostly from a climate and hydrological perspective (Maurya et al 2011; Bajracharya et al 2015; Kumar 2020). Nonglacial contributions, such as groundwater, have been ignored (Andermann et al 2012; Bookhagen 2012; Verma and Jamwal 2022), but research has shown their significant role in water basins (Williams et al 2016; Wilson et al 2016; Wood et al 2020; Yao et al 2021).

The case of Nepal well represents all these challenges. The country is one of the richest in terms of water but ranked among the lowest in terms of drinking water supply (Thakur et al 2017; Gurung, Adhikari, Chauhan, Thakuri,

Nakarmi, Ghale, et al 2019; Komatsu et al 2020; Joseph and Shrestha 2022). Mid-hill areas, home to 43% of the national population, are of particular interest (Wester et al 2019) because of their unique characteristics. Indeed, unlike higher-elevation areas that depend on glacier-fed water sources to meet their water needs, these regions primarily rely on groundwater, predominantly sourced from water springs (Agarwal et al 2012; Tambe et al 2012; NITI Aayog 2018).

In recent years, several studies have focused on different aspects of water springs, such as defining conceptual models for mountain water aquifers (Tambe et al 2020), spring hydrological classification (Daniel et al 2021), methods of investigation (Hersch 1993; Smakhtin 2001; Tarafdar 2013; Chinnasamy and Prathapar 2016), understanding discharge trends (Agarwal et al 2012; Chapagain et al 2019; Gurung, Adhikari, Chauhan, Thakuri, Nakarmi, Ghale, et al 2019; Pandit et al 2019; Adhikari et al 2021; Dhakal 2021), causes of spring degradation (Poudel and Duex 2017; Poudel 2020), water quality assessment (Tiwari 2000; Gurung, Adhikari, Chauhan, Thakuri, Nakarmi, Rijal, et al 2019; Bhat and Pandit 2020; Khadka and Rijal 2020; Thapa, Pant, et al 2020), and spring distribution modeling (Al-Manmi and Saleh 2019; Ghimire et al 2019).

Efforts have been made to understand the state of these springs, with estimates suggesting that more than half of the 3 million perennial springs in the Indian Himalayas (NITI

TABLE 1 Summary of study site characteristics at the 2 revived springs in Namobuddha municipality.

Characteristic	Bhagwate Pakhako Kuwa	Patalako Dhara
Location	27°34'08"N; 85°37'47"W	27°34'17"N; 85°37'28"W
Spring type	Depression	Depression
Elevation (masl)	1235	1415
Ethnicity of main users	Lama	Dalit
Recharge area type	Community forest	Private fields and community forest
Ecosystem area	2 hectares (upstream and downstream)	1 hectare (up and downstream)
Water flow increase	from 0 to 0.74 lpm	from 2.84 to 3.37 lpm
Users as primary source	1 household	21 households
Users as secondary source	24 households	14 households
Fetches water	Used for irrigation or as emergency source	Currently piped to tank
Intervention	5 ponds, 16 trenches, 4 check dams, 8 palisades	9 trenches, 1 pond

Aayog 2018) have either dried up or become seasonal (Valdiya and Bartarya 1991; Tiwari 2000; Kumar and Sen 2018; Thapa et al 2023). The trend of declining spring discharge is also observed in the Nepali mid-hill region, where over 30% of monitored springs show a decrease in discharge (Chapagain et al 2019). Different factors contribute to this decline in both water quantity and quality. These include higher temperatures (Pandey et al 2017), changing precipitation patterns (Agarwal et al 2012; Macchi et al 2015), land use and soil erosion (Rautela 2015; Pandey et al 2017), changes in forest types (Ghimire et al 2014; Naudiyal and Schmerbeck 2017), infrastructure (Huber and Joshi 2015; Mukherji et al 2018), and sporadic events, such as earthquakes (Lamichhane et al 2020).

Alongside the hydrological and environmental perspectives, researching water management is crucial from a social science perspective. It is closely connected to key aspects of social inequalities related to caste, gender, and power dynamics, often overlooked in water security initiatives (Leder et al 2017; Shrestha and Clement 2019). In the context of rural domestic water management, women bear significant responsibilities, but they often lack equal representation in decision-making processes (Wali et al 2020). This disparity is rooted in factors such as male-dominated representation on decision boards and deeply ingrained historical and societal norms (Raut 2023). The biophysical perspective has thus evolved, first, by incorporating management considerations and subsequently by integrating socioeconomic dimensions. Currently, approaches such as sociohydrology (Sivapalan et al 2014; Blair and Buytaert 2016; Nüsser 2017; Herrera-Franco et al 2021) conceptualize human–water systems as interconnected entities, recognizing the interactions between society and nature as essential for accurately modeling water systems (Di Baldassarre et al 2015; Gober and Wheeler 2015; Troy et al 2015).

Revival projects are being implemented across the Hindu Kush Himalaya (HKH) with varying levels of community involvement. Institutions such as ICIMOD (Shrestha et al 2018) and Aayog–International Water

Management Institute (IWMI) (Rathod et al 2021) have provided practical guidelines, yet significant gaps persist. These include the need for better documentation of spring revival projects, deeper understanding of spring mechanisms, and more robust assessments of the effectiveness and benefit–cost ratios of techniques, as well as improved integration of physical sciences with governance and social dimensions (Kumar et al 2023). This study addresses these gaps by offering, to the authors' knowledge, the first known benefit–cost analysis (BCA) of spring revival activities in the mid-hills to inform regulatory efforts.

Methods

Study sites

The 2 study sites consist of the water springs Bhagwate Pakhako Kuwa (BHA; 27°34'08"N; 85°37'47"W) and Patalako Dhara (PAT; 27°34'17"N; 85°37'28"W) and their surroundings. Both are located in Namobuddha municipality (Table 1; Figure 1), 40 km from the capital (GGGI 2018). Despite being situated in a water-rich district, the municipality has recently faced water depletion (Khatri et al 2021). The examined project, a collaboration between ICIMOD and Agricultural Policy, Research and Extension Development (CEAPRED), focused on marginalized groups and aimed to address the drying of springs through the initiation of revival activities (Dhakal 2021; Khadka et al 2021). The project was part of ICIMOD's broader Resilient Mountain Solutions pilot initiative, which emphasizes gender-inclusive, socially resilient solutions (Pokhrel et al 2019). The project employed a “water tower” approach (Tambe et al 2020), which considers the mountain aquifer as the primary unit of planning, rather than individual springsheds. This emphasis is justified by the aquifers' key role in linking precipitation and spring discharge within the mountain water system. The process followed ICIMOD's protocol (Shrestha et al 2018) which outlines 6 steps for spring mapping and revival: mapping, data monitoring setup, understanding governance systems, hydrogeological mapping, spring management protocol, and measuring

FIGURE 1 Overview of the study sites. (A) Location of Fulbari water tower in relation to Kathmandu. (B) Fulbari water tower and location of the 2 study sites within it. (C) Patalako Dhara spring and its surroundings. (D) Bhagwate Pakhako Kuwa spring and its surroundings.





FIGURE 2 Diagram showing 2 different ways of conceptualizing water resources and households in a village. A simplified one where each household is supported by one source (A) and another model (B1, B2) where households can have multiple sources with a primary or secondary role that may change across the seasons. In a benefit–cost analysis (BCA), the 2 models would lead to different weighting for each water source and likely to different results. Arrows point to the water source used by a household.



impact. The project identified 50 springs (28 active, 22 dry) in the Fulbari water tower (Pokhrel et al 2019; Khadka et al 2021), 2 of which were selected for revival.

The Bhagwate Pakhako Kuwa spring is located at 1235 masl, with 1 household (HH) using it as its primary water source and 24 HHs as a secondary/alternative source. The recharge area of the spring lies within the community forest. Downhill from the spring, there is a plains area with several dug wells. The spring was drying mainly due to the rapid increase in the number of dug wells and consequent overextraction. Research shows that it dried up in 2019, possibly due to lowering of the water table (Pokhrel et al 2019). For the revival activities, 5 ponds, 16 trenches, 4 check dams, and 8 palisades were constructed.

The Patalako Dhara spring is located at 1412 masl in Damaidada village, with 21 HHs using it as their primary water source and 14 HHs as a secondary one. The recharge area is partially within community forest and partially within private fields. The spring has been experiencing a decline due to factors such as variability in rainfall intensity and duration, the impact of the 2015 earthquake, and the abandonment of traditional ponds. For the revival activities, 9 trenches were dug and a pond was constructed to enhance groundwater recharge. In 2022 the community further improved their water supply system by installing a pump to transfer water to a water tank. This upgrade allowed for the gravity-driven distribution of water to 21 HHs directly.

The application of the BCA method in this study, though limited to 2 sites, is representative of the spring system in the mid-hill region of the Himalaya. The mid-hills, characterized by longitudinal parallel zones, share regional similarities, particularly in their geology, which influences the spring system (Dhital 2015). The cases studied focus on depression-type springs, the most common type in the mid-hills. Both formal and informal systems exist for managing these spring sources throughout the region. The spring–HH dependency structure illustrated in Figure 2 is also typical of other parts of the mid-hills.

Along with these study sites (scenarios PAT and BHA), 2 more generic scenarios were included in the analysis. The aim was to estimate the potential costs and benefits

(1) where ecosystem benefits are fully developed (GEB, for “generic, ecosystem benefits”) and (2) where supply needs of the surrounding households are fully provided for (GWS, for “generic, water supply”).

Qualitative mapping

Before the BCA economic valuation, qualitative mapping was done to identify the effects of spring revival activities for later quantification. To do so, a literature review was performed from May to July 2023 using Scopus and Semantic Scholar. Key themes were explored with search strings like “water security AND (Himalaya OR Nepal)” (38 relevant documents), “climate change AND Himalaya” (17 documents), and “(spring OR springshed) AND (Himalaya OR Nepal OR HKH OR Hindu Kush OR revival)” (38 documents). Expert opinions (4) were also gathered from within ICIMOD in September 2023 using semistructured interviews focusing on specific themes according to the experts’ background.

Furthermore, in September 2023, focus group discussions (FGDs) were held with local communities residing near the springs. Participants were primarily selected from water user groups (WUGs). WUGs are typically structured to reflect the demographic diversity of the municipality, ensuring balanced representation in terms of gender, age, and other social characteristics. In this case, the groups included 2 males and 6 females at Patalako Dhara, and 5 males and 3 females at Bhagwate Pakhako Kuwa, spanning various age groups and including minority populations.

Given the time constraints of the study and the communities’ familiarity with this method, FGDs were chosen for their effectiveness in capturing diverse perspectives and fostering equal participation of those most impacted by spring revival activities (Sim 1998; Krueger and Casey 2015; Sangaramoorthy and Kroeger 2020).

Benefit–cost analysis of spring revival

After the qualitative mapping, an economic valuation through BCA was performed (Hanley and Barbier 2009; OECD 2015; Harris and Roach 2017; Boardman et al 2018; Abelson 2020). BCA is a methodology that compares the total costs of implementing an activity (in this case the revival of a

spring), against the expected benefits quantified in the same currency. The aim is to determine whether the benefits outweigh the costs. Although this method has been widely used for policy advice, its application to environmental remediation is more recent (Brauman et al 2007; Atkinson and Mourato 2008; Mendelsohn and Olmstead 2009; Ratnaweera et al 2021). In Nepal, BCAs have been carried out to estimate the willingness to pay for conservation activities (Lamsal et al 2015), environmental costs (Pakhtigian and Jeuland 2019), ecosystem services (Kc et al 2013; Thapa, Wang, et al 2020), households' commitment to contributing to river restoration (Khatiwada et al 2023), and, in Sikkim, soil and water conservation (Mishra and Rai 2014). In this study, the BCA adopts the national standing (Abelson 2020), aligning with Boardman (2018) and national guidelines (Abelson 2020). The chosen perspective affects whether a factor is seen as a cost or benefit.

Results

Cause–effect mapping

The qualitative mapping data and findings were organized into benefits and costs, with each category associated with a numerical equation to facilitate quantitative estimation. The costs were sourced from project data and categorized as follows: material costs (C_{mat}), skilled labor for project management (C_{lab}), skilled labor for mapping and design of infiltrations (C_{lab_wt}), site supervision and unskilled labor for manual work (C_{unlab}), operation and maintenance (O&M) costs for things such as pumps, operator salaries, and repairs (C_{om}), as well as financial and opportunity costs linked to loan interest and other possible investments (C_{fin}).

The identified benefits included ecosystem benefits (B_{ecos}), increased household crop productivity due to greater water availability (B_{prod}), time-saving benefits (B_{time}) from reduced water fetching, and health benefits (B_{health}), particularly from reduced gastrointestinal disease. Additional benefits, such as cultural benefits (B_{cult}) linked to the spring, benefits from land price increase (B_{land}) due to proximity of revived water source, and increased social capital (B_{soc}) were only identified but not quantified. The parameters and equations used for the quantification are provided in Tables 2 and 3.

All results and data, unless stated otherwise, are expressed in NPR 100,000 (LNPR) equal to EUR 707, USD 750, or CNY 5477 as per October 2023. Most parameters were defined using statistical distributions such as uniform ($U(\min, \max)$) and normal ($N(\text{mean}, SD)$). Risks and uncertainty were addressed using a Monte Carlo simulation with R package MC2D (Pouillot and Delignette-Muller 2010). Global sensitivity analysis was carried out using the Multisensi package in R (Bidot et al 2018).

A key finding from the mapping phase highlights the complexity of local water supply sources. HHs often rely on multiple sources, including nearby springs, distant piped water, and wells. Although HHs may have a preferred primary source, their choice often shifts due to factors such as seasonal changes, temporary scarcity, and ongoing initiatives that modify the system. Consequently, the impact of each spring varies depending on its role within the broader local water supply system (Figure 2).

The effects of spring revival activities were synthesized and presented in terms of cause and effect (Figure 3). The 2 main activities related to the revival of springs—mapping with capacity building and construction of infiltration structures—yield various outputs. While some short-term effects, such as time savings, health benefits, ecosystem services (ES), employment increase, and productivity, were used in the BCA analysis, others, such as land price and increased social capital, were noted but excluded from the quantification. Regarding land price, while some literature on Nepal exists (Joshi et al 2017), it was not possible to retrieve enough updated information on the study sites to include the effect in the BCA. Social capital increase, on a short time range, was considered not to be significant from a quantification point of view. Long-term effects, such as improvements in employment, children's health, and nutritional development, were acknowledged but omitted due to the difficulty of establishing a direct causal relationship with spring revival.

Specific case studies illustrate the varied impacts of these revival activities. The Patalko Dhara spring serves as the sole water source for 7 HHs and the primary source for 14 others. The revival led to additional infrastructure improvements, including a cement path, fencing, and a water pump, supported by a nongovernmental organization, that now deliver water directly to households via a gravity-fed pipeline. These enhancements resulted in benefits from reduced expenses for water trucking, increased livestock, and higher crop production. However, benefits from ES were minimal, as the surrounding area is little used for productive activities.

In contrast, the Bhagwate Pakhako Kuwa case presents a different scenario. Here, the spring is located near multiple water sources (3 springs, 20 hand wells, and 1 borehole) that diminish its direct impact on the local community, relegating it to the status of an emergency water source. The spring water is primarily used as an additional source for irrigation of nearby fields and as a drinking source during agricultural work in the vicinity. Regarding ecosystem benefits, the community has observed increased vegetation in the recharge area and is considering planting *timmur* (*Zanthoxylum armatum*) and *tite pati* (*Artemisia vulgaris*) (Equation 1 in Table 3) due to their potential income generation.

BCA results

Skilled labor for mapping and design of infiltration was the main cost in 3 scenarios (GEB, GWS, BHA), while cost of operations was the highest for PAT. Total costs were in the range of LNPR 12–16 for 3 scenarios (GEB, GWS, BHA), whereas they were higher in PAT (LNPR 31) due to the O&M costs of the pump (Table 4). Over a 10-year period, benefits were the lowest for BHA (LNPR 6.28), due to the limited use of the spring, but were higher in the 2 generic scenarios (LNPR 32.25 for GEB and LNPR 44.09 for GWS), and the highest for PAT (LNPR 52.72) because of the recently established piped water supply to 21 HHs. Therefore, the benefit–cost ratio (BCR) was below 1 for BHA (0.24) and highest for GEB (4.03) and GWS (3.49). The internal rate of return (IRR)—a parameter commonly used to identify the profitability of an investment by determining

TABLE 2 Summary of parameters used in the BCA. Uniform distribution parameters are written as $U(\min, \max)$, and normal distributions are written as $N(\text{mean}, \text{standard deviation})$. (Table continued on next page.)

Parameter	Description	Bhagwate (BHA)	Patal (PAT)	General environmental benefit (GEB)	General water supply (GWS)
C_{mat}	Construction material (stone, wood), land and tools used for the physical construction of revival measures (LNPR per spring)	$N(1.2, 0.3)$	$N(0.8, 0.3)$	$N(1.2, 0.3)$	$N(1.2, 0.3)$
C_{lab}	Skilled labor for setting up data monitoring and project management (LNPR per spring)	$U(1.5, 2.5)$			
C_{unlab}	Includes labor used for the construction and for site supervision. This is likely to be sourced locally (LNPR per spring)	$U(0.8, 1.3)$			
C_{om}	Includes costs to cover maintenance of infiltration measures as well as costs for energy consumption and labor in the case of pump installation (LNPR per spring)	$U(0.1, 0.2)$	$U(1.4, 1.8)$	$U(0.1, 0.2)$	$U(1.4, 1.8)$
C_{fin}	Includes cost of capital, assumed at 8% (LNPR per year)	$U(0.16, 0.26)$			
C_{lab_wt}	Includes fees paid to experts and technician for the water tower mapping, hydrogeological survey, measuring impact and detailed revival measures	$U(8, 10)$	$U(8, 10)$	$U(2, 10)$	$U(2, 10)$
R_{ew}	Rate of improvement caused by the additional water flow on the ecosystem assumption based on expert opinion (LNPR per project)	$U(0.15, 0.30)$			
V_{ecos}	Value of the ecosystem services in LNPR/hectare based on Pant (2009) and Taye et al (2021)	$U(0.6, 1.4)$			
A_{ecos}	Area of affected ecosystem as the sum of the recharge area and the downstream area until a road or stream is met	$U(1, 3.5)$			
W_{saved}	Water saved from purchasing due to flow increase, liters per year		$U(9000, 15000)$		$U(9000, 15000)$
W_{cost}	Price per liter in NPR		$U(0.7, 1.2)$		$U(0.7, 1.2)$
HH	Number of households using the spring	$U(1, 5)$	$U(11, 21)$	$U(5, 30)$	$U(5, 30)$
F_{usage}	Fraction of HH saving water, this is to account for the fact not all houses will be saving water	0.66			
W_{prod}	Water productivity at HH level related to crop production, based on Dhakal (2020) with production of NPR 6000/year and in this case only for 6 months, therefore LNPR 0.03	Not used, see Equations 2 and 3 in Table 3	$U(0.02, 0.04)$	Not used, see Equations 2 and 3 in Table 3	$U(0.02, 0.04)$
Q_{rev}	Increased water flow due to revival activity, liters per day.	$U(800, 1900)$	Not used, see Equations 2 and 3 in Table 3	$U(500, 2000)$	Not used, see Equations 2 and 3 in Table 3

TABLE 2 Continued. (First part of Table 2 on previous page.)

Parameter	Description	Bhagwate (BHA)	Patal (PAT)	General environmental benefit (GEB)	General water supply (GWS)
$V_{w,prod}$	Water productivity, NPR/1000 L	$U(5.6, 12)$		$U(5.6, 12)$	
T_{save}	Time saved from a single trip to the spring in hours. Based on community consultation.	$U(0, 0.1)$	$U(0.1, 0.5)$	$U(0.1, 1)$	$U(0.1, 1)$
VOT	Value of time, based on a value of 50–80% (Whittington and Cook 2019) of daily unskilled labor salary (NPR 1000/day). Estimate also considers the 133 NPR/hour given to water pump operator in PAT (NPR 7000/month)	$U(62, 100)$			
$R_{t,prod}$	Fraction of time reallocated to productive activities considering weekly productive hours as 23–28 (Yokying et al 2023; Picchioni 2020)	$U(0.15, 0.5)$			
$R_{loc,l}$	Ratio of labor procured locally	Not used in this BCA, see equation 6 in Table 3			
$R_{loc,m}$	Ratio of material purchased locally	Not used in this BCA, see equation 6 in Table 3			
DW	Disability weight for gastroenteritis as 0.18 DALY (IHME 2020) \times 2 events per year (Shlim et al 1999) lasting 4 days	1.44 days			
$R_{ill,red}$	Reduction of illness due to increased water, based on Howard et al (2020)	20%			
GDP_{pppy}	Gross domestic product per person per day based on US\$1336 per year (data.world.bank.org)	NPR 485			
COI	Cost of illness that would include medicine and travel to a doctor	NPR 1000			

Note: DALY, disability-adjusted life years.

the discount rate that would lead to a Net Present Value of 0—was negative for BHA and in the range of 144% and 405% for the other scenarios (Figure 4). A comparison across scenarios is detailed in Figure 5. Linear regression on the 2 generic scenarios (GEB, GWS) showed that BCR and IRR can be predicted ($R^2 = 0.85$) with HH and T_{save} for GEB, and with HH, T_{save} , and $R_{t,prod}$ for GWS (Table 3).

Discussion and recommendations

This study aligns with previous research in identifying direct-use benefits from natural resources (Tenge 2005; Mishra and Rai 2014), employment opportunities (Kihila et al 2014), and ecosystem benefits (Thapa, Pant, et al 2020; Taye et al 2021). However, it also emphasizes indirect benefits related to human systems, such as enhancing social capital through knowledge acquisition, reducing community tensions, and appreciating land values—factors often overlooked in BCAs (Kc et al 2013; Mishra and Rai 2014; Acharya and Dhungel 2021; Mcharo and Maghenda 2021). While these benefits were identified, they were not quantified, underscoring the need for further research on

community participation (Bhandari et al 2005; Longworth 2022), water management conflict (Upreti et al 1999; Upreti 2001; Devkota 2018), impact on land prices (Joshi et al 2017; Nepal et al 2017), and cultural benefits (Roebeling et al 2016).

Cost quantification revealed that hydrogeological mapping activities ($C_{lab,wt}$) were the most impactful, except in scenarios where water supply fees were applied (C_{om} in PAT). This suggests that using an aquifer-based approach for revival can provide economies of scale benefits since mapping costs would be spread across different springs.

Time-saving benefits were particularly significant. However, their quantification is highly dependent on how the saved time is reallocated (Yokying et al 2023) and the difficulty in isolating the factors influencing this decision (Cooke 1998; Koolwal and Walle 2016). Consequently, these estimates should be interpreted with caution and further research is recommended, especially since the value of time (VOT) in rural areas is likely to change in the coming years.

ES and productive benefits were 2 other relevant categories. Although ES were primarily focused on water

TABLE 3 Summary of equations used in the BCA.

Equation no.	Symbol	Equation	Description
1	B_{ecos}	$B_{\text{ecos}} = R_{\text{ew}} \times V_{\text{ecos}} \times A_{\text{ecos}}$	Ecosystem benefits for GEB and BHA scenario
2	B_{ecos}	$B_{\text{ecos}} = W_{\text{saved}} \times W_{\text{cost}} \times \text{HH} \times F_{\text{usage}}$	Ecosystem benefits for GWS and PAT scenario
3	B_{prod}	$B_{\text{prod}} = W_{\text{prod}} \times \text{HH}$	Economic benefits from increased productivity due to increase in water use for PAT and GWS
4	B_{prod}	$B_{\text{prod}} = Q_{\text{rev}} \times 365 \times V_{\text{w.prod}}$	Economic benefits from increased productivity due to increase in water use for GEB and BHA
5	B_{time}	$B_{\text{time}} = T_{\text{save}} \times 4 \times \text{VOT} \times \text{HH} \times 365 \times R_{\text{t.prod}}$	Economic benefits from time saving assuming 4 trips per day
6	B_{impl}	$B_{\text{impl}} = (C_{\text{lab}} + C_{\text{unlab}}) \times R_{\text{loc.l}} + C_{\text{mat}} \times R_{\text{loc.m}}$	Benefits from activity implementation. Giving the standing of this BCA this item would be in the costs as well as the benefits and therefore was not accounted for. Different BCA standing would bring different results.
7	B_{health}	$\text{HH} \times 4 \times \text{DW} \times R_{\text{ill.red}} \times (\text{GDP}_{\text{pppy}}/365 + \text{COI})$	Health benefit for population ($\text{HH} \times 4$ people)
8	PV	$\text{PV}(X_t) = X_t [(1+i)^{-t}]$	Present value, where t stands for time (10 years in this case) and i is the discount rate. The discount rate was modeled as $U(6\%, 12\%)$ considering previous studies with 6% (Mishra and Rai 2014) and 12% (Kc et al 2013).
9	NPV	$\text{NPV} = \sum B_t(1+i)^{-t} - \sum C_t(1+i)^{-t}$	Net present value, where B_t is the total benefit, C_t the total costs, i the interest rate, and t the time period in years.
10	BCR	$\text{BCR} = \frac{B_t}{C_t}$	Benefit–cost ratio, where a value above 1 is considered positive, as it indicates that benefits outweigh costs, whereas a value below 1 is considered negative.
11	IRR	$0 = \text{NPV} = \sum_{t=1}^T \frac{C_t}{(1 + \text{IRR})^t} - C_0$	Internal rate of return, where C_t is the net cash flow during period t , and C_0 is the total initial investment cost.
12	BCR_{GEB}	$\text{BCR}_{\text{GEB}} = -1.5 + 0.2\text{HH} + 3.7T_{\text{save}}$	Linear regression for BCR in the GEB scenario, R^2 of 0.85 and sensitivity index of $T_{\text{save}} = 0.29$ and $\text{HH} = 0.39$.
13	IRR_{GEB}	$\text{IRR}_{\text{GEB}} = -0.35 + 0.18\text{HH} + 0.52T_{\text{save}}$	Linear regression for IRR in the GEB scenario, R^2 of 0.85 and sensitivity index of $T_{\text{save}} = 0.29$ and $\text{HH} = 0.39$.
14	BCR_{GWS}	$\text{BCR}_{\text{GWS}} = -3.93 + 0.2\text{HH} + 3.5T_{\text{save}} + 10R_{\text{t.prod}}$	Linear regression for BCR in the GWS scenario, $R^2 = 0.85$ and GSI as $\text{HH} = 0.57$, $T_{\text{save}} = 0.2$, $R_{\text{t.prod}} = 0.08$
15	IRR_{GWS}	$\text{IRR}_{\text{GWS}} = -0.6 + 0.026\text{HH} + 0.42T_{\text{save}} + 1.2R_{\text{t.prod}}$	Linear regression for IRR in the GWS scenario, $R^2 = 0.85$ and GSI as $\text{HH} = 0.57$, $T_{\text{save}} = 0.2$, $R_{\text{t.prod}} = 0.08$

Note: GSI, Generalized Sensitivity Indices (Bidot et al 2018).

provision, they may have been underestimated, as prior research indicates a broader range of potential benefits (Taye et al 2021). Productive benefits could not be fully assessed due to the lack of relevant agricultural activities linked to the springs. These factors resulted in BCR outcomes in the 4 scenarios with ratios (mean 2.35, min 0.24, max 4.05) consistent with published studies on watershed management in India by Joshi et al (2008) that reported BCR with mean of 2.00, minimum of 0.80, and maximum of 7.30. The higher BCR in the generic scenarios suggests that springs have greater potential if their full

capacity is developed. This could be achieved by focusing on springs that serve more HHs, increasing water productivity through crop cultivation, reducing O&M costs via semiautomatic controls, and reducing mapping costs through citizen science.

Data limitations prevented disaggregation by gender, age, or sociocultural groups. However, existing studies show that women disproportionately bear the burden of water fetching (Shrestha and Clement 2019; Wali et al 2020; Raut 2023), which affects their time (Yokying et al 2023) and energy (Picchioni et al 2020). This exacerbates women's dual roles in productive

FIGURE 3 Cascade diagram, based on literature and community consultation, showing activities, results, and outcomes of spring revival and what benefits it may generate.




TABLE 4 Summary of BCA results for the 4 scenarios (all values in NPR 100,000): Bhagwate Pakhako Kuwa (BHA), Patalko Dhara (PAT), a generic scenario with fully developed ecosystem benefits (GEB), and a generic scenario with avoided water purchasing (GWS).

Parameter	GEB	BHA	PAT	GWS
C_{mat}	1.14	1.15	1.15	1.15
C_{unlab}	1.06	1.05	1.04	1.05
C_{om}	1.52	1.51	16.03	1.50
C_{lab}	2.00	1.99	1.98	2.02
C_{fin}	2.10	2.12	2.11	2.09
C_{lab_wt}	4.24	8.96	8.98	4.26
Total costs	12.07	16.78	31.29	12.07
B_{ecos}	4.92	3.07	13.30	13.06
B_{prod}	0.57	0.43	5.05	5.20
B_{time}	23.66	0.40	31.29	22.66
B_{health}	0.89	0.17	0.89	0.97
IRR	0.22	-0.24	0.18	0.36
Total benefits	32.25	6.28	52.72	44.09
NPV	8.53	-4.52	9.05	13.62
BCR	4.05	0.24	1.62	3.49

Note: See Table 2 for a detailed explanation of the parameters.

and reproductive work. Although research on children’s involvement in water fetching is limited (Whittington and Cook 2019), evidence suggests that reducing this burden can improve school attendance and health outcomes for children (Nankhuni and Findeis 2004; Koolwal and Walle 2016; Nauges and Strand 2017). These findings suggest that spring revival activities are likely to benefit more vulnerable groups and contribute to reducing gender disparities.

Last, the BCA has some limitations related to its temporal scope. The 10-year time frame may be insufficient given the rapid socioeconomic changes happening in Nepal. Furthermore, the hierarchical and dynamic nature of water source systems was not fully captured, which could benefit from incorporating methods such as system mapping, network analysis (Hevey 2018), and longitudinal data collection to calibrate a dynamic model.

Conclusion

The research employed a mixed-method approach to comprehensively examine the qualitative and quantitative ramifications of spring revival activities in the mid-hill region of Nepal. This mixed methodology proved beneficial in capturing the diverse and complex ramification of revival activities.

The BCA analysis, run on 4 different scenarios over a 10-year timeframe, showed a positive net present value (NPV) and BCR in 3 scenarios (PAT, GEB, GWS) and negative results for 1 scenario (BHA). Overall, costs were mainly influenced by mapping activities, while benefits stem from

FIGURE 4 Histogram showing benefit–cost ratio (BCR) in the 4 scenarios (top row) and internal rate of return (IRR) (bottom row). Red line in BCR histograms at BCR = 1 and IRR histograms at IRR = 0.3 as a reference value.





FIGURE 5 Comparison between (A) cost types across the 4 scenarios and (B) contributions of each cost type over a 10-year period for each scenario, and between (C) benefit types across the 4 scenarios and (D) contributions of each benefit type over a 10-year period for each scenario. Values are in 100,000 NPR. See Table 2 for a detailed explanation of the parameters.



time-saving and ecosystem improvements. A linear regression on BCR and IRR results showed that for the 2 generic scenarios, 3 parameters (household number, time saved, and share of time allocated to productive activities) could accurately predict final estimates.

From a sustainability perspective, revival activities impact surrounding communities and the environment in multiple ways, generally with benefits outweighing costs. These activities address the 3 pillars of sustainability (environment, economy, people) and can therefore be considered an effective solution for developing sustainable communities.

Policy recommendations

This study indicates that spring revival activities can yield a positive BCR, despite the fact that some benefits, such as acquired skills, increased social capital, and conflict reduction, cannot be easily monetized. Policymakers should consider this limitation when relying on BCA.

Furthermore, spring revival activities were found to be catalysts for other infrastructural improvements. This indicates that such activities need to be thought of and implemented in synergy with other upgrades happening in the community.

While the suggestion to adopt a springshed (Dahal et al 2021) or an aquifer-based approach is valid, it may find challenges at governance level when the intervention area spans across different wards and municipalities. Policies should account for this aspect and support coordination across actors.

Raising capital for O&M costs is a crucial policy intervention (Rathod et al 2021), as municipalities often fail to allocate adequate budgets for this purpose (Dahal et al 2021; Thapa et al 2023). Although many benefits of spring revival activities are not marketable, this study indicates a willingness to pay for water supply services, particularly in the PAT scenario. The findings demonstrate long-term economic and social benefits of spring revival activities, justifying investments in fund allocation and potentially attracting private investments by offering a clear economic rationale. Policymakers should prioritize securing funds for O&M costs and leverage community willingness to pay to sustain water services. Additionally, policymakers should justify and support investments in these activities.

As demonstrated, this study enhances our understanding of the impact of spring revival in mid-hill rural settings, showing that revival activities are likely to have high BCR if all potential benefits for the surrounding community are fully realized. Therefore, it is recommended that longitudinal monitoring mechanisms be established within the project's beneficiary communities to gain deeper insights into the evolving impacts of spring rejuvenation over time.

ACKNOWLEDGMENTS

We thank Madhav Dhakal, Anju Pandit, Abid Hussain, Sunita Chaudhary, Erica Udas, Bandana Shakya, and Farid Ahmad from ICIMOD and Rosalind Bryce from the University of the Highlands and Islands (UHI) for their advice and support. We also express our gratitude to the communities living around Fulbari water tower, Bhakundebesi, Kavrepalanchok, Nepal, for their patient support. This article was supported by ICIMOD's Himalayan Resilience Enabling Action Programme (HI-REAP), funded by United Kingdom International Development (UKID) through its Foreign Commonwealth and Development Office (FCDO). ICIMOD and its regional member countries gratefully acknowledge the generous support of Austria,

Norway, Sweden, and Switzerland for core and program funding, and Australia, Canada's International Development Research Centre, the European Union, Finland, Germany, the United Kingdom, the United States, and the World Bank.

DISCLAIMER

The views and interpretations in this publication are those of the authors and are not necessarily attributable to ICIMOD and do not imply the expression of any opinion by ICIMOD concerning the legal status of any country, territory, city, or area of its authority, or concerning the delimitation of its frontiers or boundaries, or the endorsement of any product.

REFERENCES

- Abelson P.** 2020. A partial review of seven official guidelines for cost-benefit analysis. *Journal of Benefit-Cost Analysis* 11(2):272–293. <https://doi.org/10.1017/bca.2020.3>.
- Acharya KR, Dhungel BD.** 2021. Cost benefit analysis of mushroom farming in Nepal. *Asian Journal of Economics, Finance and Management* 5(4):16–25.
- Adhikari S, Gurung A, Chauhan R, Rijal D, Dongol BS, Aryal D, Talchabhadel R.** 2021. Status of springs in mountain watershed of western Nepal. *Water Policy* 23(1):142–156. <https://doi.org/10.2166/wp.2020.187>.
- Agarwal A, Bhatnaga NK, Nema RK, Agrawal NK.** 2012. Rainfall dependence of springs in the midwestern Himalayan hills of Uttarakhand. *Mountain Research and Development* 32(4):446–455. <https://doi.org/10.1659/MRD-JOURNAL-D-12-00054.1>.
- Akhtar M, Ahmad N, Booij MJ.** 2008. The impact of climate change on the water resources of Hindukush–Karakorum–Himalaya region under different glacier coverage scenarios. *Journal of Hydrology* 355(1–4):148–163. <https://doi.org/10.1016/j.jhydrol.2008.03.015>.
- Al-Manni DAMA, Saleh KA.** 2019. Delineation of spring protection zone and vulnerability mapping of selected springs in Sulaymaniyah area, Kurdistan, Iraq. *Environmental Earth Sciences* 78(21):622. <https://doi.org/10.1007/s12665-019-8632-2>.
- Andermann C, Longuevergne L, Bonnet S, Crave A, Davy P, Gloaguen R.** 2012. Impact of transient groundwater storage on the discharge of Himalayan rivers. *Nature Geoscience* 5(2):127–132. <https://doi.org/10.1038/ngeo1356>.
- Atkinson G, Mourato S.** 2008. Environmental cost-benefit analysis. *Annual Review of Environment and Resources* 33(1):317–344. <https://doi.org/10.1146/annurev.enviro.33.020107.112927>.
- Bajracharya SR, Maharjan SB, Shrestha F, Guo W, Liu S, Immerzeel W, Shrestha B.** 2015. The glaciers of the Hindu Kush Himalayas: Current status and observed changes from the 1980s to 2010. *International Journal of Water Resources Development* 31(2):161–173. <https://doi.org/10.1080/07900627.2015.1005731>.
- Bhandari BS, Grant M, Pokharel D.** 2005. Sustainable community water: Managing supply systems in the mid-hills of Nepal. *Water Policy* 7(2):201–214. <https://doi.org/10.2166/wp.2005.0013>.
- Bharti N, Khandekar N, Sengupta P, Bhadwal S, Kochhar I.** 2020. Dynamics of urban water supply management of two Himalayan towns in India. *Water Policy* 22(S1):65–89. <https://doi.org/10.2166/wp.2019.203>.
- Bhat SU, Pandit AK.** 2020. Water quality assessment and monitoring of Kashmir Himalayan freshwater springs: A case study. *Aquatic Ecosystem Health and Management* 23(3):274–287. <https://doi.org/10.1080/14634988.2020.1816771>.
- Bidot C, Monod H, Taupin M-L, Paris-Saclay U.** 2018. A quick guide to multisensi, an R package for multivariate sensitivity analyses. CRAN R-universe. <https://cran.r-universe.dev/multisensi/doc/multisensi-vignette.pdf>; accessed on 20 October 2023.
- Blair P, Buytaert W.** 2016. Socio-hydrological modelling: A review asking “why, what and how?” *Hydrology and Earth System Sciences* 20(1):443–478. <https://doi.org/10.5194/hess-20-443-2016>.
- Boardman AE, Greenberg DH, Vining AR, Weimer DL.** 2018. *Cost-Benefit Analysis: Concepts and Practice*. 5th edition (1st edition 1996). Cambridge: Cambridge University Press.
- Bookhagen B.** 2012. Himalayan groundwater. *Nature Geoscience* 5(2):97–98. <https://doi.org/10.1038/ngeo1366>.
- Brauman KA, Daily GC, Duarte TK, Mooney HA.** 2007. The nature and value of ecosystem services: An overview highlighting hydrologic services. *Annual Review of Environment and Resources* 32(1):67–98. <https://doi.org/10.1146/annurev.energy.32.031306.102758>.
- Chapagain PS, Ghimire M, Shrestha S.** 2019. Status of natural springs in the Melamchi region of the Nepal Himalayas in the context of climate change. *Environment, Development and Sustainability* 21(1):263–280. <https://doi.org/10.1007/s10668-017-0036-4>.
- Chen Z, Lucianetti G, Hartmann A.** 2023. Understanding groundwater storage and drainage dynamics of a high mountain catchment with complex geology using a semi-distributed process-based modelling approach. *Journal of Hydrology* 625:130067. <https://doi.org/10.1016/j.jhydrol.2023.130067>.
- Chinnasamy P, Prathapar SA.** 2016. Methods to Investigate the Hydrology of the Himalayan Springs: A Review. *IWMI Working Paper* 169. Colombo, Sri Lanka: IWMI [International Water Management Institute]. <http://www.iwmi.cgiar.org/publications/iwmi-working-papers/iwmi-working-paper-169/>; accessed on 15 June 2023.

- Cooke PA.** 1998. The effect of environmental good scarcity on own-farm labor allocation: The case of agricultural households in rural Nepal. *Environment and Development Economics* 3(4):443–469. <https://doi.org/10.1017/S1355770X98000230>.
- Dahal N, Bhattarai, Chiranjibi, Thapa B, Tiwari S, Shrestha K, Agrawal K, Buchar S, Shresth BB, Dhakal M.** 2021. Reviving springs of the Himalayas an imperative for policy support. Policy brief. Kathmandu, Nepal: ICIMOD [International Center for Integrated Mountain Development] and NWC [Nepal Water Conservation Foundation]. <https://waternepal.org.np/project/reviving-springs-of-the-himalayas-an-imperative-for-policy-support/detail>; accessed on 15 August 2024.
- Daniel D, Anandhi A, Sen S.** 2021. Conceptual model for the vulnerability assessment of springs in the Indian Himalayas. *Climate* 9(8):121. <https://doi.org/10.3390/cli9080121>.
- Devkota N, Phuyal RK.** 2018. Adoption practice of climate change adaptation options among Nepalese rice farmers: Role of information and communication technologies (ICTs). *American Journal of Climate Change* 7(2):135–152. <https://doi.org/10.4236/ajcc.2018.72010>.
- Dhakal MP.** 2021. *Springs in the Godavari Landscape, Nepal Mapping, Governance, and Revival*. Kathmandu, Nepal: ICIMOD [International Center for Integrated Mountain Development]. <https://lib.icimod.org/record/35671>; accessed on 10 July 2023.
- Dhital R.** 2015. *Geology of the Nepal Himalaya*. Cham, Switzerland: Springer. <http://link.springer.com/10.1007/978-3-319-02496-7>; accessed on 20 September 2024.
- Di Baldassarre G, Viglione A, Carr G, Kuil L, Yan K, Brandimarte L, Blöschl G.** 2015. Debates—Perspectives on socio-hydrology: Capturing feedbacks between physical and social processes. *Water Resources Research* 51(6):4770–4781. <https://doi.org/10.1002/2014WR016416>.
- Faulon M, Sacareau I.** 2020. Tourism, social management of water and climate change in an area of high altitude: The Everest massif in Nepal. *Revue de Géographie Alpine* 108(1). <https://doi.org/10.4000/rga.6779>.
- GGGI.** 2018. *Namobuddha Municipality, Nepal Situation Analysis for Green Municipal Development*. Seoul, South Korea: Global Green Growth Institute. https://gggi.org/wp-content/uploads/2018/07/GGGI_GMD-Assessment_Namobuddha.pdf; accessed on 20 July 2023.
- Ghimire CP, Buijzelaar LA, Lubczynski MW, Bonell M.** 2014. Negative trade-off between changes in vegetation water use and infiltration recovery after reforesting degraded pasture land in the Nepalese Lesser Himalaya. *Hydrology and Earth System Sciences Discussions* 11:3437–3479. <https://doi.org/10.5194/hessd-11-3437-2014>.
- Ghimire M, Chapagain PS, Shrestha S.** 2019. Mapping of groundwater spring potential zone using geospatial techniques in the Central Nepal Himalayas: A case example of Melamchi–Larke area. *Journal of Earth System Science* 128(2):26. <https://doi.org/10.1007/s12040-018-1048-7>.
- Gober P, Wheeler HS.** 2015. Debates—Perspectives on socio-hydrology: Modeling flood risk as a public policy problem. *Water Resources Research* 51(6):4782–4788. <https://doi.org/10.1002/2015WR016945>.
- Gurung A, Adhikari S, Chauhan R, Thakuri S, Nakarmi S, Ghale S, Dongol BS, Rijal D.** 2019. Water crises in a water-rich country: Case studies from rural watersheds of Nepal's mid-hills. *Water Policy* 21(4):826–847. <https://doi.org/10.2166/wp.2019.245>.
- Gurung A, Adhikari S, Chauhan R, Thakuri S, Nakarmi S, Rijal D, Dongol BS.** 2019. Assessment of spring water quality in the rural watersheds of western Nepal. *Journal of Geoscience and Environment Protection* 7(11):39–53. <https://doi.org/10.4236/gep.2019.711004>.
- Hanley N, Barbier E.** 2009. *Pricing Nature: Cost–Benefit Analysis and Environmental Policy*. Cheltenham, UK, Northampton, MA: Edward Elgar.
- Harris JM, Roach B.** 2017. *Environmental and Natural Resource Economics: A Contemporary Approach*. 4th edition (1st edition 2002). New York, NY: Routledge.
- Herrera-Franco G, Montalván-Burbano N, Carrión-Mero P, Bravo-Montero L.** 2021. Worldwide research on socio-hydrology: A bibliometric analysis. *Water* 13(9):1283. <https://doi.org/10.3390/w13091283>.
- Herschty R.** 1993. The velocity–area method. *Flow Measurement and Instrumentation* 4(1):7–10. [https://doi.org/10.1016/0955-5986\(93\)90004-3](https://doi.org/10.1016/0955-5986(93)90004-3).
- Hevey D.** 2018. Network analysis: A brief overview and tutorial. *Health Psychology and Behavioral Medicine* 6(1):301–328. <https://doi.org/10.1080/21642850.2018.1521283>.
- Huber A, Joshi D.** 2015. Hydropower, anti-politics, and the opening of new political spaces in the eastern Himalayas. *World Development* 76:13–25. <https://doi.org/10.1016/j.worlddev.2015.06.006>.
- IHME [Institute for Health Metrics and Evaluation].** 2020. *Global Burden of Disease Study 2019 (GBD 2019) Disability Weights*. Seattle, WA: IHME. <https://doi.org/10.6069/1W19-VX76>.
- Joseph G, Shrestha A.** 2022. *Glaciers, Rivers, and Springs: A Water Sector Diagnostic of Nepal*. Washington, DC: World Bank. <http://elibrary.worldbank.org/doi/book/10.1596/37236>; accessed on 8 June 2023.
- Joshi J, Ali M, Berrens RP.** 2017. Valuing farm access to irrigation in Nepal: A hedonic pricing model. *Agricultural Water Management* 181:35–46. <https://doi.org/10.1016/j.agwat.2016.11.020>.
- Joshi P, Jha A, Wani S, Sreedevi T, Shaheen F.** 2008. *Impact of Watershed Program and Conditions for Success: A Meta-Analysis Approach*. Global Theme on Agrosystems Report No. 46. Andhra Pradesh, India; International Crops Research Institute for the Semi-Arid Tropics. <https://oar.icrisat.org/2351/>; accessed on 10 July 2023.
- Kc B, Kandel P, Adhikari S.** 2013. Economic valuation of ecosystem services in protected areas: A case study from Nepal. *Banko Janakari* 23(1):42–50. <https://doi.org/10.3126/banko.v23i1.9466>.
- Khadka K, Khanal S, Dhakal M, Buchar S, Bisht S.** 2021. A Socioeconomic Survey Report on Springs in Phulbari Water Tower, Ward 2, Namobuddha Municipality, Kavrepalanchok District, Nepal. Kathmandu, Nepal: ICIMOD [International Centre for Integrated Mountain Development]; accessed on 15 July 2023.
- Khadka K, Rijal ML.** 2020. Hydrogeochemical assessment of spring water resources around Melamchi, central Nepal. *Water Practice and Technology* 15(3):748–758. <https://doi.org/10.2166/wpt.2020.066>.
- Khatiwada N, Bohara AK, Kunwar SB.** 2023. Household preferences for time and monetary contributions for river restoration: A study from the Danda River Basin in Nepal. *Water Resources and Economics* 43:100228. <https://doi.org/10.1016/j.wre.2023.100228>.
- Khatrri D, Neupane KR, Devkota K, editors.** 2021. *Dhulikhel's Journey towards Water Security*. Kathmandu, Nepal: Southasia Institute of Advanced Studies. https://sias-southasia.org/wp-content/uploads/2021/02/Book-Layout_final_forward.pdf; accessed on 20 July 2023.
- Kihila JM, Mtei K, Njau KN.** 2014. Development of a cost-benefit analysis approach for water reuse in irrigation. *International Journal of Environmental Protection and Policy* 2(5):179. <https://doi.org/10.11648/j.ijepp.20140205.16>.
- Komatsu S, Yamamoto Y, Ito Y, Kaneko S, Dhital RP.** 2020. Water for life: Ceaseless routine efforts for collecting drinking water in remote mountainous villages of Nepal. *Environment, Development and Sustainability* 22(8):7909–7925. <https://doi.org/10.1007/s10668-019-00552-9>.
- Koolwal G, Walle D.** 2016. *Access to Water, Women's Work and Child Outcomes*. The World Bank Policy Research Working Paper Report No. 5302. Washington, DC: World Bank. <https://documents.worldbank.org/en/publication/documents-reports/documentdetail/101601468336027439/access-to-water-womens-work-and-child-outcomes>; accessed on 28 July 2023.
- Krueger RA, Casey MA.** 2015. *Focus Groups: A Practical Guide for Applied Research*. 5th edition (1st edition 1994). Thousand Oaks, CA: Sage.
- Kumar M, Rathod R, Mukherji A.** 2023. Water security and spring conservation in the Himalaya. In: Ojha H, Schofield N, Camkin J, editors. *Climate Risks to Water Security: Framing Effective Response in Asia and the Pacific*. Cham, Switzerland: Springer, pp 15–36. https://doi.org/10.1007/978-3-031-16648-8_2.
- Kumar R.** 2020. Repercussion of environmental flows allocation in Ramganga basin plan-2045. *International Journal of Emerging Technology and Advanced Engineering* 10(4):54–65. <https://www.scopus.com/inward/record.uri?eid=2-s2.0-85103519311&partnerID=40&md5=d34f7d035276f2c0b45c871fb6a200a3>; accessed on 22 July 2023.
- Kumar V, Sen S.** 2018. Evaluation of spring discharge dynamics using recession curve analysis: A case study in data-scarce region, Lesser Himalayas, India. *Sustainable Water Resources Management* 4(3):539–557. <https://doi.org/10.1007/s40899-017-0138-z>.
- Lamichhane N, Dandekhya S, Shrestha K, Shrestha R.** 2020. *Springshed Research: A Study from Chorghare Village Development Committee in the Mid-hills of Nepal*. HI-AWARE Report No. 25. Kathmandu, Nepal: ICIMOD [International Centre for Integrated Mountain Development]. <https://lib.icimod.org/record/34933>; accessed on 13 June 2023.
- Lamsal P, Atreya K, Pant KP, Kumar L.** 2015. An analysis of willingness to pay for community-based conservation activities at the Ghodaghodi Lake Complex, Nepal. *International Journal of Biodiversity Science, Ecosystem Services & Management* 11(4):341–348. <https://doi.org/10.1080/21513732.2015.1055338>.
- Leder S, Clement F, Karki E.** 2017. Reframing women's empowerment in water security programmes in Western Nepal. *Gender & Development* 25(2):235–251. <https://doi.org/10.1080/13552074.2017.1335452>.
- Longworth B.** 2022. Individualized community-led drinking water supply projects in rural Nepal: A more effective and sustainable methodology than generalized approaches. *Crossings* 2(1):131–138. <https://doi.org/10.29173/crossings90>.
- Macchi M, Gurung AM, Hoermann B.** 2015. Community perceptions and responses to climate variability and change in the Himalayas. *Climate and Development* 7(5):414–425. <https://doi.org/10.1080/17565529.2014.966046>.
- Maurya AS, Shah M, Deshpande RD, Bhardwaj RM, Prasad A, Gupta SK.** 2011. Hydrograph separation and precipitation source identification using stable water isotopes and conductivity: River Ganga at Himalayan foothills. *Hydrological Processes* 25(10):1521–1530. <https://doi.org/10.1002/hyp.7912>.
- Mcharo M, Maghenda M.** 2021. Cost-benefit analysis of sustainable land and water management practices in selected highland water catchments of Kenya. *Scientific African* 12:e00779. <https://doi.org/10.1016/j.sciaf.2021.e00779>.
- Mendelsohn R, Olmstead S.** 2009. The economic valuation of environmental amenities and disamenities: Methods and applications. *Annual Review of Environment and Resources* 34(1):325–347. <https://doi.org/10.1146/annurev-environ-011509-1352011>.
- Mishra PK, Rai SC.** 2014. A cost-benefit analysis of indigenous soil and water conservation measures in Sikkim Himalaya, India. *Mountain Research and Development* 34(1):27–35. <https://doi.org/10.1659/MRD-JOURNAL-D-12-00013.1>.
- Mukherji A, Scott C, Molden D, Maharjan A.** 2018. Megatrends in Hindu Kush Himalaya: Climate change, urbanisation and migration and their implications for water, energy and food. In: Biswas AK, Tortajada C, Rohner P, editors. *Assessing Global Water Megatrends*. Singapore: Springer, pp 125–146. https://doi.org/10.1007/978-981-10-6695-5_8.

- Nankhuni FJ, Findeis JL.** 2004. Natural resource-collection work and children's schooling in Malawi. *Agricultural Economics* 31(2–3):123–134. <https://doi.org/10.1111/j.1574-0862.2004.tb00251.x>.
- Naudiyal N, Schmerbeck J.** 2017. The changing Himalayan landscape: Pine-oak forest dynamics and the supply of ecosystem services. *Journal of Forestry Research* 28(3):431–443. <https://doi.org/10.1007/s11676-016-0338-7>.
- Nauges C, Strand J.** 2017. Water hauling and girls' school attendance: Some new evidence from Ghana. *Environmental Resource Economics* 66(1):65–88. <https://doi.org/10.1007/s10640-015-9938-5>.
- Nepal M, Nepal AK, Berrens RP.** 2017. Where gathering firewood matters: Proximity and forest management effects in hedonic pricing models for rural Nepal. *Journal of Forest Economics* 27:28–37. <https://doi.org/10.1016/j.jfe.2017.02.005>.
- Nie Y, Pritchard HD, Liu Q, Hennig T, Wang W, Wang X, Liu S, Nepal S, Samyn D, Hewitt K, et al.** 2021. Glacial change and hydrological implications in the Himalaya and Karakoram. *Nature Reviews Earth & Environment* 2(2):91–106. <https://doi.org/10.1038/s43017-020-00124-w>.
- NITI Aayog.** 2018. *Inventory and Revival of Springs in the Himalayas for Water Security*. New Delhi, India: NITI Aayog.
- Nüsser M.** 2017. Socio-hydrology: A new perspective on mountain waterscapes at the nexus of natural and social processes. *Mountain Research and Development* 37(4):518–520. <https://doi.org/10.1659/MRD-JOURNAL-D-17-00101.1>.
- OECD [Organisation for Economic Cooperation and Development].** 2015. *Cost-Benefit Analysis and the Environment*. Report No. 97. Paris, France: OECD. https://www.oecd-ilibrary.org/environment/cost-benefit-analysis-and-the-environment_5jrp6w76tstg-en; accessed on 24 April 2023.
- Ojha H, Neupane KR, Pandey CL, Singh V, Bajracharya R, Dahal N.** 2020. Scarcity amidst plenty: Lower Himalayan cities struggling for water security. *Water* 12(2):567. <https://doi.org/10.3390/w12020567>.
- Pakhtigian EL, Jeuland M.** 2019. Valuing the environmental costs of local development: Evidence from households in western Nepal. *Ecological Economics* 158:158–167. <https://doi.org/10.1016/j.ecolecon.2018.12.021>.
- Pandey R, Aretano R, Gupta AK, Meena D, Kumar B, Alatalo JM.** 2017. Agroecology as a climate change adaptation strategy for smallholders of Tehri-Garhwal in the Indian Himalayan region. *Small-Scale Forestry* 16(1):53–63. <https://doi.org/10.1007/s11842-016-9342-1>.
- Pandit S, Shakya N, Shrestha SR.** 2019. Distribution and classification of springs in Bansbari area of Melamchi Municipality, Sindhupalchowk, Nepal. *Journal of Nepal Geological Society* 59:49–58. <https://doi.org/10.3126/jngs.v59i0.24985>.
- Pant KP.** 2009. Price-based valuation of rice genetic diversity in Nepal. Contributed paper prepared for presentation at the International Association of Agricultural Economists Conference, Beijing, China, August 16–22, 2009. <https://doi.org/10.22004/ag.econ.51642>.
- Picchioni F, Zanello G, Srinivasan CS, Wyatt AJ, Webb P.** 2020. Gender, time-use, and energy expenditures in rural communities in India and Nepal. *World Development* 136:105137. <https://doi.org/10.1016/j.worlddev.2020.105137>.
- Pokhrel G, Khadka K, Dhakal M.** 2019. Study of springs and springshed management in Namobuddha Municipality, Kavrepalchowk District. Report under review. Available from corresponding author of this article.
- Poudel DD.** 2020. Asta-Ja Framework: A peaceful approach to food, water, climate, and environmental security coupled with sustainable economic development and social inclusion in Nepal. *Strategic Planning for Energy and the Environment* 39(3–4):243–318. <https://doi.org/10.13052/spee1048-4236.391412>.
- Poudel DD, Duex TW.** 2017. Vanishing springs in Nepalese mountains: Assessment of water sources, farmers' perceptions, and climate change adaptation. *Mountain Research and Development* 37(1):35–46. <https://doi.org/10.1659/MRD-JOURNAL-D-16-00039.1>.
- Pouillot R, Delignette-Muller ML.** 2010. Evaluating variability and uncertainty separately in microbial quantitative risk assessment using two R packages. *International Journal of Food Microbiology* 142(3):330–340. <https://doi.org/10.1016/j.ijfoodmicro.2010.07.011>.
- Rathod R, Kumar M, Mukherji A, Sikka A, Satapathy KK, Mishra A, Goel S, Khan M.** 2021. *Resource Book on Springshed Management in the Indian Himalayan Region: Guidelines for Policy Makers and Development Practitioners*. New Delhi, India: IWMI [International Water Management Institute], NITI Aayog Government of India, and SDC [Swiss Agency for Development and Cooperation]. <https://www.iwmi.cgiar.org/Publications/Other/Reports/resource-book-on-springshed-management-in-the-indian-himalayan-region.pdf>; accessed on 29 August 2023.
- Ratnaweera D, Heistad A, Navrud S.** 2021. The current use and potential of cost benefit analysis in water sector projects. *Water Supply* 21(4):1438–1449. <https://doi.org/10.2166/ws.2020.364>.
- Raut M.** 2023. Gender and social inclusion in community water resource management: Lessons from two districts in the Himalayan foothills and the Terai in Nepal. *Water International* 48(4):547–566. <https://doi.org/10.1080/02508060.2023.2213962>.
- Rautela P.** 2015. Traditional practices of the people of Uttarakhand Himalaya in India and relevance of these in disaster risk reduction in present times. *International Journal of Disaster Risk Reduction* 13:281–290. <https://doi.org/10.1016/j.ijdrr.2015.07.004>.
- Roebeling P, Abrantes N, Ribeiro S, Almeida P.** 2016. Estimating cultural benefits from surface water status improvements in freshwater wetland ecosystems. *Science of the Total Environment* 545–546:219–226. <https://doi.org/10.1016/j.scitotenv.2015.12.063>.
- Sangaramoorthy T, Kroeger KA.** 2020. *Rapid Ethnographic Assessments: A Practical Approach and Toolkit for Collaborative Community Research*. London, United Kingdom: Routledge. <https://www.taylorfrancis.com/books/9781000001419>; accessed on 4 September 2023.
- Shea JM, Immerzeel WW, Wagnon P, Vincent C, Bajracharya S.** 2015. Modelling glacier change in the Everest region, Nepal Himalaya. *The Cryosphere* 9(3):1105–1128. <https://doi.org/10.5194/tc-9-1105-2015/>.
- Shlim DR, Hoge CW, Rajah R, Scott RM, Pandey P, Echeverria P.** 1999. Persistent high risk of diarrhea among foreigners in Nepal during the first 2 years of residence. *Clinical Infectious Diseases* 29(3):613–616. <https://doi.org/10.1086/598642>.
- Shrestha AB, Aryal R.** 2011. Climate change in Nepal and its impact on Himalayan glaciers. *Regional Environmental Change* 11(S1):65–77. <https://doi.org/10.1007/s10113-010-0174-9>.
- Shrestha AB, Desai J, Mukherji A, Dhakal M, Kulkarni, Mahamuni K, Buchar S, Bajracharya S.** 2018. *Protocol for Reviving Springs in the Hindu Kush Himalayas: A Practitioner's Manual*. Report No. 2018/4. Kathmandu, Nepal: ICIMOD [International Centre for Integrated Mountain Development].
- Shrestha G, Clement F.** 2019. Unravelling gendered practices in the public water sector in Nepal. *Water Policy* 21(5):1017–1033. <https://doi.org/10.2166/wp.2019.238>.
- Sim J.** 1998. Collecting and analysing qualitative data: Issues raised by the focus group. *Journal of Advanced Nursing* 28(2):345–352. <https://doi.org/10.1046/j.1365-2648.1998.00692.x>.
- Sivapalan M, Konar M, Srinivasan V, Chhatre A, Wutich A, Scott CA, Wescoat JL, Rodriguez-Iturbe I.** 2014. Socio-hydrology: Use-inspired water sustainability science for the Anthropocene. *Earth's Future* 2(4):225–230. <https://doi.org/10.1002/2013EF000164>.
- Smakhtin VU.** 2001. Low flow hydrology: A review. *Journal of Hydrology* 240(3–4):147–186. [https://doi.org/10.1016/S0022-1694\(00\)00340-1](https://doi.org/10.1016/S0022-1694(00)00340-1).
- Tambe S, Dhakal S, Dhakal Deepak, Sharma G, Sherpa PN, Kulkarni H, Bhutia NT, Dhakal Dinesh, Pradhan S, Sinha UK, et al.** 2020. Scaling up spring revival in the Himalaya: Graduating from spring-centric to aquifer-centric nature-based solutions. In: Dhyani S, Gupta AK, Karki M, editors. *Nature-Based Solutions for Resilient Ecosystems and Societies*. Singapore: Springer, pp 29–50. https://doi.org/10.1007/978-981-15-4712-6_2.
- Tambe S, Kharel G, Arrawatia ML, Kulkarni H, Mahamuni K, Ganeriwala AK.** 2012. Reviving dying springs: Climate change adaptation experiments from the Sikkim Himalaya. *Mountain Research and Development* 32(1):62–72. <https://doi.org/10.1659/MRD-JOURNAL-D-11-00079.1>.
- Tarafdar S.** 2013. Understanding the dynamics of high and low spring flow: A key to managing the water resources in a small urbanized hillslope of Lesser Himalaya, India. *Environmental Earth Sciences* 70(5):2107–2114. <https://doi.org/10.1007/s12665-011-1493-y>.
- Taye FA, Foltkes MV, Fleming CM, Buckwell A, Mackey B, Diwakar KC, Le D, Hasan S, Ange CS.** 2021. The economic values of global forest ecosystem services: A meta-analysis. *Ecological Economics* 189:107145. <https://doi.org/10.1016/j.ecolecon.2021.107145>.
- Tenge AJ, de Graffe J, Hella JP.** 2005. Financial efficiency of major soil and water conservation measures in West Usambara highlands, Tanzania. *Applied Geography* 25(4):348–366. <https://doi.org/10.1016/j.apgeog.2005.08.003>.
- Thakur JK, Neupane M, Mohanan AA.** 2017. Water poverty in upper Bagmati River Basin in Nepal. *Water Science* 31(1):93–108. <https://doi.org/10.1016/j.wsj.2016.12.001>.
- Thapa B, Bhattarai C, Dahal N, Tiwari S, Jacobsen D.** 2023. Drying of springs in the Himalayan region of Nepal: Perspectives of local government leaders on causes, consequences, and conservation efforts. *Mountain Research and Development* 43(4):R9–R15. <https://doi.org/10.1659/mrd.2023.00007>.
- Thapa B, Pant RR, Thakuri S, Pond G.** 2020. Assessment of spring water quality in Jhimruk River Watershed, Lesser Himalaya, Nepal. *Environmental Earth Science* 79(22):504. <https://doi.org/10.1007/s12665-020-09252-4>.
- Thapa S, Wang L, Koira A, Shrestha S, Bhattarai S, Aye WN.** 2020. Valuation of ecosystem services from an important wetland of Nepal: A study from Begnas watershed system. *Wetlands* 40(5):1071–1083. <https://doi.org/10.1007/s13157-020-01303-7>.
- Tiwari PC.** 2000. Land-use changes in Himalaya and their impact on the plains ecosystem: Need for sustainable land use. *Land Use Policy* 17(2):101–111. [https://doi.org/10.1016/S0264-8377\(00\)00002-8](https://doi.org/10.1016/S0264-8377(00)00002-8).
- Troy TJ, Pavao-Zuckerman M, Evans TP.** 2015. Debates—Perspectives on socio-hydrology: Socio-hydrologic modeling: Tradeoffs, hypothesis testing, and validation. *Water Resources Research* 51(6):4806–4814. <https://doi.org/10.1002/2015WR017046>.
- Upreti BR.** 2001. *Conflict Management in Natural Resources, A Study of Land, Water and Forest Conflicts in Nepal* [PhD dissertation]. Wageningen, the Netherlands: Wageningen University.
- Upreti BR, Dierolf TS, Krain E, Kramer E, Tarmudji MS, Nasution A.** 1999. *Managing Local Conflicts over Water Resources: A Case Study from Nepal*. Network Paper No. 95. London, United Kingdom: Agricultural Research and Extension Network.
- Valdiya KS, Bartarya SK.** 1991. Hydrogeological studies of springs in the catchment of the Gaula River, Kumaun Lesser Himalaya, India. *Mountain Research and Development* 11(3):239–258. <https://doi.org/10.2307/3673618>.

- Verma R, Jamwal P.** 2022. Sustenance of Himalayan springs in an emerging water crisis. *Environmental Monitoring and Assessment* 194(2):87. <https://doi.org/10.1007/s10661-021-09731-6>.
- Wali N, Georgeou N, Simmons O, Gautam MS, Gurung S.** 2020. Women and WASH in Nepal: A scoping review of existing literature. *Water International* 45(3):222–245. <https://doi.org/10.1080/02508060.2020.1754564>.
- Wester P, Mishra A, Mukherji A, Shrestha AB, editors.** 2019. *The Hindu Kush Himalaya Assessment: Mountains, Climate Change, Sustainability and People*. Cham, Switzerland: Springer. <https://doi.org/10.1007/978-3-319-92288-1>.
- Whittington D, Cook J.** 2019. Valuing changes in time use in low- and middle-income countries. *Journal of Benefit-Cost Analysis* 10(S1):51–72. <https://doi.org/10.1017/bca.2018.21>.
- Williams MW, Wilson A, Tshering D, Thapa P, Kayastha RB.** 2016. Using geochemical and isotopic chemistry to evaluate glacier melt contributions to the Chamkar Chhu (river), Bhutan. *Annals of Glaciology* 57(71):339–348. <https://doi.org/10.3189/2016AoG71A068>.
- Wilson AM, Williams MW, Kayastha RB, Racoviteanu A.** 2016. Use of a hydrologic mixing model to examine the roles of meltwater, precipitation and groundwater in the Langtang River basin, Nepal. *Annals of Glaciology* 57(71):155–168. <https://doi.org/10.3189/2016AoG71A067>.
- Wood LR, Neumann K, Nicholson KN, Bird BW, Dowling CB, Sharma S.** 2020. Melting Himalayan glaciers threaten domestic water resources in the Mount Everest region, Nepal. *Frontiers in Earth Science* 8:128. <https://doi.org/10.3389/feart.2020.00128>.
- Yao Y, Zheng C, Andrews CB, Scanlon BR, Kuang X, Zeng Z, Jeong S, Lancia M, Wu Y, Li G.** 2021. Role of groundwater in sustaining northern Himalayan rivers. *Geophysical Research Letters* 48(10):e2020GL092354. <https://doi.org/10.1029/2020GL092354>.
- Yokying P, Saksena S, Fox J.** 2023. Impacts of migration on time allocation of those who remain at home in rural Nepal. *Journal of International Development* 35(7):2067–2106. <https://doi.org/10.1002/jid.3765>.