



Sexual Identification of *Caiman latirostris* Hatchlings by Cloacal Inspection

Authors: Ciocan, Hernán, Leiva, Pamela M.L., and Simoncini, Melina S.

Source: South American Journal of Herpetology, 16(1) : 50-55

Published By: Brazilian Society of Herpetology

URL: <https://doi.org/10.2994/SAJH-D-18-00075.1>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Sexual Identification of *Caiman latirostris* Hatchlings by Cloacal Inspection

Hernán Ciocan^{1,2,3*}, Pamela M.L. Leiva^{1,3}, Melina S. Simoncini^{1,3}

¹ Proyecto Yacaré, Laboratorio de Zoología Aplicada: Anexo Vertebrados (Facultad de Humanidades y Ciencias, Universidad Nacional del Litoral/ Ministerio de Medio Ambiente), Aristóbulo del Valle 8700, CP 3000 Santa Fe, Santa Fe, Argentina.

² Ministerio de Medio Ambiente de Santa Fe, Dirección General de Manejo Sustentable de Fauna, Patricio Cullen 6161, CP S3004IYC, Santa Fe, Argentina.

³ Centro de Investigación Científica y de Transferencia Tecnológica a la Producción (Consejo Nacional de Investigaciones Científicas y Técnicas, Provincia Entre Ríos, Universidad Autónoma de Entre Ríos). España 149, CP 3105 Diamante, Entre Ríos, Argentina.

* Corresponding author. Email: hernanciocan@gmail.com

Abstract. Sexual identification of crocodylians is important in population studies and provides useful information for conservation and management plans and monitoring populations over time. It is possible to distinguish between male and female *Caiman latirostris* by cloacal palpation or eversion of the penis in individuals larger than 75 cm total length, but smaller animals possess a barely differentiable cliteropenis. In hatchlings, sex determination methods involve surgical examination, necropsy, or analysis of cranial dimorphism, which cannot be applied in the field. We classified hatchlings of *C. latirostris* by observing the color and shape of their genitals. The penis is a milky white organ with a rounded shape at the tip and a purple hue at the end, whereas the clitoris is shorter, whitish, and has a pointed end. The procedure was tested on hatchlings from three nests at the “Proyecto Yacaré” study area (Santa Fe province); half of the eggs of each nest were incubated at a constant temperature of 31°C (producing females) and the other half at 33°C (producing males). To observe the sexual organs by cloacal inspection, we used a modified instrument whose function during palpation is like that of a finger applied in large animals to evert the penis or clitoris. In the first days after hatching we correctly scored the sex of 80% of the individuals. The number of correct identifications was slightly lower for males than for females. This technique might be a useful tool for field studies, as it allows the sex of small caimans to be estimated in situ.

Keywords. Broad Snouted Caiman; Cliteropenis; Cloacal sexing; Genital morphology; Sex identification; Sex ratio.

Resumen. La identificación del sexo de los cocodrilos es de importancia en estudios sobre ecología de poblaciones, ya que este dato permite elaborar o proponer planes de conservación y manejo, así como también, el seguimiento de estas poblaciones en el tiempo. En la naturaleza, es posible diferenciar machos y hembras de *Caiman latirostris*, mediante la palpación cloacal o eversión del pene en animales de tamaño mayor a 75 cm de largo total. Mientras que en animales pequeños se observa un cliteropene difícilmente diferenciable. En el caso de los neonatos, los métodos de determinación del sexo implican exámenes quirúrgicos, necropsias o análisis de dimorfismo craneal, los cuales no son aplicables a las condiciones de muestreo en campo. En este trabajo, clasificamos ejemplares recién nacidos de *C. latirostris*, observando el color y forma de sus genitales. Pudimos caracterizar al pene como un órgano de color blanco lechoso con forma redondeada en la punta y una tonalidad púrpura en el extremo; mientras que el clitoris es más corto, blanquecino y con el extremo puntiagudo. El procedimiento se testeó en crías de tres nidos, provenientes de sitios de trabajo de “Proyecto Yacaré” (Provincia de Santa Fe); la mitad de los huevos de cada uno fueron incubados a una temperatura constante de 31°C (productora de hembras) y la otra mitad a 33°C (productora de machos). Para observar los órganos sexuales mediante inspección cloacal, utilizamos un instrumento que modificamos y cuya función es similar a la de un dedo, como frecuentemente se utiliza en animales grandes para realizar la eversión del pene o clitoris. De acuerdo con nuestras observaciones, en los primeros días de vida asignamos correctamente el sexo del 80% de los individuos. La cantidad de aciertos fue ligeramente menor para los machos que para las hembras. Esta técnica sería una herramienta de utilidad para los estudios de campo, ya que posibilita la identificación del sexo de animales pequeños in situ.

INTRODUCTION

All species of crocodylians undergo temperature-dependent sex determination (Lang and Andrews, 1994), with the temperatures that produce males or females varying according to the species (Janzen and Paukstis, 1991; Valenzuela and Lance, 2004; Piña et al., 2015). In nature, nest temperature cannot be used to deduce hatchling gender because researchers are unaware of the temperatures at which the embryos developed, and these temperatures

vary due to the effect of environmental conditions or nesting habitat (Campos, 1993; Lang and Andrews, 1994; Simoncini et al., 2014).

Many crocodile species, including *Crocodylus acutus* (Cuvier, 1807), *Crocodylus moreletii* (Duméril and Bibron, 1851), and *Crocodylus porosus* Schneider, 1801, show sexual size dimorphism when mature, and it is possible to identify males when they exceed the maximum size of females; however, males cannot be distinguished from females of the same size (Platt et al., 2009; Platt et al.,

How to cite this article: Ciocan H., Leiva P.M.L., Simoncini M.S. 2020. Sexual identification of *Caiman latirostris* hatchlings by cloacal inspection. *South American Journal of Herpetology* 16: 50–55. <http://doi.org/10.2994/SAJH-D-18-00075.1>

Submitted: 06 November 2018
Accepted: 11 September 2019
Available Online: 31 August 2020

Handling Editor: Luciano Martins Verdade
<http://doi.org/10.2994/SAJH-D-18-00075.1>

2011; Barrios-Quiroz et al., 2012; Fukuda et al., 2013). Similarly, individuals can be sexed by cloacal palpation or eversion of the penis once they reach a total length greater than approximately 60 cm (Chabreck, 1963; Webb et al., 1984) or 75 cm (Honegger, 1978; Combrink et al., 2012). Studies on *Alligator mississippiensis* (Daudin, 1802) and *Caiman latirostris* (Daudin, 1802) report that hatchlings possess an undifferentiable cliteropenis (Joanen and McNease, 1978; Ziegler and Olbort, 2007; Nuñez-Otaño et al., 2010); in contrast, in other species of crocodylians the hatchling penis and the clitoris can be differentiated, as for example in *Caiman crocodilus* (Linnaeus, 1758) and several species of *Crocodylus porosus* and *Crocodylus johnstoni* Krefft, 1873 (Webb et al., 1984), *Crocodylus niloticus* Laurenti, 1768 (Hutton, 1987), and *Crocodylus palustris* (Lesson, 1831) (Lang et al., 1989; Lang and Andrews, 1994).

The penis of crocodylians is normally hidden within the cloaca (Ziegler and Olbort, 2007; Johnston et al., 2014), and its size depends on the species and the size of the animal (Moore and Kelly, 2015). It is usually cylindrical, somewhat laterally compressed, and reinforced by connective tissue (Ziegler and Olbort, 2007; Moore et al., 2012; Kelly, 2013). Protrusion during copulation is produced by muscular pressure and vascular dilatation (Ziegler and Olbort, 2007; Moore et al., 2012), and the penis has a dorsal groove that functions as a spermatic channel (Cabrera and García, 2007; Moore and Kelly, 2015); this groove is also found in the clitoris of females, but it lacks any known function (Ziegler and Olbort, 2007). At the end of the genital organ, a loop-like structure is observed, called the tip, on to which the spermatic groove extends, and a softer structure is demarcated from the tip by a small groove. For mating, the penis protrudes forward (cephalad) from the cloaca and is bent towards the venter (Ziegler and Olbort, 2007; Johnston et al., 2014).

The cloacal palpation method used to evaluate the sex of crocodylians usually consists of introducing a finger or speculum in the cloaca and everting the penis (Chabreck, 1967; Brazaitis, 1969). However, this method can only be used in animals larger than about 75 cm in total length (Honegger, 1978; Ziegler and Olbort, 2007; Combrink et al., 2012), whose minimum cloacal size allows the insertion of a finger. In the case of smaller specimens, a speculum can be used to observe differences in external genital morphology, although it is difficult to apply (Hutton, 1984; Allsteadt and Lang, 1995; Leslie, 1997; Combrink et al., 2012). Another method proposed by Whitaker et al. (1980) consists of everting the penis by compressing the cloaca laterally and flexing the tail towards the ventral part. Differentiation of genital structures occurs at very small sizes in some species and at sizes of at least 1 m in others (e.g., *Gavialis gangeticus* [Gmelin, 1789], Lal and Basu, 1982). Dissection of *Alligator mississippiensis* and *Caiman latirostris* hatchlings have revealed that the penis is

long and has a rounded shape and reddish coloration, while the clitoris is short and pale white (Allsteadt and Lang, 1995; Nuñez-Otaño et al., 2010; Gredler et al., 2015).

In the case of neonates, palpation is not possible due to their small size and undifferentiated penis and clitoris (Ziegler and Olbort, 2007; Nuñez-Otaño et al., 2010). Therefore, for the evaluation of sex in neonates, invasive methods such as surgical examination of the gonads (Magnusson et al., 1990) or necropsy are applied (Nuñez-Otaño et al., 2010; Murray et al., 2016). Cranial dimorphism can also be analyzed, but it has an efficacy of approximately 70% and is not applicable in the field (Piña et al., 2007a), or one must for the animals to grow, which requires a lot of time (Piña et al., 2007b).

The ability to identify the sex of crocodylian hatchlings and small juveniles in the field would provide an important tool to obtaining data without transporting or collecting the animals or impacting the study population. This is important because sex identification is fundamental to understand population dynamics and ecological/biological processes, and knowing the proportions of males and females is essential for management plans, either for conservation or harvest (Ziegler and Olbort, 2007; Platt et al., 2009; Fukuda et al., 2013). Consequently, the objective of this study is to validate a technique that allows sex identification of *Caiman latirostris* hatchlings through observation with the naked eye. We also test if the results are affected by the size of the hatchlings and the nest of origin.

MATERIALS AND METHODS

We collected 100 eggs of *Caiman latirostris* from three nests collected in the localities of Cacique Ariacaiquín, Espín, and Intiyaco, which are areas of the “Proyecto Yacaré,” a sustainable use and conservation program (Leiva et al., 2019). Once the nests were collected and transferred to the laboratory, we determined the embryonic development time of the nest according to the criteria of Lungman et al. (2008). This was done by observing the opaque band (no greater than 37% of egg surface) and the morphological characteristics of the embryo. We only used nests at a developmental stage less than or equal to 5 d (stage 5). All caimans of the three nests hatched on 7–8 March 2016. We randomly separated the eggs of each nest into two equal groups (50 each) and incubated them at 31°C (which produces a 100% female development) and 33°C (to produce 100% male development; Piña et al., 2007b; Parachú-Marcó et al., 2017). After birth, we identified the animals individually by scute clipping (Richardson et al., 2002).

We carried out four measurement events in which we considered different morphometric parameters: total length (TL), snout–vent length (SVL), and weight (W).

The first measurement and observation were carried out at the month of birth; thereafter, measurements and observations were made every 2 months until the hatchlings reached 7 months of age. We considered ranges of size of the animals to observe the sex according to the TL; these size intervals in total length (cm) were: 22–27, > 27–33, > 33–36+. Hatchlings that remained at the same size interval at the next measurement period were re-examined and their sex re-evaluated. However, the second measurement and sex identity were not included in the analysis. If the second sex identity was the same as the first, we classified it as successful; if not, we classified it as failure, regardless of which identification was confirmed by necropsy (see below).

For data collection, we obtained hatchlings from unidentified plastic trays and measured and weighed each individual. Later, the same experienced observer identified the sex—not knowing the mark, nest of origin, or incubation temperature—throughout the experiment. To observe the sexual organs by cloacal inspection, we used an instrument that we modified and whose function during palpation is like a finger applied in large animals to perform the eversion of the penis or clitoris. This instrument consists of a curved forceps (London College Tweezer) covered with latex to increase adhesion and prevent injury (Fig. 1). However, it can also be built with other materials, and we have also constructed versions made of wood and aluminum covered with latex. At the end of the study, we corroborated the sex of the animals via direct observation of their gonads by necropsy (Nuñez-Otaño et al., 2010; Murray et al., 2016) by randomly selecting 10% of each group to confirm that the hatchlings from eggs incubated at 31°C were females and those incubated at 33°C were males.

We analyzed the data using the Chi-square test, considering successes and failures in sex identification for each TL interval; this analysis was performed for each sex separately and with all individuals pooled. We considered no differences in the identification between size intervals studied if $P > 0.05$. For each growth interval, the reliability of the method was also calculated as successes/total individuals. Further, we evaluated if there was an effect of the nest of origin on the effectiveness of the identification in the animals whose origin was known.

RESULTS

Among the 100 incubated eggs, hatching success was 76%, including 41 females and 35 males (i.e., 54% F; 46% M). The observer identified males by the presence of a longer sexual organ with a broad, rounded tip and milky whitish coloration with a purple hue on the tip (Fig. 2). In contrast, females had a shorter clitoris with a fine tip termination and whitish coloration (Fig. 3).

We did not observe a size effect in the correct assignment of the sex of individuals in general ($X^2 = 0.74$, $df = 2$; $P = 0.6896$) or for males ($X^2 = 0.97$, $df = 2$; $P = 0.6146$) or females ($X^2 = 1.09$, $df = 2$, $P = 0.5785$). The Chi-square was calculated to observe if there was influence of the origin nest in the identification of sex, and we did not find differences between the different nests ($X^2 = 0.43$, $df = 2$; $P = 0.8049$).

DISCUSSION

According to Leslie (1997), the use of the cloacal test to identify sex is not reliable in *Crocodylus niloticus* from the Lake St. Lucia region during the first year of development. At approximately 14 months, the morphological differences between the male and female genitalia, mainly the length and shape of the cliteropenis, were more evident and enabled external sex identification. However, that study did not mention the size of those animals at 14 months. In contrast, Hutton (1984) was able to identify the sex of small 4-week-old *C. niloticus* with 80–85% effectiveness, which increased to 95% in animals larger than 40 cm TL. In our study, we achieved a success rate of 80.5% (71.4% for males and 87.5% for females) for individuals that were 4 weeks old and 22–27 cm TL—approximately half the size reported by Hutton (1984). Although we did not observe statistically significant differences in reliability according to the increase in size of the sexed individuals, the reliability was 88.9% (75.0% in males and 92.8% in females) for individuals of 33–36+cm TL. It is worth mentioning that the caimans in this study reached the last interval of TL at approximately 7 months under captive conditions (held at a controlled temperature of 32°C and fed three times per week).

For individuals of *Crocodylus porosus* and *Crocodylus johnstoni* of 10–14 cm SVL (~20–30 cm TL), the penis and clitoris can be distinguished with 95% accuracy using forceps (Webb et al., 1984). As reported by Webb et al. (1984), there is a bias in sex identification, confusing the long clitoris of females with penises, although it is rare for males to be classified as females. In contrast, in our study errors occurred when a small penis was mistaken for a clitoris, resulting in males being misclassified as females (Tables 1, 2). According to Moore et al. (2018), there are morphological differences between the sex organs of crocodiles and alligators. These differences could be observed in hatchlings, whereby it would be easier to identify sex in crocodiles, whose genital structures are more prominent than those of alligators. However, we were able to obtain a high percentage of successes in *Caiman latirostris*, although we did not test it in other alligatorids; therefore, we recommend that the tests should be extended to other species. According to Joanen and McNease (1978), *Alligator mississippiensis* individuals less




Figure 1. Curved forceps, closed at the end and lined with latex, used like a finger to evert the cliteropenis. Total length: 165 mm. Width from the rubber tip to elbow: 2–4 mm. Scale bar = 2 cm.




Figure 2. Penis with the tip retracted (A) and penis with the tip deployed (B). The milky white coloration with purple color/tint hue at the end is observed. The instrument used to make eversion is observed behind. Scale bar = 5 mm.



Figure 3. Clitoris. The whitish coloration and the sharpest termination can be observed. Behind it is the instrument used to evert the clitoris. Scale bar = 5 mm.

than 30 cm SVL (~60 cm TL) possess and undifferentiated cliteropenis; nevertheless, Allsteadt and Lang (1995) were able to classify the sex of hatchlings and juveniles of *A. mississippiensis* with the naked eye with a high percentage of effectiveness. Other studies have found morphological differences in alligatorids, but through necropsies or histological observations (Nuñez-Otaño et al., 2010; Kelly, 2013; Gredler et al., 2015).

Previous studies have claimed that sex could be identified in most crocodylian species through direct observation of individuals greater than 60 cm (Chabreck, 1963; Webb et al., 1984) or 75 cm TL (Honegger, 1978; Com-

brink et al., 2012). In the particular case of *Caiman latirostris*, to date sex identification has been recorded only in animals between 60–120 cm TL (Piña et al., 2007b; Nuñez-Otaño et al., 2010). In this species, although the penis is wider and has a greater volume than the clitoris, this was only observed after euthanasia of the hatchlings and analysis of photographs taken using a stereoscopic binocular microscope (Nuñez-Otaño et al., 2010). Through this method of observation, Nuñez-Otaño et al. (2010) achieved approximately 85% accuracy for hatchlings, which is similar to the results obtained in our study by direct observation of live animals.

We did not find statistically significant differences in the number of successes with respect to the size of the animals in the intervals evaluated. Since the nest effect is reflected in hatchlings size, it was expected that it would not influence the identification of hatchling sex (Garnett and Murray, 1986; Hutton, 1987; Schulte and Chabreck, 1990; Ciocan et al., 2018). Regardless, we do not dismiss the possibility that there could be differences in the sexual organ development associated with the nest of origin. However, reliability seems to increase as the size of the animal increases, reaching almost 90% for both sexes (Table 3).

Table 1. Male assessment, differentiated by individual size intervals (total length). *n* = No. individuals sexed in the corresponding size range.

Males				
Total length (cm)	<i>n</i>	Successes	Failures	Reliability
22–27	56	40	16	0.714
27–33	36	29	7	0.806
33–36+	4	3	1	0.75
Mean ± SD				0.757 ± 0.046

Table 2. Female assessment, differentiated by individual size intervals (total length). *n* = No. individuals sexed in the corresponding size range.

Females				
Total length (cm)	<i>n</i>	Successes	Failures	Reliability
22–27	72	63	9	0.875
27–33	40	33	7	0.825
33–36+	14	13	1	0.928
Mean ± SD				0.876 ± 0.052

Table 3. Assessment of both sexes, differentiated by individual size intervals (total length). *n* = No. individuals sexed in the corresponding size range.

Males and females				
Total length (cm)	<i>n</i>	Successes	Failures	Reliability
22–27	128	103	25	0.804
27–33	76	62	14	0.815
33–36+	18	16	2	0.889
Mean ± SD				0.836 ± 0.046

Previous studies affirm that the size of the cliteropenis of neonates of *Alligator mississippiensis* increases according to incubation temperature among animals of the same and different sexes (Allsteadt and Lang, 1995; Gredler et al., 2015). In the present study, we worked with animals from eggs incubated at temperatures producing only males (33°C) or only females (31°C), but it is possible that *Caiman latirostris* individuals incubated at temperatures producing both males and females (e.g., 32°C; Parachú-Marco et al., 2017), external sexual morphology might be more difficult to distinguish. Given that incubation temperatures in nature vary due to environmental conditions (Simoncini et al., 2014), the results of our method of sex identification could also vary.

We can conclude that it is possible to observe and distinguish between the male and genital structure of *Caiman latirostris* at an early age and small size using the cloacal palpation method and this new tool to perform the eversion. The method is simple, inexpensive, and easy to use in field conditions, without requiring the complex logistics required to study cranial morphometry or resorting to highly invasive practices such as surgical examination or necropsy. We also found that accuracy increased as the size of the hatchlings increased. For all individuals the accuracy was 84%, being slightly higher for females (88%)

than males (76%). The percentage of failures might vary according to the experience of the observer, who should understand the method and be able to clearly distinguish between male and female genitalia. Through this study, we considerably reduced the size in which a hatchling can be classified as male or female with a high effectiveness, which allows to expand the database of sexual proportion in small animals in the field.

ACKNOWLEDGMENTS

We would like to thank CICYTTP-CONICET for support. We are grateful to all members of Proyecto Yacaré, Dr. Carlos Piña for his advice, Msc. Ricardo Ojeda and Lic. Leonardo Scarpa for their help, and Dr. Perran Ross for help and advice. Special thanks to CSG-IUCN members for their contribution.

REFERENCES

- Allsteadt J., Lang J.W. 1995. Sexual dimorphism in the genital morphology of young American alligators, *Alligator mississippiensis*. *Herpetologica* 51:314–325.
- Barrios-Quiroz G., Casas-Andreu G., Escobedo-Galván A.H. 2012. Sexual size dimorphism and allometric growth of Morelet's crocodiles in captivity. *Zoological Science* 29:198–203. DOI
- Brazaitis P.J. 1969. The determination of sex in living crocodylians. *Herpetological Journal* 4:54–58.
- Cabrera F.J., García G.C. 2007. Mucinas of the male genitalia of the Baba (*Caiman crocodilus crocodilus*). *Revista de la Facultad de Ciencias-Veterinarias* 48:77–84.
- Campos Z. 1993. Effect of habitat on survival of eggs and sex ratio of hatchlings of *Caiman crocodilus yacare* in the Pantanal, Brazil. *Journal of Herpetology* 27:127–132. DOI
- Chabreck R.H. 1963. Methods of capturing, marking and sexing alligators. Pp. 47–50, in Proceedings of the Annual Conference of the Southeastern Association of Game and Fish Commissioners, Jefferson City.
- Chabreck R.H. 1967. Alligator farming hints. Pp. 1–21, in Wildlife and Fisheries Commission Report, Grand Chenier.
- Ciocan H., Imhof A., Parachú-Marcó M.V., Isberg S.R., Siroski P.A., Larriera A. 2018. Increasing photoperiod enhances growth in captive hatchling *Caiman latirostris*. *Aquaculture* 482:193–196. DOI
- Combrink X., Warner J.K., Hofmeyr M., Govender D., Ferreira S.M. 2012. Standard operating procedure for the monitoring, capture and sampling of Nile crocodiles (*Crocodylus niloticus*). Pp. 1–14, in Swan G. (Ed.), South African National Parks, Skukuza.
- Cuvier G. 1807. Sur les différentes espèces de crocodiles vivans et sur leurs caractères distinctifs. *Annales Muséum National D'Histoire Naturelle, par les professeurs de cet établissement* 10:8–86.
- Daudin F.M. 1802. Histoire Naturelle, Générale et Particulière des Reptiles; ouvrage faisant suite à l'Histoire naturelle générale et particulière, composée par Leclerc de Buffon; et rédigée par C.S. Sonnini, membre de plusieurs sociétés savantes, Tome 2. F. Dufart, Paris. DOI
- Duméril A., Bibron G. 1851. Sauriens ou lézards. Pp. 26–197, in Duméril M.C., Duméril A. (Eds.), Catalogue Méthodique de la Collection des Reptiles. Gide et Baudry, Paris.
- Fukuda Y., Saalfeld K., Lindner G., Nichols T. 2013. Estimation of total length from head length of Saltwater crocodiles (*Crocodylus porosus*) in the Northern territory, Australia. *Journal of Herpetology* 47:34–40. DOI

- Garnett S., Murray R.M. 1986.** Parameters affecting the growth of the estuarine crocodile, *Crocodylus porosus*, in captivity. *Australian Journal of Zoology* 34:221–223. [DOI](#)
- Gmelin J.F. 1789.** Caroli a Linné systema naturae per regna tria naturae, secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis. Tomus I. (ed. 13). Laurentii, Salvii, Holmiae. [DOI](#)
- Gredler M.L., Seifert A.W., Cohn M.J. 2015.** Morphogenesis and patterning of the phallus and cloaca in the American alligator, *Alligator mississippiensis*. *Sexual Development* 9:53–67. [DOI](#)
- Hogger R.E. 1978.** Sex identification in reptiles. *Salamandra* 14:69–79.
- Hutton J.M. 1984.** The Population Ecology of the Nile Crocodile, *Crocodylus niloticus* Laurenti, 1768, at Ngezi, Zimbabwe. Ph.D. Thesis, University of Zimbabwe, Zimbabwe.
- Hutton J.M. 1987.** Incubation temperatures, sex ratios, and sex determination in a population of Nile crocodiles (*Crocodylus niloticus*). *Journal of Zoology* 211:143–155. [DOI](#)
- Iungman J., Piña C.I., Siroski P. 2008.** Embryological development of *Caiman latirostris* (Crocodylia: Alligatoridae). *Genesis* 46:401–417. [DOI](#)
- Janzen F.J., Paukstis G.L. 1991.** Environmental sex determination in reptiles: ecology, evolution, and experimental design. *The Quarterly Review of Biology* 66:149–179. [DOI](#)
- Joanen T., McNease L. 1978.** The cloaca sexing method for immature alligators. *Proceedings of the Annual Conference of the Southeastern Association of Fish and Wildlife Agencies, Oklahoma* 32:179–181.
- Johnston S.D., Lever J., McLeod R., Oishi M., Qualischefski E., Omanga C., ... D'Occhio M. 2014.** Semen collection and seminal characteristics of the Australian saltwater crocodile (*Crocodylus porosus*). *Aquaculture* 422–423:25–35. [DOI](#)
- Kelly D.A. 2013.** Penile anatomy and hypotheses of erectile function in the American alligator (*Alligator mississippiensis*): muscular eversion and elastic retraction. *The Anatomical Record* 296:488–494. [DOI](#)
- Lal S., Basu D. 1982.** Sex and sex ratios of gharial (*Gavialis gangeticus*) raised in captivity. *Journal Bombay Natural History Society* 79:688–691.
- Lang J.W., Andrews H.V. 1994.** Temperature-dependent sex determination in crocodylians. *Journal Experimental Zoology* 270:28–44. [DOI](#)
- Lang J.W., Andrews H., Whitaker R. 1989.** Sex determination and sex ratios in *Crocodylus palustris*. *American Zoologist* 29:935–952. [DOI](#)
- Laurenti J.N. 1768.** Specimen medicum, exhibens synopsis reptilium emendatam cum experimentis circa venena et antidota reptilium austriacorum. Joan. Thom. Nob. de Trattren, Viennae. [DOI](#)
- Leiva P.M.L., Simoncini M.S., Portelinha T.C.G., Larriera A., Piña C.I. 2019.** Size of nesting female broad-snouted caimans (*Caiman latirostris* Daudin 1802). *Brazilian Journal of Biology* 79:139–143. [DOI](#)
- Leslie A. 1997.** The Ecology and Physiology of the Nile Crocodile, *Crocodylus niloticus*, in Lake St Lucia KwaZulu Natal, South Africa. Ph.D. Thesis, Drexel University, United States.
- Lesson R.-P. 1834.** Reptiles. Pp. 291–336, in Bélanger I.G.S. (Ed.), Voyage aux Indes-Orientales, par le Nord de l'Europe: Les Provinces du Caucase, la Géorgie, l'Arménie Et la Perse, Suivi de Détails Topographiques Et Autres sur le Pégou, les Îles de Java, de Maurice Et de Bourbon, sur le Cap de Bonne-Espérance Et Sainte-Hélène, Pendant les Années 1825, 1826, 1827, 1828 et 1829. Zoologie. A. Bertrand, Paris.
- Linnaeus C. 1758.** Systema naturae per regna tria naturae, secundum classes, ordines, genera, species, cum characteribus, differentiali, synonymis, locis, Tomus I. Editio decima, reformata. Laurentii Salvii, Holmiae. [DOI](#)
- Magnusson W.E., Lima A.P., Hero J.M., Sanaiotii T.M., Yamakoshi M. 1990.** *Paleosuchus trigonatus* nests: sources of heat and embryo sex ratios. *Journal of Herpetology* 24:397–400. [DOI](#)
- Moore B.C., Kelly D.A. 2015.** Histological investigation of the adult alligator phallic sulcus. *South American Journal of Herpetology* 10:32–40. [DOI](#)
- Moore B.C., Mathavan K., Guillette L.J. Jr. 2012.** Morphology and histochemistry of juvenile male American alligator (*Alligator mississippiensis*) phallus. *The Anatomical Record* 295:328–337. [DOI](#)
- Moore B.C., Myburgh J., Woodward A., Penland S., Francis R., Kelly D. 2018.** Comparing the male copulatory anatomy of American alligator and Nile crocodile: Differing forms, differing functions? Pp. 37–38, in Proceeding 25th Working Meeting of the Crocodile Specialist Group, Santa Fe.
- Murray C.M., Easter M., Merchant M., Rheubert J.L., Wilson K.A., Cooper A., ... Guyer C. 2016.** Methyltestosterone alters sex determination in the American alligator (*Alligator mississippiensis*). *General and Comparative Endocrinology* 236:63–69. [DOI](#)
- Núñez-Otaño N.B., Imhof A., Bolcatto P.G., Larriera A. 2010.** Sex differences in the genitalia of hatchling *Caiman latirostris*. *Herpetological Review* 41:32–35.
- Parachú-Marcó M.V., Leiva P., Iungman J.L., Simoncini M.S., Piña C.I. 2017.** New evidence characterizing temperature-dependent sex determination in broad-snouted caiman, *Caiman latirostris*. *Herpetological Conservation and Biology* 12:78–84.
- Piña C.I., Larriera A., Siroski P., Verdade L.M. 2007a.** Cranial sexual discrimination in hatchling broad-snouted caiman (*Caiman latirostris*). *Iheringia* 97:17–20. [DOI](#)
- Piña C.I., Siroski P., Larriera A., Lance V.A., Verdade L.M. 2007b.** The temperature-sensitive period (TSP) during incubation of broad-snouted caiman (*Caiman latirostris*) eggs. *Amphibia Reptilia* 28:123–128. [DOI](#)
- Piña C.I., Merchant M.E., Verdade L.M. 2015.** Introduction: Reproduction in crocodylians. *South American Journal of Herpetology* 10:1–3. [DOI](#)
- Platt S.G., Rainwater T.R., Thorbjarnarson J.B., Finger A.G., Anderson T.A., McMurry S.T. 2009.** Size estimation, morphometrics, sex ratio, sexual size dimorphism, and biomass of Morelet's crocodile in northern Belize. *Caribbean Journal of Science* 45:80–93. [DOI](#)
- Platt S.G., Rainwater T.R., Thorbjarnarson J.B., Martin D. 2011.** Size estimation, morphometrics, sex ratio, sexual size dimorphism, and biomass of *Crocodylus acutus* in the coastal zone of Belize. *Salamandra* 47:179–192.
- Richardson K.C., Webb G., Manolis S.C. 2002.** Crocodiles: Inside Out. A Guide to the Crocodylians and Their Functional Morphology. Surrey Beatty & Sons, Chipping Norton.
- Schulte D., Chabreck R. 1990.** Effects of nest and egg characteristics on size and early development of American alligators. Pp. 177–187, in Proceedings of the 10th Working Meeting of Crocodile Specialist Group, IUCN-The World Conservation Union. Gland.
- Simoncini M.S., Cruz F.B., Larriera A., Piña C.I. 2014.** Effects of climatic conditions on sex ratios in nests of Broad-snouted Caiman. *Journal of Zoology* 293:243–251. [DOI](#)
- Valenzuela N., Lance V. 2004.** Temperature-dependent Sex Determination in Vertebrates. Smithsonian Books, Washington.
- Webb G.J.W., Manolis S.C., Sack G.C. 1984.** Cloacal sexing of hatchling crocodiles. *Australian Wildlife Research* 11:201–202. [DOI](#)
- Whitaker R., Whitaker Z., Vaughan A. 1980.** Notes on sexing crocodylians. *Journal of the Bombay Natural History Society* 77:341–343.
- Ziegler T., Olbort S. 2007.** Genital structures and sex identification in crocodiles. *Crocodile Specialist Group Newsletter* 26:16–17.