

## SUPPLEMENTARY ELECTRONIC MATERIAL

ARDEOLA 64(1)

MORPHOLOGICAL AND GENETIC DIFFERENTIATION AMONG KENTISH  
PLOVER *CHARADRIUS ALEXANDRINUS* POPULATIONS IN MACARONESIA

DIFERENCIACIÓN GENÉTICA Y MORFOLÓGICA ENTRE LAS POBLACIONES  
DE CHORLITEJO PATINEGRO *CHARADRIUS ALEXANDRINUS* DE LA  
MACARONESIA

Mohammed ALMALKI<sup>1, 2, 3, 10 \*</sup>, Krisztina KUPÁN<sup>4</sup>, María Cristina CARMONA-ISUNZA<sup>2</sup>,  
Pedrin LÓPEZ<sup>5</sup>, Ana VEIGA<sup>6</sup>, András KOSZTOLÁNYI<sup>7, 8</sup>, Tamás SZÉKELY<sup>2, 9, 10</sup> and  
Clemens KÜPPER<sup>3, 4, 10</sup>

<sup>1</sup> Department of Biology, College of Science, Taif University, P. O. 888, Taif, Kingdom of Saudi Arabia.

<sup>2</sup> Department of Biology and Biochemistry, University of Bath, Claverton Down, Bath BA2 7AY, UK.

<sup>3</sup> NERC-Biomolecular Analysis Facility, Department of Animal and Plant Sciences, University of Sheffield, Western Bank, Sheffield, S10 2TN, UK.

<sup>4</sup> Institute of Zoology, Universitätsplatz 2, A-8010 Graz, Austria.

<sup>5</sup> BIOS-CV, Boa Vista, Republic of Cape Verde.

<sup>6</sup> Directorate General of Ambient Environment, Praia, Republic of Cape Verde.

<sup>7</sup> MTA-DE 'Lendület' Behavioural Ecology Research Group, Department of Evolutionary Zoology, University of Debrecen, Egyetem tér 1., H-4032 Debrecen, Hungary.

<sup>8</sup> Department of Ecology, Faculty of Veterinary Science, Szent István University, Rottenbiller u.  
50, H-1077 Budapest, Hungary.

<sup>9</sup> Maio Biodiversity Foundation, Isla do Maio, Republic of Cape Verde.

<sup>10</sup> These authors contributed equally.

\* Corresponding author: m11m116@hotmail.com

#### TABLE S1

Result of pairwise Tukey HSD tests for wing and tarsus length of Kentish Plovers from different Macaronesian populations. Significant comparisons are presented in bold.

*[Resultados del test de Tukey HSD por pares para la longitud del ala y la longitud del tarso de chorlitejos patinegros de diferentes poblaciones macaronésicas. Las diferencias significativas se muestran en negrita]*

Comparison	Wing		Tarsus	
	Diff	<i>P</i>	Diff	<i>P</i>
FUV-STM	-0.15	1.00	-2.99	<b>&lt;0.001</b>
FUS-STM	-0.17	1.00	-0.54	0.05
CVM-STM	-4.15	<b>&lt;0.001</b>	0.50	<b>0.003</b>
OUG-STM	1.66	0.19	-0.92	0.01
SAM-STM	-1.43	<b>0.04</b>	-1.27	<b>&lt;0.001</b>
FUS-FUV	-0.02	1.00	2.45	<b>&lt;0.001</b>
CVM-FUV	-4.00	<b>&lt;0.001</b>	3.49	<b>&lt;0.001</b>
OUG-FUV	1.82	0.27	2.07	<b>&lt;0.001</b>
SAM-FUV	-1.27	0.39	1.72	<b>&lt;0.001</b>
CVM-FUS	-3.98	<b>&lt;0.001</b>	1.04	<b>&lt;0.001</b>
OUG-FUS	1.83	0.11	-0.38	0.76
SAM-FUS	-1.26	0.10	-0.73	<b>0.003</b>
OUG-CVM	5.81	<b>&lt;0.001</b>	-1.42	<b>&lt;0.001</b>
SAM-CVM	2.72	<b>&lt;0.001</b>	-1.77	<b>&lt;0.001</b>
SAM-OUG	-3.09	<b>&lt;0.001</b>	-0.36	0.82