

**Supplement Tab. 1:** List of the type specimens hosted in the arachnological collections of the MSNVR

Taxon	Data	Type	Collection	Jar/tube N°.	Notes
<b>Araneae</b>					
<b>Agelenidae</b>					
<i>Aterigena aliquoi</i> (Brignoli, 1971)	1 ♂, ITALY, Sicily, Piano della Battaglia, Madonie, 1600 m, 28. Jul. 1968, Aliquò leg.	Holotype	Generale	77	as <i>Tegenaria aliquoi</i>
<i>Coelotes arganoi</i> Brignoli, 1978	1 ♀, TURKEY, Vil. Erzincan, Sakaltutan Geçidi, 12. Jun. 1973, Argano, Boitani & Cottarelli leg.	Paratype	Brignoli	535	
<i>Coelotes coenobita</i> Brignoli, 1978	3 ♀♀ TURKEY, Vil. Trabzon, Sümela, 10. Jun. 1969, Brignoli leg.	Paratypes	Brignoli	535	
<i>Coelotes italicus</i> Kritscher, 1956	1 ♀, ITALY, Monti Sibillini, Valle del Lago Pilato, circa 1600–1900 m, 30. Jun. 1955, Ruffo e Biancheri leg.	Holotype	Generale	82	
	1 ♂, ITALY, Monte Gorzano, M.ti della Laga, 1800–2300 m, IX.1968, Osella leg.	Allotype			
	1 ♂, 3 ♀♀, same data as the allotype	Paratypes			
	3 ♀♀, ITALY, Monti Sibillini, Forca Viola 1939 m, 9. Jul. 1954, Ruffo leg.; 1 ♀, same data as the holotype	Syntypes		6	
<i>Coelotes luculli</i> Brignoli, 1978	1 ♀, TURKEY, Vil. Tokat, Ardiçli, near Niksar, 100 m, 7. Jun. 1969, Osella leg.	Holotype	Generale	100	
<i>Coelotes osellai</i> de Blauwe, 1973	1 ♂, ITALY, Maritime Alps, Besio valley, Rifugio Garelli mountain lodge, Jul. 1968, Osella leg.	Holotype	Generale	79	
	6 ♂♂, 33 ♀♀, ITALY, Cottian Alps, Demonte, Colle del Mulo mountain, 2500 m; Cottian Alps, Val Varaita valley, S. Anna di Bellino, 2000–2300, 1. Jul. 1977; Maritime Alps, Col Lombarda mountain; Val Pesio valley, Marguareis Lakes; Maritime Alps, Val Pesio, Passo delle Gorre; Maritime Alps, Val Gesso valley, Rifugio Soria mountain refuge; Maritime Alps, Val Pesio valley, Rifugio Garelli mountain refuge, Osella leg.	Paratypes			
<i>Coelotes titaniacus</i> Brignoli, 1977	1 ♀, same data as the holotype	Allotype			
<i>Coelotes vignai</i> Brignoli, 1978	1 ♀, GREECE, Larissa, around Karitsa, Mt. Ossa, 900–1200 m, 20. Nov. 1974, Vigna leg.	Holotype	Brignoli	535	
<i>Draconarius gurkha</i> (Brignoli, 1976)	2 ♀♀, TURKEY, Vil. Bolu, Abant, 1400 m, 17. Jul. 1971, Brignoli leg.	Paratypes	Brignoli	535	
<i>Draconarius lama</i> (Brignoli, 1976)	1 ♀, NEPAL, Pangpoche, Yaral, 3900 m, 25. May 1961, Janetschek leg.	Paratype	Brignoli	531	as <i>Coelotes gurkha</i>
	1 ♀, NEPAL, Mingbo-Tal, airstrip of Hillary–Makalu Expedition, 4800 m, 28. May 1961, Janetschek leg.	Paratype	Brignoli	531	as <i>Draconarius lama</i>
<i>Draconarius latellai</i> Marusik & Ballarin, 2011	1 ♂, PAKISTAN, Northern Areas, Gilgit–Baltistan, Gilgit District, Bagrot Valley, Burche Glacier, 27. Oct. 2008, Latella leg.	Holotype	Generale	129	
	2 ♀♀, PAKISTAN, Northern Areas, Gilgit District, Naltar Valley, 1. Oct. 2008, Ahmed leg.	Paratypes			
<i>Eratigena arganoi</i> (Brignoli, 1971)	1 ♀, ITALY, Lazio, Roma, Altopiano di Arcinazzo, 800 m, 17. Apr. 1966, Brignoli leg.	Holotype	Brignoli	62	as <i>Cicurina arganoi</i>
<i>Eratigena balearica</i> (Brignoli, 1978)	1 ♀, SPAIN, Majorque, pres de Valldemossa, 4. Apr. 1968, Hauser leg.	Paratype	Brignoli	68	as <i>Malthonica balearica</i>
<i>Eratigena florea</i> (Brignoli, 1974)	1 ♀, MEXICO, Chiapas, Comitan, Cueva de las Floresillas cave, 2265 m, 18. Mar. 1971, Zullini leg.	Holotype	Brignoli	491	as <i>Tegenaria florea</i>
<i>Eratigena sardoa</i> (Brignoli, 1977)	1 ♀, ITALY, Sardegna, Oristano, around Asuni, 20. Feb. 1971, Vigna leg.	Holotype	Brignoli	62	as <i>Malthonica sardoa</i>
	1 ♀, ITALY, Sardegna, Carbonia–Iglesias, Domusnovas, Grotta di S. Giovanni cave, 5. Mar. 1972, Puddu leg.; 1 ♀, Sassari, Limbara, Cantoniera Gaddau, 30. Mar. 1972, Brignoli leg.	Paratypes			
<i>Eratigena sicana</i> (Brignoli, 1976)	1 ♀, ITALY, Sicily, Palermo, Piana degli Albanesi, 20. Nov. 1972, Aliquò leg.	Holotype	Brignoli	62	as <i>Malthonica sicana</i>
	1 ♂, 18 ♀♀, ITALY, Sicily, Palermo, Carini, Grotta del Puntale cave, 30. Apr. 1974; 1 ♀, Mt. Maganoce, 2. May 1974; 1 ♀, Bosco Ficuzza, 4. May 1974, Vigna leg.	Paratypes			

Taxon	Data	Type	Collection	Jar/tube N°.	Notes
<i>Eratigena vomeroi</i> (Brignoli, 1977)	1 ♂, ITALY, Basilicata, Potenza, Lagonegro, Grotta del Cervaro cave, 9. Dec. 1966, Sbordoni leg.	Holotype	Brignoli	541	as <i>Tegenaria vomeroi</i>
	1 ♂, 1 ♀, same data as the holotype; 1 ♂, ITALY, Basilicata, Potenza, Tramutola, Risorgenza dell'Aquila, 900 m, 9. Apr. 1970, Sbordoni leg.	Paratypes			
<i>Himalcoelotes sherpa</i> (Brignoli, 1976)	1 ♀, NEPAL, Basislager Yaral (Pang Poche), Lichtfang (61–207), 3900 m, 25. May 1961, Janeztschek leg.	Paratype	Brignoli	531	as <i>Coelotes sherpa</i>
<i>Histopona hauseri</i> (Brignoli, 1972)	1 ♂, GREECE, Corfu, upper Bodolakos cave, 26. Mar. 1967, Hofer leg.; 2 ♀♀, same locality, 12. Apr. 1972, Hauser leg.	Paratypes	Brignoli	68	as <i>Roeweriana hauseri</i>
<i>Histopona isolata</i> Deeleman-Reinhold, 1983	1 ♀, GREECE, Crete, Topolia, Aghia Sofia, 19. Mar. 1981, Deeleman-Reinhold leg.	Paratype	Brignoli	68	
<i>Histopona italicica</i> Brignoli, 1977	1 ♀, ITALY, Lazio, Colli Albani, Mt. Cavo, 5. May 1968, Brignoli leg.	Holotype	Brignoli	68	
	1 ♀, ITALY, Tuscany, Firenze, Borgo San Lorenzo, Polcanto, 11. Mar. 1972, Magini leg.; 1 ♀, Molise, Campobasso, Matese, Boiano, 500–700 m, 18. Sep. 1967, Minelli leg.; 2 ♀♀, Roccamandolfi, 820 m, 29. Jun. 1967, Riggio & Osella leg.; 1 ♀, San Massimo, Piani di Campitello, 1500 m, 22. Sep. 1967, Minelli leg.; 1 ♀, Piemonte, Cuneo, Monviso, Crissolo, 1300 m, VII.1967, Osella leg.; 3 ♀♀, Maritime Alps, Val Pesio, Laghetti del Marguareis, 2100 m, VII.1968, osella leg.; 1 ♀, Lazio, Rieti, Monti della Laga, Amatrice, Capricchia, 1150–1300 m, IX.1968, Osella leg.	Paratypes			
<i>Histopona leonardoi</i> Bolzern, Pantini & Isaia, 2013	1 ♀, ITALY, Piemonte, Verbania, Cusio, Ossola, Grotta di S. Carlo Varzo cave, 4. Jun. 1979, Casale leg.	Paratype	Brignoli	68	
<i>Histopona palaeolithica</i> (Brignoli, 1971)	1 ♀, ITALY, Liguria, Savona, Finale Ligure, Loano, Arma delle Arene Candide cave, 34/Li, 25. Nov. 1967, Vigna leg.	Holotype	Brignoli	68	as <i>Chorizomma palaeolithicum</i>
<i>Histopona strinatii</i> (Brignoli, 1976)	1 ♂, 1 ♀, GREECE, Morea, Ton-Limnon cave, Kastria Pel 74/31., 17. May 1974, B. Hauser leg.	Paratypes	Brignoli	68	as <i>Roeweriana strinati</i>
<i>Histopona vignai</i> Brignoli, 1980	1 ♂, GREECE, Larissa, near Karixa, Mt. Ossa, 900–1200 m, 20. Nov. 1974, Vigna leg.; 1 ♀, GREECE, Kastarini, Mt. Olympus, Litochoran, 700 m, 1. Nov. 1974, Vigna leg.; 1 ♀, Larissa, near Karixa, Mt. Ossa, 900–1200 m, 20. Nov. 1974, Vigna leg.	Holotype	Brignoli	68	
		Paratypes			
<i>Inermocoelotes microlepidus</i> (de Blauwe, 1973)	1 ♀, ITALY, Veneto, Verona, Montecchio, 2. May 1968, Osella, Lazzeroni & Montolli leg.	Holotype	Generale	81	as <i>Coelotes microlepidus</i>
<i>Maimuna cariae</i> Brignoli, 1978	1 ♂, 6 ♀♀, TURKEY, Vil. Mugla, Gökböl, 30. Apr. 1973, Vigna leg.; 3 ♀♀, near Ferthiye, 29. Apr. 1973, Brignoli leg.	Paratypes	Brignoli	533	
<i>Malthonica paraschiae</i> Brignoli, 1984	1 ♀, GREECE, Naxos, 4. May 1982, Paraschi leg.	Holotype	Brignoli	62	
<i>Mistaria fagei</i> (Caporiacco, 1949)	2 ♀♀, KENYA, Mau, 150 km North of Nairobi, 2500–3000 m, Jan. 1946, Toschi, Meneghetti leg.	Paralectotype	Di Caporiacco	15	as <i>Agelena fagei</i>
<i>Mistaria nairobi</i> (Caporiacco, 1949)	1 ♂, 1 ♀, KENYA, Nairobi, 1500 m, 1944, Toschi, Meneghetti leg.	Paralectotype	Di Caporiacco	15	as <i>Agelena leucopyga nairobi</i>
<i>Pireneitega armeniaca</i> (Brignoli, 1978)	2 ♂♂, 5 ♀♀ TURKEY, Vil. Gümüşhane, Zigana geçidi, 2100 m, 12. Jul. 1971, Vigna leg.; 1 ♀ same data, Boitani leg.; 2 ♂♂, 7 ♀♀, same data, Osella leg.; 1 ♀ same locality, 2025 m, 12. Jun. 1969, Cottarelli leg.; 1 ♀ same locality 12. Jun. 1969, 1500 m, Osella leg.	Paratypes	Brignoli/ Generale	535 (Bri.), 99 (Gen.)	as <i>Coelotes armeniacus</i>
<i>Pireneitega cottarellii</i> (Brignoli, 1978)	2 ♀♀, TURKEY, Vil. Sinop, Drannaz dag, 1300 m, 29. Oct. 1974, Vigna leg.	Paratypes	Brignoli	535	as <i>Coelotes cottarellii</i>
<i>Pireneitega garibaldii</i> (Kritscher, 1969)	1 ♀, ITALY, Calabria, Reggio Calabria, S. Stefano in Aspromonte, Gambarie, 6. Jul. 1962, Kritscher leg.	Holotype	Generale	81	as <i>Coelotes garibaldii</i>
<i>Rualena pasquinii</i> Brignoli, 1974	1 ♀, ITALY, Calabria, Reggio Calabria, Aspromonte, Cippo Garibaldi, 6. Jul. 1962, Kritscher leg.	Paratype		52, 81	
	1 ♀, MEXICO, Chiapas, Comitán, S. Agostin, 2320 m, 6. Mar. 1971, Zullini leg.	Holotype	Brignoli	544	

Taxon	Data	Type	Collection	Jar/tube N°.	Notes
<i>Tegenaria agnoletti</i> Brignoli, 1978	1 ♀, TURKEY, Vil. Antalya, Dosemeatti, Yagca, Grotta Mustanini, 300 m, 30. Apr. 1973, Agnoletti leg.	Holotype	Brignoli	541	
<i>Tegenaria anhela</i> Brignoli, 1972	1 ♂, TURKEY, Vil. Antalya, Grotta Karain Magarasi, 14. Aug. 1967, Brignoli leg.	Holotype	Brignoli	541	
<i>Tegenaria ariadnae</i> Brignoli, 1984	1 ♀, GREECE, Crete, Perama, Melidoni Spilea cave, 14. Mar. 79, Hauser leg. (Kar 79/17)	Paratype	Brignoli	542	
<i>Tegenaria averni</i> Brignoli, 1978	1 ♀, TURKEY, Vil. Mersin, Silifke, Cennet Magarasi, 26. Apr. 1973, Brignoli leg.	Holotype	Brignoli	542	
<i>Tegenaria capolongoi</i> Brignoli, 1977	1 ♀, ITALY, Napoli, San Sebastiano al Vesuvio, Grotticella vesuviana cave, 10. Mar. 1973, Capolongo leg.	Holotype	Brignoli	543	
<i>Tegenaria commena</i> Brignoli, 1978	1 ♀, TURKEY, Vil. Trabzon, Sümela (Maçka), 1200 m, 13. Jul. 1971, Brignoli leg. (Bri.); 1 ♀, same data, Osella leg. (Gen.)	Paratypes	Brignoli/ Generale	544 (Bri.), 100 (Gen.)	
<i>Tegenaria cottarelli</i> Brignoli, 1978	1 ♀, TURKEY, Vil. Artvin, Borçka, 15–18. Jun. 1969, Cottarelli leg. (Bri.); 1 ♀, same data, Osella leg.? (Gen.)	Paratypes	Brignoli/ Generale	544 (Bri.), 100 (Gen.)	
<i>Tegenaria drescoi</i> Brignoli, 1971	1 ♀, ITALY, Sardegna, Sassari, Castelsardo, 28. Apr. 1967, Vigna leg.	Holotype	Brignoli	543	
<i>Tegenaria eleonorae</i> Brignoli, 1974	1 ♂, ITALY, Sardegna, Carbonia–Iglesias, Domusnovas, Grotta di S. Giovanni cave, 81 Sa/Ca, 6. Oct. 1968, Pilia, Usai leg.  2 ♀♀, ITALY, Sardegna, Carbonia–Iglesias, Domusnovas, Grotta di S. Giovanni cave, 81 Sa/Ca, 8. Sep. 1968, Pirodda, Latte, Pinna leg.; 1 ♀, same locality, 28. Feb. 1971, Vigna leg.	Holotype Paratypes	Brignoli	543 Paratypes	
<i>Tegenaria elysii</i> Brignoli, 1978	1 ♀, TURKEY, Vil. Mersin, Silifke, Dilek cave, Magrasi, 26. May 1973, Vigna leg.  8 ♀♀, TURKEY, Vil. Mersin, Silifke, Magrasi, Dilek cave, 26. Apr. 1973, Brignoli leg.; 1 ♀, Cemet cave, 26. Apr. 1973, Brignoli leg.; 1 ♀, same data, Di Rao leg.	Holotype Paratypes	Brignoli	542 Paratypes	
<i>Tegenaria faniapollinis</i> Brignoli, 1978	1 ♀, TURKEY, Vil. Antakya, near Harbiye, Harbiye cave, 26. Jun. 1971, Boitani leg.  3 ♀♀, same data as the holotype, Brignoli leg. (Bri.); 1 ♀, same data, Boitani leg. (Gen.)	Holotype Paratypes	Brignoli Brignoli/Ge- nerale	542 542 (Bri.), 100 (Gen.)	
<i>Tegenaria forestiero</i> Brignoli, 1978	1 ♀, TURKEY, Vil. Isparta, Kuruçaova, Asar Ini cave, 1250 m, 19. Apr. 1973, Forestiero leg.  3 ♀♀, same data as the holotype, Sbordoni leg.	Holotype Paratypes	Brignoli	541 Paratypes	
<i>Tegenaria hauseri</i> Brignoli, 1979	1 ♀, GREECE, Citera, Aghia Sophia cave, Mylopotamos, Cy77–10, 17. Apr. 1977, Hauser leg.	Paratype	Brignoli	542	
<i>Tegenaria karaman</i> Brignoli, 1978	1 ♀, TURKEY, Vil. Konya, Seydisehi, Ferzen Magarasi cave, 1410 m, 23. Apr. 1973, Forestiero leg.	Holotype	Brignoli	543	
<i>Tegenaria lyncea</i> Brignoli, 1978	1 ♂, TURKEY, Vil. Erzuru, Aşkale, 1800 m, 10. Jul. 1971, Osella leg.	Paratype	Generale	100	
<i>Tegenaria melbae</i> Brignoli, 1972	1 ♀, TURKEY, Vil. Diyarbakir, Lice, Korkha cave, 23. Jun. 1968, Brignoli leg.	Holotype	Brignoli	541	
<i>Tegenaria michae</i> Brignoli, 1978	1 ♀, LEBANON, Caza, Bcharre, Cedres de Brharri, 1250 m, 2–5. Jun. 1972, Brignoli leg.  1 ♀, same data as the holotype	Holotype Paratype	Brignoli	544 Paratype	
<i>Tegenaria parmenidis</i> Brignoli, 1971	1 ♀, ITALY, Campania, Salerno, Novi Velia, 2. Oct. 1967, Brignoli leg.	Holotype	Brignoli	543	
<i>Tegenaria pasquinii</i> Brignoli, 1978	1 ♀, TURKEY, Vil. Agri, Tahir Geçidi, 2500 m, 4. Jul. 1971, Brignoli leg.  9 ♀♀, TURKEY, Vil. Artvin, Sayşat, 2450 m, 6. Jul. 1971, Brignoli leg.; 1 ♀, same locality and date, Boitani L. leg. (Bri.); 1 ♀, same locality, Osella leg. (Gen.), 1 ♀, Vil. Kars, Çıldır, 5. Jul. 1971, Brignoli leg.; 1 ♀, same locality, 5. Jul. 1971, Boitani leg., 1 ♀, Ardahar, 2000 m, 6. Jul. 1971, Vigna leg. (Bri.)	Holotype Paratypes	Generale Brignoli/Ge- nerale	100 544 (Bri.), 100 (Gen.)	
<i>Tegenaria percuriosa</i> Brignoli, 1972	1 ♀, TURKEY, Vil. Isparta, Anamas, Zindan magarasi, 11. Aug. 1967, Sbordoni leg.	Holotype	Brignoli	543	
<i>Tegenaria pieperi</i> Brignoli, 1979	1 ♀, GREECE, Crete, Sitia, Megalo Katafigi, Ag. Georgios, 21. May 1977, Pieper leg.	Holotype	Brignoli	542	

Taxon	Data	Type	Collection	Jar/tube N°.	Notes
<i>Tegenaria regispyrri</i> Brignoli, 1976	1 ♂, GREECE, Epirus, Trikkala, Malakasi, 28. Sep. 1966, Brignoli leg. 2 ♀♀, GREECE, Epirus, Trikkala, Malakasi, 28. Sep. 1966, Brignoli leg. (Bri.); 1 ♀, Ioannina, Metsovon, 1400 m, 22. Jul. 1971, Osella leg. (Gen.)	Holotype Paratypes	Brignoli Brignoli/ Generale	544 544 (Bri.), 100 (Gen.)	
<i>Tegenaria sbordonii</i> Brignoli, 1971	1 ♂, ITALY, Lazio, Frosinone, Esperia, Grotta dei Serini cave, 480 La, 1000 m, 13. Dec. 1970, Sbordoni leg. 1 ♀, ITALY, Lazio, Frosinone, Esperia, Grotta dei Serini cave, 480 La, 1000 m, 13. Dec. 1970, Sbordoni leg.; 1 ♀, same locality, 17. May 1970, Sbordoni leg.; same locality 17. Jan. 1971, C.S.R. leg.	Holotype Paratypes	Brignoli	543	
<i>Tegenaria schmalfussi</i> Brignoli, 1976	1 ♀, GREECE, Milatos cave, 23. Aug. 1972, Schmalfus leg.; 1 ♀, same data.	Holotype	Brignoli	544	
<i>Tegenaria vallei</i> Brignoli, 1972	1 ♂, LYBIA, Cyrenaica, Benghazi, Lete cave, 1. Jan. 1968, Valle leg.	Paratype	Brignoli	543	
<i>Tegenaria wittmeri</i> Brignoli, 1978	1 ♂, BHUTAN, Thimphu, 31. May 1972, Nat. Hist. Museum Basel–Bhutan Expedition 1972 leg.	Paratype	Brignoli	542	
<i>Urocoras matesianus</i> (de Blauwe, 1973)	1 ♀, ITALY, Molise, Isernia, Matese, Roccamandolfi, 840 m, 27. Apr. 1967, Riggio & Osella leg. 1 ♀, same data as the holotype, 820 m, Osella leg.	Holotype Paratype	Generale	81	as <i>Coelotes matesianus</i>
<i>Urocoras nicomedis</i> (Brignoli, 1978)	1 ♀, TURKEY, Vil. Bolu, Abant, 1450 m, 17. Jul. 1971, Osella leg.	Paratype	Generale	99	as <i>Coelotes nicomedis</i>
<b>Amaurobiidae</b>					
<i>Amaurobius koponeni</i> Marusik, Ballarin & Omelko, 2012	1 ♂, INDIA, Uttar Pradesh, Farrukhabad District, Kaimganj City, 23. Mar. 2003, Abrescia leg.	Holotype	Generale	129	
<i>Amaurobius pesarinii</i> Ballarin & Pantini, 2017	1 ♂, ITALY, Lazio, Roma, Canale Monterano, Guallo, 179 m, 18. Sep.–16. Oct. 2007, Rizzo & Vigna leg. 4 ♂♂, same data as the holotype; 1 ♀, same locality, 18. Apr.–15. May 2007, all Rizzo & Vigna leg.	Holotype Paratypes	Generale	130	
<i>Amaurobius ruffoi</i> Thaler, 1990	1 ♂, 2 ♀♀, ITALY, Veneto, Verona, Bosco Chiesanuova, Lessini hills, Podestaria, 5. Sep. 1972, Osella leg.; 7 ♀♀, same locality, 1600 m, 17. Aug. 1972, Osella, Ruffo leg.; 1 ♀, Vallon di Malera, 23. Jun. 1966, Ruffo leg.; 1 ♀, same locality, 6. Jun. 1971, Osella leg.; 3 ♀♀, Tracchi, 29. Aug. 1969, Osella leg.; 3 ♀♀, Mt. Corno, in front of the Ciabattino cave, 29. May 1968, Lazzeroni, Gioco, Montolli leg.; 1 ♀, Passo Fittanze, 1450 m, 29. May 1968, Lazzeroni, Gioco, Montolli leg.; 2 ♀♀, Mt. Lessini, Alta Val Squaranto valley, 500 m, 27. Oct. 1968, Amici della montagna leg.; 1 ♀, Mt. Lessini, Malga Bagorno, 27. Oct. 1968, Amici della montagna leg.; 1 ♂, 1 ♀, Velo Veronese, 8. May 1968, Lazzeroni, Gioco leg.; 1 ♂, Ponte di Veja, 6. Dec. 1967, Lazzeroni, Osella leg.	Paratypes	Generale	131	
<b>Anapidae</b>					
<i>Metanapis montisemodi</i> (Brignoli, 1978)	1 ♂, 1 ♀, NEPAL, Thakkola, Nebrikot, 2700 m, 5–13. Nov. 1969, Martens leg.	Paratypes	Brignoli	74	as <i>Pseudanapis montisemodi</i>
<i>Pseudanapis schauenbergi</i> Brignoli, 1981	1 ♂, 1 ♀, MAURITIUS, Chamarel, 360 m, 22. Dec. 1974, P. Schauenberg leg. (Mau 75/49)	Paratypes	Brignoli	74	
<i>Teutoniella plaumanni</i> Brignoli, 1981	3 ♂♂, 2 ♀♀, BRAZIL, Nova Teutonia, 300–500 m, Oct. 1965, Plaumann leg.	Paratypes	Brignoli	74	
<i>Zangherella apuliae</i> (Caporiacco, 1949)	1 ♂, 1 ♀, ITALY, Apulia, Lecce, Castro, Castromarina, Zinzulusa cave, 3. Oct. 1948, Ruffo, Conci leg. 2 ♀♀, same locality as the lectotype, 24. Sep. 1949, Ruffo leg.	Lectotype Paralectotypes	Generale	70	as <i>Pseudanapis apuliae</i>
<b>Araneidae</b>					
<i>Araneus mauensis</i> Caporiacco, 1949	1 ♀, KENYA, Mau, 150 km North–West of Nairobi, 2500–3000 m, Jan. 1946, Toschi, Meneghetti leg.	Syntype	Di Caporiacco	73	

Taxon	Data	Type	Collection	Jar/tube N°.	Notes
<i>Araneus mauensis ocellatus</i> Caporiacco, 1949	2 ♀♀, KENYA, Mau, 150 km North-West of Nairobi, 2500–3000 m, Dec. 1945, Toschi, Meneghetti leg.	Syntypes	Di Capo-riacco	73	
<i>Cyclosa lawrencei</i> Caporiacco, 1949	1 ♀, KENYA, Kabete, 13 West of Nairobi, Aug. 1946, Toschi, Meneghetti leg.	Syntype	Di Capo-riacco	67	
<i>Hypsosinga lithyphantoides dealbata</i> Caporiacco, 1949	1 ♂, 1 ♀, KENYA, Nairobi, 1500 m, 1944, Toschi, Meneghetti leg.	Syntypes	Di Capo-riacco	71	
<i>Micrathena margerita</i> Levi, 1985	1 ♀, MEXICO, Chiapas, Lake Tziscao, 17–18. Aug. 1973, Brignoli leg.	Paratype	Brignoli	508	
<i>Micrathena tziscao</i> Levi, 1985	1 ♀, MEXICO, Chiapas, Laguna Tziscao, 17–18. Aug. 1973, Brignoli leg.	Paratype	Brignoli	508	
<b>Cheiracanthiidae</b>					
<i>Cheiracanthium punctipedellum</i> Caporiacco, 1949	1 ♀, KENYA, Mau, 150 km North-West of Nairobi, 2500–3000 m, 1944, Toschi, Meneghetti leg.	Syntype	Di Capo-riacco	86	
<i>Cheiracanthium schenkeli</i> Caporiacco, 1949	1 ♀, KENYA, Nairobi, 1500 m, 1944, Toschi, Meneghetti leg.	Syntype	Di Capo-riacco	86	
<i>Cheiracanthium tetragnathoides</i> Caporiacco, 1949	2 ♂♂, KENYA, Mau, 150 km North-West of Nairobi, 2500–3000 m, Jan. 1946, Toschi, Meneghetti leg.	Syntypes	Di Capo-riacco	86	
<b>Cybaeidae</b>					
<i>Cybaeus vignai</i> Brignoli, 1977	1 ♀, ITALY, Ligurian Alps, Certosa di Pesio, underground, 859 m, 17. Sep. 1964, Vigna leg.	Holotype	Brignoli	535	
<b>Corninnidae</b>					
<i>Brachyphaea simpliciauleata</i> Caporiacco, 1949	1 ♂, KENYA, Elmenteita, 100 km North of Nairobi, 1500 m, Dec. 1945, Toschi, Meneghetti leg.	Syntype	Di Capo-riacco	92	
<b>Dictynidae</b>					
<i>Dictyna tullgreni</i> Caporiacco, 1949	1 ♂, KENYA, Mau, 150 km North-West of Nairobi, 2500–3000 m, Jan. 1946, Toschi, Meneghetti leg.	Syntype	Di Capo-riacco	7	
<b>Dysderidae</b>					
<i>Dasumia mariandyна</i> Brignoli, 1979	1 ♀, TURKEY, Bolu Akgakoka, boreal woods, 300–400 m, 29.VI–10. May 1976, Korge leg.	Paratype	Brignoli	455	
<i>Dasumia sancticedri</i> Brignoli, 1978	1 ♂, LEBANON, Caza Bcharré, Cédres de Bcharré, 1950 m, 3–5. Jun. 1972, Brignoli leg.	Holotype	Brignoli	455	
<i>Dysdera apenninica</i> Alicata, 1964	1 ♂, ITALY, Molise, Campobasso, S. massimo, Campitello Matese, 1500 m, 16. Jun. 1962, Ruffo leg.	Holotype	Generale	4	
<i>Dysdera bottazziae</i> Caporiacco, 1951	1 ♂, ITALY, Apulia, Foresta Umbra, Gargano, Oct. 1950, Bottazzia leg.	Holotype	Generale	21	
<i>Dysdera brignoliana</i> Gasparo, 2000	1 ♂, ITALY, Lazio, Latina, Isola Ponza, Piana di Punta d'Incenso, 8. Dec. 1966, Brignoli leg.	Holotype	Brignoli	552	
<i>Dysdera fustigans</i> Alicata, 1966	2 ♀♀, same data as the holotype; 1 ♀, same locality 19. Jan. 1967, Argano leg.	Paratypes			
<i>Dysdera maronita</i> Gasparo, 2003	1 ♂, ITALY, Campania, Salerno, San Rufo, Monte Serra Nuda, 950 m, 9. May 1965, Ruffo leg.	Holotype	Generale	84	
<i>Dysdera monterossoi</i> Alicata, 1964	1 ♂ 1 ♀, ITALY, Calabria, Reggio Calabria, Gambarie, Aspromonte, 8. May 1957, Ruffo leg.	Paratypes			
<i>Dysdera osellai</i> Alicata, 1973	1 ♂, ITALY, Sicily, Trapani, Favignana, Marettimo, Isole Egadi, 21–23. Oct. 1967, Osella leg.	Holotype	Generale	97	
<i>Dysdera romana</i> Gasparo & Di Franco, 2008	1 ♀, ITALY, Lazio, Roma, Ciampino, Fiorano, 120 m, 2. Dec. 2004; 2 ♂♂, 16. Dec. 2004; 1 ♂, 15. Apr. 2005; 2 ♀♀, 29. Jun. 2005	Paratypes	Generale	84	

Taxon	Data	Type	Collection	Jar/tube N°.	Notes
<i>Dysdera silana</i> Alicata, 1965	1 ♂, ITALY, Calabria, Sila, Monte Botte Donato, 25. Jun. 1960, Ruffo leg.	Holotype	Generale	4	
<i>Dysdera sibyllinica</i> Kritscher, 1956	1 ♂, ITALY, Marche, Monti Sibillini, Alta Valle del Pilato, 1800–1900 m, 11. Jul. 1954, Ruffo leg.	Holotype	Generale	84	as <i>Dysdera ninnii sibyllinica</i>
<i>Dysdera vignai</i> Gasparo, 2003	1 ♂, LEBANON, Les Cèdres, Biblos, 2300 m, 26. Jun. 1971, Osella leg.	Holotype	Brignoli	143	
<i>Harpactea agnoletti</i> Brignoli, 1978	1 ♂, LEBANON, Les Cèdres, Biblos, 2300 m, 26. Jun. 1971, Brignoli leg.	Paratype			
<i>Harpactea alicatai</i> Brignoli, 1979	1 ♀, TURKEY, Vil. Isparta, Kuruçaova, Inogu Ini cave, 1280 m, 24. Apr. 1973, Agnoletti leg.	Holotype	Brignoli	517	
<i>Harpactea catholica</i> (Brignoli, 1984)	1 ♀, ITALY, Sardegna, Sassari, Bono, Passo S'Uccaidu, 1000 m, 31. Mar. 1972, Brignoli leg.	Holotype	Brignoli	517	
<i>Harpactea coccifera</i> Brignoli, 1984	1 ♀, GREECE, Crete, Hania, Akrotiri, Mon. Gonverneto, Grotta Katholica cave, 600 m, 10. Aug. 1974, Sbordoni leg.	Holotype	Brignoli	455	as <i>Minotauria catholica</i>
<i>Harpactea diraoi</i> Brignoli, 1978	1 ♂, 1 ♀, GREECE, gorge near Sfaka, along the road Sitia-Iraklion, 13. Mar. 1979, Hauser leg., ex coll. MHNG	Paratype	Brignoli	517	
<i>Harpactea fageli</i> Brignoli, 1980	1 ♂, 1 ♀, TURKEY, Vil. Marsin, near Mut, 25. Apr. 1973, Di Rao leg.; 1 ♀, same data, Vigna leg.	Paratypes	Brignoli	517	
<i>Harpactea isaurica</i> Brignoli, 1978	1 ♂, SPAIN, Salamanca, Puerto de Bejar, Jan. 1957, Fagel leg.	Paratypes	Brignoli	517	
<i>Harpactea nausicaae</i> Brignoli, 1976	2 ♀♀, TURKEY, Vil. Konya, çamlik Dalayman, near Koruk ini, 1200 m, 24. Apr. 1973, Di Rao leg.; 2 ♀♀, same data, Vigna leg.	Paratypes	Brignoli	517	
<i>Harpactea osellai</i> Brignoli, 1978	2 ♀♀, GREECE, Epirus, Greveniti, 1050 m, 27. Apr. 1973, Löbl leg.; 1 ♂, Ag. Komasos, 280 m, 25. Apr. 1973, Löbl leg.	Paratypes	Brignoli	517	
<i>Harpactea parthica</i> Brignoli, 1980	1 ♂, TURKEY, Vil. Amasya, Borabay Gölu, 1100 m, 4. Apr. 1969, Brignoli & Osella leg.	Holotype	Generale	100	
<i>Harpactea pisidica</i> Brignoli, 1978	1 ♀, IRAN, Teheran, Dowlat, Abad, loc. 20, 500 m, 18. Jul. 1975, Di Rad leg.	Holotype	Brignoli	517	
<i>Harpactea strinatii</i> Brignoli, 1979	1 ♀, TURKEY, Vil. Jsparta, Egridir, cave 6km from the crossroad to Anamas, 1300 m, 17. Apr. 1973, Sbordoni leg.	Holotype	Brignoli	517	
<i>Harpactea zannonensis</i> Alicata, 1966	1 ♂, GREECE, Morea, Pyrgos Dirov, Glyphada cave, 19. May 1976, Hauser leg.	Paratype	Brignoli	517	
<i>Harpactocrates troglophilus</i> Brignoli, 1978	1 ♂, ITALY, Lazio, Isola di Zannone, wood, 29. Jan. 1966, Argano & Vigna leg.	Holotype	Brignoli	516	
<i>Kaemis circe</i> (Brignoli, 1975)	1 ♀, 1 ♀, ITALY, Lazio, Isola di Zannone, wood, 29. Jan. 1966, Argano & Vigna leg.; 1 ♀, Caponegro, 28. Feb. 1966, Brignoli leg.	Paratypes			
<i>Parachtes romandiola</i> (Caporiacco, 1949)	1 ♂, TURKEY, Vil Isparta, Anamas, Gr. Zindan Magarasi, 1200 m, 17. Apr. 1973, Di Rao leg.	Holotype	Brignoli	469	
<i>Stalagtia argus</i> Brignoli, 1976	1 ♂, ITALY, Lazio, Circeo, bosco presso Villa Agnet, 30. Nov. 1968, Argano leg.	Holotype	Brignoli	79	as <i>Harpassa circe</i>
<i>Stalagtia kratochvili</i> Brignoli, 1976	1 ♀, ITALY, Lazio, Isola di Zannone, Caponegro, 23. Feb. 1966, Argano leg.	Paratype			
<i>Euagridae</i>	4 ♂♀, ITALY, Emilia–Romagna, Monte Gemelli, beech forest litter, 7. Apr. 1946; 1 ♂, 4 ♀♀, Campania, forest litter, 8. Aug. 1945; 1 ♂, 1 ♀, S. Marino, wood litter, 3. Oct. 1945; all Zangheri leg.	Syntypes	Zangheri	25325, 25334, 25335	as <i>Harpactocrates romandiola</i>
<i>Euagrus lynceus</i> Brignoli, 1974	1 ♀, GREECE, Epirus, Kopani 460 m, 3. May 1973, Löbl leg.; 1 ♂, GREECE, Epirus, Psaka lake, 5. Nov. 1974, Vigna leg.;	Paratypes	Brignoli	469, 517	
	1 ♀, GREECE, Epirus, Aristi, 620 m, 1. May 1973, Löbl leg.	Paratype	Brignoli	469	
	1 ♂, MEXICO, Chapas, Comitàn, S. Agostin, 2320 m, 6. Mar. 1971, Zullini leg.	Holotype	Brignoli	42	
	2 ♀♀, same data as the holotype.				

Taxon	Data	Type	Collection	Jar/tube N°.	Notes
<b>Filistatidae</b>					
<i>Kukulcania brignolii</i> (Alayón, 1981)	1 ♂, MEXICO, Oaxaca, Monte Alban, 7. Aug. 1973, Brignoli leg.	Holotype	Brignoli	19	as <i>Filistata brignolii</i>
<i>Zaitunia alexandri</i> Brignoli, 1982	1 ♀, IRAN, Fars, Kuhendjan, loc. 13, 27. May 1976, Zerunian leg.	Holotype	Brignoli	65	
<i>Zaitunia medica</i> Brignoli, 1982	1 ♀, IRAN, Esthan, Laybid, loc. 4, 2100 m, 7. Jul. 1975, Di Rao leg.	Holotype	Brignoli	65	
	1 ♀, IRAN, Esthan, Laybid, 2100 m, 7. Jul. 1975, Brignoli leg.	Paratype			
<i>Zaitunia persica</i> Brignoli, 1982	1 ♀, IRAN, Fars Dehbid, loc. 6, 2100 m, 24. May 1976, Brignoli leg.	Holotype	Brignoli	65	
	1 ♀, IRAN, Fars Dehbid, loc. 6, 24. May 1976, Brignoli leg.	Paratype			
<b>Gnaphosidae</b>					
<i>Drassodes braendegardi</i> Caporiacco, 1949	1 ♂, KENYA, Elmementeita, 100 km North of Nairobi, 1500 m, Dec. 1945, Toschi, Meneghetti leg.	Syntype	Di Capori- acco	80	
<i>Leptodrassus diomedae</i> Caporiacco, 1951	1 ♂, ITALY, Apulia, Tremiti Islands, S. Domino Is., 11. May 1948, leg.?	Holotype	Generale	24	
<i>Marinarozelotes adriaticus</i> (Caporiacco, 1951)	1 ♂, ITALY, Apulia, Tremiti Islands, S. Domino Is., 13. May 1948, leg.?	Holotype	Generale	23	as <i>Zelotes adriaticus</i>
<i>Scotophaeus affinis</i> Caporiacco, 1949	1 ♀, KENYA, Mau, 150 km North-West of Nairobi, 2500–3000 m, Jan. 1946, Toschi, Meneghetti leg.	Syntype	Di Capori- acco	82	
<i>Scotophaeus schenkeli</i> Caporiacco, 1949	1 ♂, KENYA, Nairobi, 1500 m, 1944, Toschi, Meneghetti leg.	Syntype	Di Capori- acco	82	
<i>Xerophaeus thomasi</i> (Caporiacco, 1949)	1 ♀, KENYA, Elmementeita, 100 km North of Nairobi, 1500 m, Dec. 1945, Toschi, Meneghetti leg	Syntype	Di Capori- acco	83	as <i>Zelotes thomasi</i>
<b>Hahniidae</b>					
<i>Alistra mendanai</i> Brignoli, 1986	1 ♀, SOLOMON ISLANDS, Guadalcanal Is., Mt. Austin, 7. Feb. 1982, Brignoli leg.	Holotype	Brignoli	566	
<i>Cicurina paplagoniae</i> Brignoli, 1978	1 ♀, TURKEY, Vil. Kostamonu, Ballidag, 1350 m, 31. May 1969, Osella leg.	Paratype	Generale	101	
<i>Hahnia harmae</i> Brignoli, 1977	1 ♂, TUNISIA, Jendouba, Aïn Draham, 1. Oct. 1972, Brignoli leg.	Holotype	Brignoli	58	
<i>Hahnia hauseri</i> Brignoli, 1978	1 ♀, SPAIN, Majorca, along the route Palma–Manacor, near Algaida, 1. Apr. 1968, Hauser leg.	Paratype	Brignoli	58	
<i>Hahnia maginii</i> Brignoli, 1977	1 ♀, ITALY, Tuscany, Vetta delle Croci, 7. Jan. 1973, Magini leg.	Holotype	Brignoli	58	
<i>Hahnia musica</i> Brignoli, 1978	1 ♂, 1 ♀, BHUTAN, 87 km from Phuntsholing–Thimphu, 2215 m, 22. May 1972, Nat. Hist. Museum Basel–Bhutan expedition leg.	Paratypes	Brignoli	58	
<i>Hahnia rossii</i> Brignoli, 1977	1 ♀, ITALY, Marche, Ancona, Frasassi, Grotta del Fiume cave, 10. Jun. 1972, Rossi leg.	Holotype	Brignoli	58	
<i>Hahnia tikaderi</i> Brignoli, 1978	1 ♀, BHUTAN, Dechhi Paka, loc. 48, 3300 m, 20. Jun. 1972, Nat. Hist. Museum Basel–Bhutan expedition leg.	Paratype	Brignoli	58	
<i>Neoantistea janetscheki</i> Brignoli, 1976	2 ♂♂, 1 ♀, NEPAL, Valley of the Nare Drangka, trail to the Green Hut Hillary–Malcalu expedition, 5250 m, 22. May–3. Jun. 1961, Janetschek leg.	Paratypes	Brignoli	58	
<i>Neoantistea maxima</i> (Caporiacco, 1935)	1 ♀, PAKISTAN, Karakorum, meadows of Lopsang Bransa, near Mustagh glacier, 4500 m, 8. Jun. 1929, di Caporiacco leg.	Paralectotype	Brignoli	58	as <i>Hahnia maxima</i>
<b>Idiopidae</b>					
<i>Idiops melloleitaoi</i> (Caporiacco, 1949)	1 ♀, KENYA, Mackinnon Road, 80 km West of Mombasa, 500 m, Dec. 1946, Toschi, Meneghetti leg.	Syntype	Di Capori- acco	2	as <i>Titanidiops melloleitaoi</i>

Taxon	Data	Type	Collection	Jar/tube N°.	Notes
<b>Leptonetidae</b>					
<i>Barusia laconica</i> (Brignoli, 1974)	1 ♀, GREECE, Laconia, Pyrgos Dyrou, 18. Aug. 1970, Senglet leg.	Paratype	Brignoli	483	as <i>Paraleptoneta laconica</i>
<i>Cataleptoneta aesculapii</i> (Brignoli, 1968)	1 ♀, TURKEY, Alanya, Vil. Antalya, Grotta Damlatas, 15. Aug. 1967, Brignoli & Sbordoni leg. 7 ♀♀, TURKEY, Alanya, Vil. Antalya, Grotta Damlatas, 15. Jul. 1967, Brignoli & Sbordoni leg.	Holotype Paratypes	Brignoli	468	as <i>Paraleptoneta aesculapii</i>
<i>Cataleptoneta sordonii</i> (Brignoli, 1968)	1 ♂, TURKEY, Vil. Burdur, dint. Burdur, Jnsuyu cave, 1250 m, 13. Aug. 1967, Sbordoni leg. 1 ♀, same data as the holotype	Holotype Paratypes	Brignoli	483	as <i>Paraleptoneta sordonii</i>
<i>Cataleptoneta sengleti</i> (Brignoli, 1974)	1 ♂, 1 ♀, GREECE, Crete, Rethimnon caves, Mélidionion, 30. Jun. 1970, Senglet leg.	Paratypes	Brignoli	483	as <i>Paraleptoneta sengleti</i>
<i>Darkoneta arganoi</i> (Brignoli, 1974)	1 ♀, MEXICO, Chiapas, Cueva Grande de S. Agostin, Comitan, 2430 m, 4. Mar. 1971, Argano leg.	Holotype	Brignoli	545	as <i>Neoleptoneta arganoi</i>
<i>Leptonetela strinatii</i> (Brignoli, 1976)	1 ♀, GREECE, Attica, Keratea, Spilia tou Panos cave, 13. May 1974, Hauser leg.	Paratype	Brignoli	39	as <i>Proteleptoneta strinatii</i>
<i>Neoleptoneta caliginosa</i> Brignoli, 1977	1 ♀, MEXICO, Oaxaca, 1800 m, 6–8. Aug. 1973, Brignoli leg.	Holotype	Brignoli	39	
<i>Proteleptoneta baccettii</i> (Brignoli, 1979)	1 ♂, ITALY, Tuscany, Riomarina, Isola d'Elba, Grotta Ugliero cave, 5. May 1976, Sammuri leg. 3 ♀♀, same data as the holotype, Sbordoni leg.	Holotype Paratypes	Brignoli	466	as <i>Leptoneta baccettii</i>
<b>Linyphiidae</b>					
<i>Acartauchenius himalayensis</i> Tanasevitch, 2011	1 ♂, PAKISTAN, Northern Areas, Gilgit District, Bagrot Valley, 2500 m, 20. Jun. 2008, Latella leg.	Holotype	Generale	129	
<i>Araeoncus gertschi</i> Caporiacco, 1949	1 ♀, KENYA, Mau, 150 km North of Nairobi, 2500–3000 m, Jan. 1946, Toschi, Meneghetti leg.	Syntype	Di Caporiacco	47	
<i>Acartauchenius minor</i> (Millidge, 1979)	1 ♂, ITALY, Isola Ponza, Piana di Punta d'Incenso, Dec. 1966, Brignoli leg.	Holotype	Brignoli	66	as <i>Trachelocamptus minor</i>
<i>Araeoncus sicanus</i> Brignoli, 1979	1 ♀, ITALY, Sicily, Catania, Grotta Marrano cave, 1. Mar. 1975, leg.	Paratype	Brignoli	66	
<i>Asthenargellus kastoni</i> Caporiacco, 1949	1 ♀, KENYA, Mau, 150 km North of Nairobi, 2500–3000 m, Jan. 1946, Toschi, Meneghetti leg.	Syntype	Di Caporiacco	44	
<i>Asthenargellus meneghettii</i> Caporiacco, 1949	1 ♀, KENYA, Mau, 150 km North of Nairobi, 2500–3000 m, Jan. 1946, Toschi, Meneghetti leg.	Syntype	Di Caporiacco	44	
<i>Centromerus bonaeviae</i> Brignoli, 1979	1 ♀, ITALY, Sardegna, Sud Sardegna, Iglesias, Grotta di Buoncammino cave, 25. Feb. 1968, Puddu leg. 2 ♀♀, same data as the holotype	Holotype Paratypes	Brignoli	533 63	
<i>Centromerus cottarelli</i> Brignoli, 1979	1 ♂, ITALY, Liguria, La Spezia, Grotta Grande di Pignone cave, 36/Li, 22. Nov. 1967, Cottarelli & Vigna leg.	Holotype	Brignoli	533	
<i>Centromerus pasquinii</i> Brignoli, 1971	2 ♀♀, same data as the holotype 1 ♂, ITALY, Piemonte, Garessio, Alpi Liguri, Arma dei Grai Inferiore, 120 Pi, 25. Apr. 1969, Casale leg. 1 ♂, ITALY, same locality as the holotype, 31. Aug. 1967; 2 ♀♀, 25. Aug. 1968, all Vigna leg.	Paratypes Holotype Paratypes		63, 533 533	

Taxon	Data	Type	Collection	Jar/tube N°.	Notes
<i>Centromerus puddui</i> Brignoli, 1979	1 ♂, ITALY, Sardegna, Sud Sardegna, Villamassargia, Monte Ollastu, Grotta dei Pipistrelli cave, 1. Feb. 1976, Puddu leg. 1 ♂, 1 ♀, same data as the holotype; 1 ♂, ITALY, Basilicata, Potenza, Tramutola, Risorgenza dell'Aquila, 900 m, 9. Apr. 1970, Sbordoni leg.	Holotype Paratypes	Brignoli	63	
<i>Centromerus tongiorgii</i> Ballarin & Pantini, 2020	1 ♂, ITALY, Veneto, Verona, Negrar, Val Borago, South-West of Montecchio village, 463 m, 8. Jan.–12. Feb. 2004, submesophilic meadow and common hornbeam (Carpinus betulus) deciduous forest, in the litter, pitfall traps, Reich leg. 2 ♂♂, same data as the holotype, 12. Feb.–18. Mar. 2004, Reich L. leg., 2 ♂♂, 1 ♀, Verona, Grezzana, Val Galina, Le Volpare, 520 m, 28. Nov. 2003–8. Jan. 2004, submesophilic meadow, in the litter, pitfall traps, Reich leg.	Holotype Paratypes	Generale	116	
<i>Diplocephalus arnoi</i> Isaia, 2005	1 ♂, ITALY, Abruzzo, Pescara, Popoli, Sorgenti di S. Callisto, 300 m, 28. Aug. 2004, Leg. Osella 5 ♂, 15 ♀, same data as the holotype	Holotype Paratypes	Generale	116	
<i>Diplocephalus turcicus</i> Brignoli, 1972	1 ♂, TURKEY, vil. Burdur, Linsuyu cave, 13. Aug. 1967, Brignoli leg. 3 ♀♀, 1 juv., TURKEY, vil. Burdur, Linsuyu cave, 13. Aug. 1967, Brignoli leg.	Holotype Paratypes	Brignoli	66	
<i>Erigonoplus simplex</i> Millidge, 1979	1 ♂, ITALY, Apulia, Massafra, Bosco delle Pianelle, May 1967, Brignoli leg.	Holotype	Brignoli	66	
<i>Gonatium petrunkevitschi</i> Caporiacco, 1949	1 ♂, KENYA, Nairobi, 1500 m, 1944, Toschi, Meneghetti leg.	Syntype	Di Caporiacco	42	
<i>Improphanes mauensis</i> (Caporiacco, 1949)	1 ♀, KENYA, Mau, 150 km North of Nairobi, 2500–3000 m, Jan. 1946, Toschi, Meneghetti leg.	Syntype	Di Caporiacco	37	as <i>Leptyphantes mauensis</i>
<i>Mansuphantes auruncus</i> (Brignoli, 1979)	1 ♂, ITALY, Lazio, Latina, Grotta di Valmarino, 599 La, Mar. 1973, Sbordoni leg. 1 ♀, same data as the holotype	Holotype Paratype	Brignoli	533 61	as <i>Leptyphantes auruncus</i>
<i>Mecopisthes latinus</i> Millidge, 1978	2 ♂♂, 3 ♀♀, 1 juv., ITALY, Lazio, Roma, Riano Flaminio, oakwood litter, 5. Mar. 1967, Sbordoni leg.; 1 ♂, Tuscany, Firenze, Birigliano, museo terra, 25. Mar. 1967, Bordoni leg.	Paratypes	Brignoli	66	
<i>Microctenonyx apuliae</i> (Caporiacco, 1951)	1 ♀, ITALY, Apulia, Brindisi, Ostuni, Sep. 1950	Paratype	Generale	21	as <i>Aulacocyba apuliae</i>
<i>Mugiphantes brunneri</i> (Thaler, 1984)	3 ♀♀, ITALY, Veneto, Dolomiti, Mt. Cristallo, Passo del Cristallo, Cima Principale, 2800–3200 m, 10. Aug. 1983, Thaler leg.	Paratypes	Generale	102	as <i>Leptyphantes brunneri</i>
<i>Mugiphantes merretti</i> (Millidge, 1975)	2 ♂♂, 2 ♀♀, ITALY, Trentino A.A., Sella Group, passo del Pordoi, Dolomiti Mt., 2900 m, 16. Jul. 1973, Thaler leg.	Paratypes	Generale	102	as <i>Leptyphantes merretti</i>
<i>Palliduphantes brignolii</i> (Kratochvíl, 1978)	3 ♀♀, CROATIA, Dalmazia, Cavtat Sipun Pecina, 21. Oct. 1968, Vigna leg.	Paralectotypes	Brignoli	61	as <i>Leptyphantes brignolii</i>
<i>Palliduphantes carusoii</i> (Brignoli, 1979)	1 ♀, ITALY, Sicily, Siracusa, Grotta Palombara cave, 9. Dec. 1973; 1 ♂, Catania, Grotta Nuovalucello cave, 10. Oct. 1974, Caruso & Brisolese leg.	Paratypes	Brignoli	61	
<i>Palliduphantes conradini</i> (Brignoli, 1971)	1 ♂, ITALY, Abruzzo, Aquila, Verrecchia, Grotta Beatrice Cenci cave 3A, 12. Nov. 1967, Sbordoni leg.	Holotype	Brignoli	61	as <i>Leptyphantes conradini</i>
<i>Palliduphantes epaminondae</i> (Brignoli, 1979)	1 ♂, 2 ♀♀, GREECE, Beozia, Arachora, Koryghion Antron cave, cy 77–17, 1340 m, 19. Apr. 1977, Tranoodakis leg.	Paratypes	Brignoli	70	as <i>Leptyphantes epaminondae</i>
<i>Palliduphantes garganicus</i> (Caporiacco, 1951)	1 ♀, ITALY, Apulia, Gargano, cave Montenero, 22. Oct. 1949, Ruffo leg. 1 ♀, same data as the lectotype	Lectotype Paralectotype	Generale	70	as <i>Leptyphantes garganicus</i>
<i>Palliduphantes sbordonii</i> (Brignoli, 1970)	1 ♀, IRAN, Ask, Demarend, Cave A6, 1200 m, 26. Aug. 1966, Sbordoni leg.	Holotype	Brignoli	61	as <i>Leptyphantes sbordonii</i>

Taxon	Data	Type	Collection	Jar/tube N°.	Notes
<i>Tapinocyba latia</i> Millidge, 1979	1 ♂, ITALY, Lazio, Monti Reatini, Monte Catabio, Oct. 1967, Cottarelli leg. 1 ♂, ITALY, Lazio, Monti Reatini, Monte Catabio, 12. Oct. 1967	Holotype Paratype	Brignoli	66	
<i>Tapinocyba lucana</i> Millidge, 1979	1 ♀, ITALY, Basilicata, Potenza, Viggiano, Mar. 1967, Vomero leg. 1 ♂, ITALY, Apulia, Foggia, S. Angelo Gargano, 25. Oct. 1966	Holotype Paratype	Brignoli	66	
<i>Tapinocyba maureri</i> Thaler, 1991	3 ♂♂, 1 ♀, ITALY, Trentino A. A., Bolzano, Lagundo Schweigl leg.	Paratypes	Generale	131	
<i>Tchatkalophantes baltistan</i> Tanasevitch, 2011	1 ♀, PAKISTAN, Karakoru m, Baltistan, Shalabot, 1700 m, 15. Aug. 1976, Osella leg.	Holotype	Generale	129	
<i>Toschia picta</i> Caporiacco, 1949	2 ♂♂, 2 ♀♀, KENYA, Mau, 150 km North of Nairobi, 2500–3000 m, Jan. 1946, Toschi, Meneghetti leg.; 1 ♂, same data	Syntypes	Di Caporiacco	43	
<i>Troglohyphantes bolognai</i> Brignoli, 1975	1 ♀, ITALY, Liguria, Imperia, Tana Bertrand cave, 1. Sep. 1974, Bologna & Bonzano leg.	Holotype	Brignoli	79	
<i>Troglohyphantes caporiaccoi</i> Brignoli, 1971	1 ♂, ITALY, Lombardia, Bergamo, Rota Imagna, Valle Imagna, Tomba del Polacco cave, 1003 lo, 5. Mar. 1966, Sbordoni & Vigna leg. 1 ♀, same data as the holotype	Holotype Paratype	Brignoli	72	
<i>Troglohyphantes cruentus</i> Brignoli, 1971	1 ♂, SLOVENIA, Mostna Soci Lom. Esterno Smojanica, 15. Oct. 1968, Vigna leg.	Holotype	Brignoli	79	
<i>Troglohyphantes exul</i> Thaler, 1987	1 ♂, ITALY, Veneto, Verona, S. Ambrogio di Valpolicella, Cava di S. Ambrogio di Valpolicella, 16. Sep. 1984, Caoduro leg.	Holotype	Generale	102	
<i>Troglohyphantes iulianae</i> Brignoli, 1971	1 ♂, 2 ♀♀, same locality as the holotype, 27. Feb. 1983; 4 ♀♀, 16. Sep. 1984, all Caoduro leg. 1 ♀, ITALY, Liguria, La Spezia, Grotta di Cassana cave, 65Li, 1. Dec. 1968, Ceccarelli & Vigna leg.	Paratypes Holotype	Brignoli	72	
<i>Troglohyphantes juris</i> Thaler, 1982	1 ♀, same data as the holotype; 1 ♀, same locality, 23. Nov. 1969, all Vigna leg.	Paratypes			
<i>Troglohyphantes konradi</i> Brignoli, 1975	2 ♀, ITALY, Friuli Venezia-Giulia, Pordenone, Vito d'Asio, Inghiottoio di Juris, Fr. 623, 540 m, 22. Jul. 1979, Gasparo leg. 1 ♂, ITALY, Piemonte, Cuneo, Val Vermenagna, sotterranei a valle di Vernante, 14. Oct. 1972, Morisi leg. 10 ♀♀, same locality as the holotype, 14. Oct. 1972 and 10. Feb. 1973; 1 ♂, 1 ♀, 17. Nov. 1974, all Morisi leg.	Paratypes Holotype Paratypes	Brignoli	79	
<i>Troglohyphantes lessinensis</i> Caporiacco, 1936	1 ♂, ITALY, Veneto, Verona, Covoli di Velo, Monti Lessini, 878 m, 21. Nov. 1934, Ruffo leg.	Lectotype	Generale	5	
<i>Troglohyphantes nigraerosae</i> Brignoli, 1971	1 ♂, ITALY, Alpi Graie, Colle dell'Arietta, 2900 m, 27. Jul. 1966, Vigna leg.	Holotype	Brignoli	72	
<i>Troglohyphantes ruffoi</i> Caporiacco, 1936	1 ♂, ITALY, Veneto, Verona, S. Anna d'Alfaedo, Ponte di Veja, Grotta Grande, Monti Lessini, 600 m, 16. Jul. 1934, Ruffo leg.	Neotype	Generale	69	
<i>Troglohyphantes pisidicus</i> Brignoli, 1971	1 ♂, TURKEY, Vil. Konya, Beysihir lake Island Hadji Akif cave, 10. Aug. 1967, Brignoli leg.	Holotype	Brignoli	79	
<i>Troglohyphantes sbordonii</i> Brignoli, 1975	1 ♂, ITALY, Friuli Venezia Giulia, Pradis di sotto, Grotta Mainarie dal Punt I cave, 500 m, 13. Feb. 1975, Sbordoni V. leg.	Holotype	Brignoli	79	
<i>Troglohyphantes vignai</i> Brignoli, 1971	1 ♂, same data as the holotype 1 ♂, ITALY, Piemonte, Crissolo, Alpi Cozie, Buco di Valenza cave, 1009 pi, 24. Aug. 1967, Vigna leg. 4 ♀♀ (Bri.), 2 ♀♀ (Gen.), same data as the holotype	Paratype Holotype Paratypes	Brignoli Brignoli Brignoli/ Generale	72 72 72 (Bri.), 70 (Gen.)	

Taxon	Data	Type	Collection	Jar/tube N°.	Notes
<i>Walckenaeria angelica</i> Millidge, 1979	1 ♀, ITALY, Apulia, Foggia, S. Angelo Gargano, Oct. 1966, Sbordoni leg. 1 ♀, same data as the holotype	Holotype Paratype	Brignoli	66	
<i>Walckenaeria insperata</i> Millidge, 1979	1 ♂, ITALY, Isola Ponza, Piana di Punta d'Incenso, Dec. 1966, Vigna leg. 1 ♀, ITALY, Isola Ponza, Piana di Punta d'Incenso, 12. Nov. 1966, Cottarelli leg.; 1 ♂, M. Rosa beach, 23. Feb. 1968, Vigna leg.	Holotype Paratypes	Brignoli	66	
<i>Walckenaeria plumata</i> Millidge, 1979	1 ♂, ITALY, Isola Ponza, Piana di Punta d'Incenso, Dec. 1966, Vigna leg. 1 ♀, ITALY, Calabria, Crotone, Capocolonna, 24. Aug. 1966, Lunare leg.; 1 ♀, Tuscany, Firenze, along the road, Apr. 1967, ? leg.; 2 ♂♂, Friuli Venezia Giulia, Gorizia, Doberdò, 2. Nov. 1968, Brignoli leg.	Holotype Paratypes	Brignoli	66	
<b>Liocranidae</b>					
<i>Cybaeodes alicatai</i> Platnick & Di Franco, 1992	1 ♀, TUNISIA, Ain Draha m, Oued Zeed, 13. Apr. 1969, Alicata leg.	Holotype	Generale	103	
<i>Cybaeodes avolensis</i> Platnick & Di Franco, 1992	1 ♂, ITALY, Sicily, Siracusa, Cava grande di Avola, 6. Sep.–12. Oct. 1989, Adorno leg. 1 ♀, same data as the holotype	Holotype Allotype	Generale	103	
<i>Cybaeodes carusoi</i> Platnick & Di Franco, 1992	1 ♂, ALGERIA, Bejaïa, Pic des Singes, 400 m, 3. Nov. 1984, Marcellino leg.	Holotype	Generale	103	
<i>Cybaeodes sardus</i> Platnick & Di Franco, 1992	1 ♀, ITALY, Sardinia, Oristano, Parco Laconi, 550 m, 1. Oct. 1963, Alicata leg.	Holotype	Generale	103	
<i>Mesiobelus pococki</i> Caporiacco, 1949	1 ♀, KENYA, Mau, 150 km North–West of Nairobi, 2500–3000 m, Jan. 1946, Toschi, Meneghetti leg.	Syntype	Di Caporiacco	87	
<b>Lycosidae</b>					
<i>Alopecosa fabrilis trinacriae</i> Lugetti & Tongiorgi, 1969	1 ♂, ITALY, Sicily, Catania, Mt. Etna, Rifugio Citelli, 10. Apr. 1963, Alicata leg.	Holotype	Generale	43	
<i>Alopecosa osellai</i> Lugetti & Tongiorgi, 1969	1 ♂, SPAIN, Saragoza, Sierra de Montcayo, 2000 m, 13. May 1967, Osella leg. 3 ♂♂, 7 ♀♀, same data as the holotype	Holotype Paratypes	Generale	44	
<i>Arctosa berlandi</i> (Caporiacco, 1949)	1 ♀, KENYA, Elmenteita, 100 km North of Nairobi, 1500 m, Dec. 1945, Toschi, Meneghetti leg.	Holotype	Di Caporiacco	23	as <i>Hogna berlandi</i>
<i>Pardosa gromovi</i> Ballarin, Marusik, Omelko & Koponen, 2012	2 ♂♂, 1 ♀, KAZAKHSTAN, Almaty area, 777 m, 24. Jun. 2011, Gromov leg.	Paratypes	Generale	129	
<i>Pardosa persica</i> Marusik, Ballarin & Omelko, 2012	5 ♂♂, 5 ♀♀, IRAN, Fars Province, Haft Bar m, 24. May 2000, Marusik leg.	Paratypes	Generale	129	
<i>Pardosa zonsteini</i> Ballarin, Marusik, Omelko & Koponen, 2012	1 ♂, KYRGYZSTAN, Baubanh–Atu, Arslanbob, 1000–1800 m, scrub litter and under stones, 19. May 1993, Glovatch leg.; 5 ♂♂, Inner Tien–Shang, Kokirim Mountain Range, Kainy, 3. Jul. 199?, Ovtchinnikov leg.	Paratypes	Generale	129	
<b>Mimetidae</b>					
<i>Mimetus bishopi</i> Caporiacco, 1949	1 ♀, KENYA, Nairobi, 1500 m, 1944, Toschi, Meneghetti leg.; 1 ♀, 150 km North–West of Nairobi, 2500–3000 m, Jan. 1946, Toschi, Meneghetti leg.	Syntypes	Di Caporiacco	77	
<i>Mimetus vespillo</i> Brignoli, 1980	1 ♀, INDONESIA, Sulawesi, Sultan, near Rantspao, station. 11, inside a small cave, 24. Jan. 1979, Brignoli leg.	Holotype	Brignoli	67	

Taxon	Data	Type	Collection	Jar/tube N°.	Notes
<b>Mysmenidae</b>					
<i>Maymena sbordonii</i> Brignoli, 1974	1 ♂, MEXICO, Chiapas, Ocozocantla, Rancho del Cielito, Cueva de Cerro Brugio, 1320 m, 22. Mar. 1971, Sbordoni leg. 4 ♀♀, same data as the holotype	Holotype Paratypes	Brignoli	545	
<b>Nemesiidae</b>					
<i>Raveniola niedermeieri</i> (Brignoli, 1972)	1 ♀, IRAN, Astrabad, Nierdermayer leg.	Paratype	Brignoli	42	as <i>Brachythele niedermeieri</i>
<b>Nesticidae</b>					
<i>Domitius culsu</i> Ballarin, 2020	1 ♂, ITALY, Tuscany, Garfagnana area, Lucca province, Coreglia Antelminelli municipality, Tana delle Fate di Coreglia Antelminelli cave, 141/T/LU, 260m a.s.l., 44.046336°N 10.523525°E, 21. Aug. 2018, Ballarin & Avesani leg. 1 ♀, same locality as the holotype, 04. Sep. 1967, Vigna Taglianti leg.; 1 ♀, 15. Aug. 2015, Ballarin & Gaiga leg.; 4 ♀♀, 24. Jun. 2017, Ballarin & Ballarin; 2 ♀♀, 21. Aug. 2018, leg. Ballarin & Avesani leg.	Holotype Paratypes	Generale	90	
<i>Domitius sbordonii</i> (Brignoli, 1979)	1 ♂, ITALY, Lazio, Frosinone, Supino, Valle Serena, Grotta della Croce cave, 1. Nov. 1977, Sbordoni leg. 1 ♀, ITALY, Lazio, Frosinone, Supino, Grotta della Croce cave, 8. Feb. 1976, Sbordoni leg.	Holotype Paratype	Brignoli	524	as <i>Nesticus sbordonii</i>
<i>Gaucelmus strinati</i> Brignoli, 1979	1 ♀, GUATEMALA, Alta Vera Paz, Cueva Chirrupeck, 6–8. Apr. 1973, Strinati leg.	Paratype	Brignoli	524	
<i>Nesticella aelleni</i> (Brignoli, 1972)	1 ♀, SRI LANKA, Kuruwita, Stripura caves, 22. Jan. 1970, Aellen & Strinati leg.	Paratype	Brignoli	524	as <i>Nesticus aelleni</i>
<i>Nesticus arganoi</i> Brignoli, 1972	1 ♀, MEXICO, Veracruz, Orizalba, Tlilapan, Cueva de Ojo de Agua, 1150 m, 7. Jan. 1969, Argano leg.	Holotype	Brignoli	545	
<i>Nesticus brasiliensis</i> Brignoli, 1979	1 ♀, BRAZIL, S. Catarina, Nova Teutonia, 300–500 m, Nov. 1958, Plaumann leg.	Paratype	Brignoli	524	
<i>Typhlonesticus morisii</i> (Brignoli, 1975)	1 ♂, ITALY, Piemonte, Cuneo, Ali Marittime, Sotterranei presso Vernante, 16. Oct. 1973, Morisi leg. 1 ♂, 5 ♀♀, same data as the holotype, 14. Oct. 1972; 3 ♀♀, 10. Feb. 1973; 1 ♀, 17. Sep. 1974, all Morisi leg.	Holotype Paratypes	Brignoli	524	as <i>Nesticus morisii</i>
<b>Ochyroceratidae</b>					
<i>Ochyrocera fagei</i> Brignoli, 1974	1 ♂, MEXICO, Chiapas, Teopisca, Cueva del Panteon, 1820 m, 19. Mar. 1971, Argano, Sbordoni & Zullini leg.	Holotype	Brignoli	545	
<i>Ochyrocera viridissima</i> Brignoli, 1974	9 ♂, 12 ♀, same data as the holotype, Argano, Sbordoni & Zullini leg.	Paratypes			
<b>Oonopidae</b>					
<i>Camptosaphiella strepens</i> Brignoli, 1976	1 ♀, NEPAL, Zwegrhodendron, Bestrande, beim Basislager, Yaral bei Pangpoche, 3900 m, 7. May 1961, Janetschak leg.	Paratype	Brignoli	477	
<i>Oonopinus ionicus</i> Brignoli, 1979	1 ♂, 1 ♀, GREECE, Zakynthos, Vasilikon, 22. Apr. 1972, Hauser leg.	Paratypes	Brignoli	477	
<i>Oonopoides zullinii</i> Brignoli, 1974	1 ♀, MEXICO, Chiapas, Ocozocantla, Rancho del Cielito, Cueva de Cerro Brugio, 1330 m, 22. Mar. 1971, Zullini leg.	Holotype	Brignoli	545	
<i>Toloonops chickeringi</i> (Brignoli, 1974)	1 ♂, MEXICO, Chiapas, Teopisca, Cueva del Panteon, 1820 m, 19. Mar. 1971, Sbordoni leg. 1 ♀, same data as the holotype; 1 ♀, same locality, 19. Apr. 1971, Argano leg.	Holotype Paratypes	Brignoli	545	as <i>Oonops chickeringi</i>
				477, 545	

Taxon	Data	Type	Collection	Jar/tube N°.	Notes
<b>Oxyopidae</b>					
<i>Oxyopes toschii</i> Caporiacco, 1949	1 ♂, 1 ♀, KENYA, Elmenteita, 100 km North of Nairobi, 1500 m, Dec. 1945, Toschi, Meneghetti leg.; 1 ♀, Nairobi, 1500 m, 1944, Toschi, Meneghetti leg.	Syntypes	Di Caporiacco	29	
<b>Pacullidae</b>					
<i>Lamania sheari</i> (Brignoli, 1980)	1 ♂, INDONESIA, Sulawesi, Seltan, Londa, station 10, 24. Feb. 1979, Brignoli leg.	Holotype	Brignoli	59	as <i>Paculla sheari</i>
<b>Pholcidae</b>					
<i>Anopsicus hanakash</i> (Brignoli, 1974)	1 ♀, GUATEMALA, Rio Usumacinta, Env. Yaxchilan, 1230 m, 12. Mar. 1971, Zullini leg.	Holotype	Brignoli	531	as <i>Pholcophora hanakash</i>
<i>Hoploholcus asiaeminoris</i> Brignoli, 1978	1 ♂, TURKEY, Vil. Konya, Hadim, Cikkigi, cave, 1700 m, 25. Apr. 1973, Sbordoni leg.	Holotype	Brignoli	528	several vials
	13 ♂ and 3 ♀, same data as the holotype, Sbordoni, Agnoletti, Forestiero leg.	Paratypes			
<i>Metagonia maximiliani</i> Brignoli, 1972	1 ♂, MEXICO, Quaretarro, El Jobo, Cueva del Madrono cave, 1830 m, 17. Nov. 1969, Argano, Sbordoni leg.	Holotype	Brignoli	491	
	2 ♂♂, 9 ♀♀, MEXICO, Quaretarro, El Jobo, Cueva del Madrono cave, 1830 m, 17. Nov. 1969, Argano, Sbordoni leg.	Paratypes		491, 530	several vials
<i>Metagonia strinatii</i> (Brignoli, 1972)	1 ♀, ARGENTINA, Mendoza, Cueva de las Brujas, 9. Feb. 1968, Strinati leg.	Paratype	Brignoli	530	as <i>Spermophora strinatii</i>
<i>Pholcus maronita</i> Brignoli, 1977	1 ♀, LEBANON, Cazabeharri, Hassoroun, 1500 m, 2. Jun. 1972, Brignoli leg.	Holotype	Brignoli	529	
	1 ♀, same data as the holotype	Paratype			
<i>Pholcus spasskyi</i> Brignoli, 1978	1 ♂, 2 ♀♀, TURKEY, Vil. Diyarbakir, Korkha, Lice cave, 23. Jun. 1968, Brignoli, Sbordoni leg.	Paratypes	Brignoli	529	
<b>Phyxelididae</b>					
<i>Phyxelida carcharata</i> Griswold, 1990	1 ♂, KENYA, Mau, 150 km North-West of Nairobi, 2500–3000 m, Gan. 1946, Toschi, Meneghetti leg.	Holotype	Di Caporiacco	6	as <i>Haemilla nebulosa</i>
<b>Pimoidae</b>					
<i>Pimoa thaleri</i> Trotta, 2009	1 ♂, INDIA, West Sikkim, Bakim, Bakim cave, 2510 m, 12. Apr. 1994, De Pasquale, Latella, Sbordoni leg.	Holotype	Generale	94	
<b>Psilodercidae</b>					
<i>Leclercera machadoi</i> (Brignoli, 1973)	1 ♀, NEPAL, Dhorpatan, Nadelwald, 3000–3200 m, Apr. 1970, Martens leg.	Paratype	Brignoli	59	as <i>Altephus machadoi</i>
<i>Merizocera brincki</i> Brignoli, 1975	1 ♂, SRI LANKA, Deerwood, Sabaragamuwa, Kuruwita, NNW Ratnapura, loc. 90, 18–21. Feb. 1962, leg.	Paratype	Brignoli	74	
<b>Salticidae</b>					
<i>Bianor ghigii</i> (Caporiacco, 1949)	1 ♂, 1 ♀, KENYA, Elmenteita, 100 km North of Nairobi, 1500 m, Dec. 1945, Toschi, Meneghetti leg.	Syntypes	Di Caporiacco	138	as <i>Modunda ghigii</i>
<i>Brancus hemmingi</i> Caporiacco, 1949	1 ♀, KENYA, Mau, 150 km North-West of Nairobi, 2500–3000 m, Jan. 1946, Toschi, Meneghetti leg.	Syntype	Di Caporiacco	123	
<i>Brancus poecilus</i> Caporiacco, 1949	6 ♂♂, 29 ♀♀, KENYA, Nairobi, 1500 m, 1944–May 1945, Toschi, Meneghetti leg.; 2 ♀♀, Elmenteita, 100 km North of Nairobi, 1500 m, Dec. 1945, Toschi, Meneghetti leg.	Syntypes	Di Caporiacco	123	
<i>Copocrossa albozonata</i> Caporiacco, 1949	1 ♀, KENYA, Mau, 150 km North-West of Nairobi, 2500–3000 m, Jan. 1946, Toschi, Meneghetti leg.	Syntype	Di Caporiacco	119	
<i>Dioleinus angustipes</i> Gardzińska & Źabka, 2006	1 ♂, INDONESIA, Irian Jaya, Biak, Bosnik, 13. Jul. 1984, Brignoli leg.	Holotype	Brignoli	236	

Taxon	Data	Type	Collection	Jar/tube N°.	Notes
<i>Diolenius insignitus</i> Gardzińska & Źabka, 2006	1 ♀, INDONESIA, Maluku, Ambon, Liang, 16–30. Jan. 1979, Brignoli leg. 3 ♀♀, same data as the holotype	Holotype Paratypes	Brignoli	237	
<i>Euophrus megastyla</i> Caporiacco, 1949	2 ♂♂, KENYA, Mau, 150 km North-West of Nairobi, 2500–3000 m, Jan. 1946, Toschi, Meneghetti leg.; 1 ♂, Elmenteita, 100 km North of Nairobi, 1500 m, Dec. 1945, Toschi, Meneghetti leg.	Syntypes	Di Caporiacco	135	as <i>Euophrus megastylus</i>
<i>Hispo continentalis</i> (Caporiacco, 1949)	1 juv., KENYA, Nairobi, 1500 m, 1944, Toschi, Meneghetti leg	Syntype	Di Caporiacco	118	as <i>Astaenorcheses continentalis</i> ; nomen dubium
<i>Hyllus multiaculeatus</i> Caporiacco, 1949	1 ♂, KENYA, Nairobi, 1500 m, 1944, Toschi, Meneghetti leg.	Syntype	Di Caporiacco	121	
<i>Langona minima</i> Caporiacco, 1949	1 ♀, KENYA, Nairobi, 1500 m, 1944, Toschi, Meneghetti leg.	Syntype	Di Caporiacco	130	
<i>Lyssomanes michae</i> Brignoli, 1984	1 ♂, MONTSERRAT, Soufrière, 11. Jul. 1981, Hilov leg.	Holotype	Brignoli	458	
<i>Menemerus niger</i> Caporiacco, 1949	1 ♀, same data as the holotype, 11. Jul. 1981, 1 ♀, 12. Jul. 1981, all Brignoli leg. 1 juv., KENYA, Nairobi, 1500 m, 1944, Toschi, Meneghetti leg.	Paratypes			
<i>Myrmarachne exultans</i> Caporiacco, 1949	1 ♀, KENYA, Mau, 150 km North-West of Nairobi, 2500–3000 m, Jan. 1946, Toschi, Meneghetti leg.	Syntype	Di Caporiacco	117	Nomen dubium
<i>Natta splendidissima</i> (Caporiacco, 1949)	1 ♀, KENYA, Nairobi, 1500 m, 1944, Toschi, Meneghetti leg.	Syntype	Di Caporiacco	134	as <i>Cyllobelus splendidissimus</i> ; nomen dubium
<i>Neaetha alborufula</i> Caporiacco, 1949	1 ♂, KENYA, Nairobi, 1500 m, 1944, Toschi, Meneghetti leg.	Syntype	Di Caporiacco	132	
<i>Pachyballus flavipes aurantius</i> Caporiacco, 1949	1 ♀, 1 juv., KENYA, Mau, 150 km North-West of Nairobi, 2500–3000 m, Jan. 1946, Toschi, Meneghetti leg.	Syntype	Di Caporiacco	120	
<i>Pseudicius elmenteitae</i> Caporiacco, 1949	1 ♀, KENYA, Nairobi, 1500 m, 1944, Toschi, Meneghetti leg.; 1 ♀, Elmenteita, 100 km North of Nairobi, 1500 m, Dec. 1945, Toschi, Meneghetti leg.	Syntypes	Di Caporiacco	137	
<i>Stenaelurillus albopunctatus</i> Caporiacco, 1949	1 ♂, 1 ♀, KENYA, Nairobi, 1500 m, 1944, Toschi, Meneghetti leg.; 1 ♀, Elmenteita, 100 km North of Nairobi, 1500 m, Dec. 1945, Toschi, Meneghetti leg.	Syntypes	Di Caporiacco	131	
<i>Thyene nigriceps</i> (Caporiacco, 1949)	1 ♀, KENYA, Nairobi, 1500 m, 1944, Toschi, Meneghetti leg.	Syntype	Di Caporiacco	126	as <i>Paramodunda nigriceps</i>
<b>Segestriidae</b>					
<i>Ariadna brignolii</i> Wunderlich, 2011	1 ♂, ITALY, Isola di Ponza, Scoglio Rosso di Pilato, 21. Jul. 1968, Utzeri leg. 2 ♀♀, same data as the holotype.	Holotype Paratypes	Brignoli	64	
<i>Segestria sibordoni</i> Brignoli, 1984	1 ♂, GREECE, Crete, Hania, Afrata, Kramiri cave, 3 m, 12. Aug. 1974, Sbordoni leg. 1 ♂, same data as the holotype	Holotype Paratype	Brignoli	64	
<b>Telemidae</b>					
<i>Apneumonella jacobsoni</i> Brignoli, 1977	1 ♀, INDONESIA, Sumatra, Fort de Kock, 920 m, 1926, Jacobson leg.	Paratype	Brignoli	533	
<i>Seychellia lodoiceae</i> Brignoli, 1980	1 ♀, SEYCHELLES, Praslin, Vallée da Mai, 25. May 1972, Benoit, Van Mol leg.	Paratype	Brignoli	59	

Taxon	Data	Type	Collection	Jar/tube N°.	Notes
<b>Tetrablemmidae</b>					
<i>Brignoliella bicornis</i> (Simon, 1893)	1 ♂, 1 ♀, PHILIPPINES, Manilla, Antipolo, Simon leg.	Paralecto-types	Brignoli	59	as <i>Paculla bicornis</i>
<i>Brignoliella tao</i> Ballarin & Yamasaki, 2020	3 ♂♂, 4 ♀♀, TAIWAN, Taitung County, Orchid Island (Lanyu), Hongtou neighborhood, along the trail to Datianchi lake, 106 m, rainforest litter, 13. Jun. 2019, Ballarin leg.	Paratypes	Generale	134	
<i>Monoblemma becki</i> Brignoli, 1978	1 ♂, BRAZIL, Manaus, Reserva Ducke, 14. Apr. 1966, Beck leg.	Paratype	Brignoli	59	
<i>Mariblemma pandani</i> (Brignoli, 1978)	1 ♂, 1 ♀, SEYCHELLES, Silhouette, Mare aux Cochons, 500 m, 3–4. Jul. 1972, Benoit, Van Mol leg.	Paratypes	Brignoli	59	as <i>Paculla pandani</i>
<i>Tetrablemma benoiti</i> (Brignoli, 1978)	1 ♀, SEYCHELLES, La Digue, Mt. La Digue, 300 m, 28–31. Jul. 1972, Benoit, Van Mol leg.	Paratype	Brignoli	59	as <i>Hexablemma benoiti</i>
<b>Tetragnathidae</b>					
<i>Tetragnatha kolosvaryi</i> Caporiacco, 1949	1 ♂, 2 ♀♀, KENYA, Nairobi, 1500 m, 1944, Toschi, Meneghetti leg.; 1 ♂, 1 ♀, Elmenteita, 100 km North of Nairobi, 1500 m, Dec. 1945, Toschi, Meneghetti leg.; 1 ♀, KENYA, Mau, 150 km North-West of Nairobi, 2500–3000 m, Jan. 1946, Toschi, Meneghetti leg.	Syntypes	Di Caporiacco	34	
<i>Tetragnatha roeweri</i> Caporiacco, 1949	1 ♂, KENYA, Nairobi, 1500 m, 1944, Toschi, Meneghetti leg.	Syntype	Di Caporiacco	34	
<i>Tetragnatha zangherii</i> (Caporiacco, 1926)	1 ♂, ITALY, Emilia–Romagna, Portico di Romagna, riverbed, 1. Jul. 1923, Zangheri leg.	Syntype	Zangheri	4961	as <i>Eucta zangheri</i>
<b>Theridiidae</b>					
<i>Argyrodes exlineae</i> (Caporiacco, 1949)	1 ♀, KENYA, Kabete, 13 km West of Nairobi, savannah, 1500 m, Aug. 1946, Toschi, Meneghetti leg.	Syntype	Di Caporiacco	60	as <i>Conopistha exlineae</i>
<i>Argyrodes kratochvili</i> (Caporiacco, 1949)	2 ♀♀, KENYA, Nairobi, Parkboma area, 1500 m, 1944–May 1946, Toschi, Meneghetti leg.; 1 ♀, Elmenteita, 100 km North of Nairobi, 1500 m, Dec. 1945, Toschi, Meneghetti leg.	Syntypes	Di Caporiacco	60	as <i>Conopistha kratochvili</i>
<i>Dipoena bristowei</i> Caporiacco, 1949	1 juv., KENYA, Nairobi, 1500 m, 1944, Toschi, Meneghetti leg.	Syntype	Di Caporiacco	54	
<i>Dipoena quadricuspis</i> Caporiacco, 1949	2 ♀♀, KENYA, Nairobi, 1500 m, 1944–1945, Toschi, Meneghetti leg.	Syntypes	Di Caporiacco	54	
<i>Episinus hickmani</i> Caporiacco, 1949	1 ♀, KENYA, Mau, 150 km North-West of Nairobi, 2500–3000 m, Jan. 1946, Toschi, Meneghetti leg.	Syntype	Di Caporiacco	57	
<i>Phoroncidia kibonotensis concolor</i> (Caporiacco, 1949)	1 ♀, KENYA, Mau, 150 km North-West of Nairobi, 2500–3000 m, Jan. 1945, Toschi, Meneghetti leg.	Syntype	Di Caporiacco	49	as <i>Ulesanis kibonotensis concolor</i>
<i>Theridion angusticeps</i> Caporiacco, 1949	1 ♀, KENYA, Mau, 150 km North-West of Nairobi, 2500–3000 m, Jan. 1946, Toschi, Meneghetti leg	Syntype	Di Caporiacco	55	
<i>Theridion brunneonigrum</i> Caporiacco, 1949	1 ♀, KENYA, Mau, 150 km North-West of Nairobi, 2500–3000 m, Jan. 1946, Toschi, Meneghetti leg	Syntype	Di Caporiacco	55	
<i>Theridion carpathium</i> Brignoli, 1984	1 ♀, GREECE, Karpathos, Martens leg.	Paratype	Brignoli	464	
<i>Theridion chamberlini</i> Caporiacco, 1949	1 ♀, KENYA, Mau, 2500–3000 m, Dec. 1946 Toschi, Meneghetti leg.	Syntype	Di Caporiacco	55	
<i>Theridion charitonowi</i> Caporiacco, 1949	1 ♀, KENYA, Mau, 150 km North-West of Nairobi, 2500–3000 m, Jan. 1946, Toschi, Meneghetti leg	Syntype	Di Caporiacco	55	
<i>Theridion convexisternum</i> Caporiacco, 1949	1 ? (only opisthosoma), KENYA, Mau, 150 km North-West of Nairobi, 2500–3000 m, Jan. 1946, Toschi, Meneghetti leg	Syntype	Di Caporiacco	55	

Taxon	Data	Type	Collection	Jar/tube N°.	Notes
<i>Theridion femoratissimum</i> Caporiacco, 1949	1 ♂, KENYA, Mau, 150 km North-West of Nairobi, 2500–3000 m, Jan. 1946, Toschi, Meneghetti leg.	Syntype	Di Capori- acco	55	
<i>Theridion mauense</i> Caporiacco, 1949	1 ♀, KENYA, Jan. 1945, Toschi, Meneghetti leg.	Syntype	Di Capori- acco	55	
<i>Theridion meneghettii</i> Caporiacco, 1949	1 ♀, KENYA, Mau, 150 km North-West of Nairobi, 2500–3000 m, Jan. 1946, Toschi, Meneghetti leg.	Syntype	Di Capori- acco	55	
<i>Theridion nasinotum</i> Caporiacco, 1949	1 ♂, KENYA, Mau, 150 km North-West of Nairobi, 2500–3000 m, Jan. 1945, Toschi, Meneghetti leg.	Syntype	Di Capori- acco	55	
<i>Theridion nigroplagiatum</i> Caporiacco, 1949	1 ♀, KENYA, Mau, 150 km North-West of Nairobi, 2500–3000 m, Jan. 1946, Toschi, Meneghetti leg.	Syntype	Di Capori- acco	55	
<i>Theridion teutanooides</i> Caporiacco, 1949	1 juv., KENYA, Mau, 150 km North-West of Nairobi, 2500–3000 m, Jan. 1946, Toschi, Meneghetti leg.	Syntype	Di Capori- acco	55	
<i>Theridion trizonatum</i> Caporiacco, 1949	1 ♀, KENYA, Mau, 150 km North-West of Nairobi, 2500–3000 m, Jan. 1946, Toschi, Meneghetti leg.	Syntype	Di Capori- acco	55	
<i>Theridion zebra</i> Caporiacco, 1949	1 juv., KENYA, Nairobi, 1500 m, 1944, Toschi, Meneghetti leg.	Syntype	Di Capori- acco	55	
<i>Theridula albonigra</i> Caporiacco, 1949	1 ♀, KENYA, Mau, 150 km North-West of Nairobi, 2500–3000 m, Jan. 1946, Toschi, Meneghetti leg.	Syntype	Di Capori- acco	57	
<i>Theridula albonigra vittata</i> Caporiacco, 1949	1 ♂, 1 ♀, KENYA, Mau, 150 km North-West of Nairobi, 2500–3000 m, Jan. 1946, Toschi, Meneghetti leg.	Syntypes	Di Capori- acco	57	
<b>Theridiosomatidae</b>					
<i>Plato guacharo</i> (Brignoli, 1972)	1 ♂, 2 ♀♀, VENEZUELA, Monjas, Cueva del Guacharo, 18. Feb. 1968, Strinati leg.	Paratypes	Brignoli	74	as <i>Wendilgarda guacharo</i>
<i>Plato miranda</i> (Brignoli, 1972)	2 ♀♀, VENEZUELA, Miranda, Cueva Alfredo Jahn, 21. Feb. 1968, Bordon, Strinati leg.	Paratypes	Brignoli	74	as <i>Wendilgarda miranda</i>
<b>Thomisidae</b>					
<i>Ozyptila salustri</i> Wunderlich, 2011	1 ♂, ITALY, South Tuscany, East Grosseto, ~15km, South-East of Roccastrada, near Campagnatico, Salustri, garden in a wine growing estate, in detritus under a bush, Aug. 2010, Wunderlich leg.	Paratype	Generale	130	
<i>Misumena pallescens</i> Caporiacco, 1949	1 ♀, KENYA, Nairobi, 1500 m, 1944, Toschi, Meneghetti leg.	Syntype	Di Capori- acco	108	
<i>Mystaria variabilis delesserti</i> (Caporiacco, 1949)	1 ♀, KENYA, Nairobi, 1500 m, 1944, Toschi, Meneghetti leg.	Syntype	Di Capori- acco	104	as <i>Paramystaria variabilis delesserti</i>
<i>Xysticus jugalis larvatus</i> Caporiacco, 1949	1 juv., KENYA, Mackinnon Road, 80 West of Mombasa, 500 m, Dec. 1946, Toschi, Meneghetti leg.	Syntype	Di Capori- acco	110	
<b>Trachelidae</b>					
<i>Orthobula milloti</i> Caporiacco, 1949	1 ♂, 1 ♀, KENYA, Nairobi, 1500 m, 1944, Toschi, Meneghetti leg.	Syntypes	Di Capori- acco	89	
<b>Zodariidae</b>					
<i>Diores delesserti</i> Caporiacco, 1949	1 ♀, KENYA, Nairobi, 1500 m, 1944, Toschi, Meneghetti leg.	Syntype	Di Capori- acco	30	
<i>Diores strandi</i> Caporiacco, 1949	1 ♂, KENYA, Nairobi, 1500 m, 1944, Toschi, Meneghetti leg.	Syntype	Di Capori- acco	31	

Taxon	Data	Type	Collection	Jar/tube N°.	Notes
<i>Systenoplacis falconeri</i> (Caporiacco, 1949)	1 ♂, KENYA, Mackinnon Road, 10 km West of Mombasa, 500 m, Dec. 1945, Toschi, Meneghetti leg.	Syntype	Di Caporiacco	32	as <i>Cydrela falconeri</i>
<i>Zodarion epirensis</i> Brignoli, 1984	1 ♀, GREECE, Epirus, Loannina, Eleousa, 28. Jul. 1967, Brignoli leg.; 2 ♀♀, around Ioannina, 2. Aug. 1966, Argano leg.	Paratypes	Brignoli	546	
<i>Zodarion ruffoi</i> Caporiacco, 1951	1 ♀, ITALY, Apulia, Foggia, Is. Tremiti, S. Domino Is., 9. May 1948, Ruffo leg.	Holotype	Generale	24	
<b>Zoropsidae</b>					
<i>Pseudoctenus meneghettii</i> Caporiacco, 1949	2 ♀♀, KENYA, Mau, 150 km North-West of Nairobi, 2500–3000 m, Jan. 1946, Toschi, Meneghetti leg.	Syntypes	Di Caporiacco	97	
<b>Opiliones</b>					
<b>Dicranolasmatidae</b>					
<i>Dicranolasma ponticum</i> Gruber, 1998	1 ♂, TURKEY, Vil. Ordu, Gurgentepe Geçidi, 1275 m, 8. Jun. 1969, Brignoli leg.	Paratype	Brignoli	566	
<b>Nemastomatidae</b>					
<i>Ortholasma sbordonii</i> Šilhavý, 1973	1 ♂, MEXICO, Tamauligas, Sierra de Guatemala, Gomez Faria, Cueva de la Terra, 2160 m, 23. Nov. 1969, Argano leg. 1 juv., same locality and data as the holotype 1 ♀, MEXICO, Tamauligas, Rancho del Cielo, Cueva de la Mina, 1600 m, 24. Nov. 1969, Sbordoni leg.	Holotype Paratype Allotype	Brignoli	57	
<b>Ricinulei</b>					
<b>Ricinoididae</b>					
<i>Pseudocellus sbordonii</i> (Brignoli, 1974)	1 ♂, MEXICO, Chiapas, Ocozocautla, Rancho del Celito, Cueva de las Canicas, 1350 m, 22. Mar. 1971, Sbordoni leg.	Holotype	Brignoli	482	as <i>Cryptocellus sbordonii</i>
<b>Schizomida</b>					
<b>Hubbardiidae</b>					
<i>Baalrog sbordonii</i> (Brignoli, 1973)	1 ♂, MEXICO, Vera Cruz, Cordoba, Paraje Nuevo, Cueva Ojo de Agua Grande, 550 m, 6. Nov. 1969, Sbordoni leg.	Holotype	Brignoli	482	as <i>Schizomus sbordonii</i>
<i>Schizomus arganoi</i> Brignoli, 1973	1 ♂, MEXICO, Chiapas, Bochil, Cueva de la Golondrina, 27. Mar. 1971, Argano leg.	Holotype	Brignoli	482	