

SUPPLEMENTAL MATERIAL

**PREDICTIVE HABITAT MODEL REVEALS SPECIFICITY IN A BROADLY
DISTRIBUTED FOREST RAPTOR, THE HARPY EAGLE**

JOSÉ DE JESÚS VARGAS GONZÁLEZ¹

The Peregrine Fund, 5668 West Flying Hawk Lane, Boise, ID 83709 USA

and

Fundación Rapaces y Bosques de Panamá, Apdo. 1015-00326, La Chorrera, Panamá Oeste

JENNIFER D. MCCABE, DAVID L. ANDERSON, AND MARTA CURTI

The Peregrine Fund, 5668 West Flying Hawk Lane, Boise, ID 83709 USA

DARISNEL CARPIO CÁRDENAS

Fundación Rapaces y Bosques de Panamá, Apdo. 1015-00326, La Chorrera, Panamá Oeste

FÉLIX HERNÁN VARGAS CASTILLO

The Peregrine Fund, 5668 West Flying Hawk Lane, Boise, ID 83709 USA

Table S1: Pearson's correlations coefficients for all predictor variables used in the model

	<i>Elevation</i>	<i>FC 250 m</i>	<i>FC 1 km</i>	<i>FC 4 km</i>	<i>Dist. lg. com.</i>	<i>Dist. to river</i>	<i>Slope</i>
<i>Elevation</i>	1	0.30	0.33	0.37	0.17	0.24	0.68
<i>FC 250 m</i>	0.30	1	0.86	0.75	0.24	0.05	0.38
<i>FC 1 km</i>	0.33	0.86	1	0.92	0.31	0.01	0.40
<i>FC 4 km</i>	0.37	0.75	0.92	1	0.35	-0.07	0.39
<i>Dist. lg. com.</i>	0.17	0.24	0.31	0.35	1	0	0.03
<i>Dist. to river</i>	0.24	0.05	0.01	-0.07	0	1	0.19
<i>Slope</i>	0.68	0.38	0.40	0.39	0.03	0.19	1