

The Arabidopsis Circadian System

Authors: McClung, C. Robertson, Salomé, Patrice A., and Michael, Todd P.

Source: The Arabidopsis Book, 2002(1)

Published By: The American Society of Plant Biologists

URL: <https://doi.org/10.1199/tab.0044>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

First published on March 27, 2002: e0044. doi: 10.1199/tab.0044

The Arabidopsis Circadian System

C. Robertson McClung¹, Patrice A. Salomé, and Todd P. Michael

Department of Biological Sciences, 6044 Gilman Laboratories, Dartmouth College, Hanover, New Hampshire 03755-3576

¹Corresponding Author: telephone: 603-646-3940; fax: 603-646-1347; email: mcclung@dartmouth.edu

ABSTRACT

Rhythms with periods of approximately 24 hr are widespread in nature. Those that persist in constant conditions are termed circadian rhythms and reflect the activity of an endogenous biological clock. Plants, including *Arabidopsis*, are richly rhythmic. Expression analysis, most recently on a genomic scale, indicates that the *Arabidopsis* circadian clock regulates a number of key metabolic pathways and stress responses. A number of sensitive and high-throughput assays have been developed to monitor the *Arabidopsis* clock. These assays have facilitated the identification of components of plant circadian systems through genetic and molecular biological studies. Although much remains to be learned, the framework of the *Arabidopsis* circadian system is coming into focus.

Dedication

This review is dedicated to the memory of DeLill Nasser, a wonderful mentor and an unwavering advocate of both *Arabidopsis* and circadian rhythms research.

INTRODUCTION

The evolution of life within the biosphere has taken place in a context that has changed continuously over time. It is common to think of this temporal change as linear. Cyanobacteria appeared 3.5 billion years ago in the Pre-Cambrian, land plants appeared 425 million years ago in the Silurian, and flowering plants appeared 135 million years ago in the Cretaceous. Undoubtedly, environmental changes played key roles in many of the stages in the evolution of modern species, including *Arabidopsis thaliana*. For example, the number and density of stomatal complexes have responded to changing atmospheric CO₂ concentration, contributing to the appearance of the planate leaf form 360 million years ago in the Late Paleozoic (Beerling et al., 2001).

Often neglected in this linear view of temporal environmental change is the periodic change in the immediate environment that results from the rotation of the earth on its axis every 24 hours. This alternation of day and night produces profound environmental changes that must be faced by any organism. Thus, life has evolved in an environment that shows short term (~24 hr) recurrent change that is drastic but that is entirely predictable. Time may fly like an arrow in the long term, but in the short term each dawn is followed by a successive dawn and time can more appropriately be thought of as having the trajectory of a boomerang rather than that of an arrow. Given this repetitive perspective, it seems intuitive that any organism would accrue advantage from the temporal organization of

biochemistry, physiology and behavior with respect to the environmental oscillation of day and night. Not surprisingly, most organisms express diurnal rhythms. It is less obvious that many of these rhythms should persist in the absence of environmental time cues. However, if one accepts the adage that “the early bird gets the worm,” it would seem to be to the advantage of the worm to anticipate the dawn and hide so as not to serve as the early bird’s breakfast. No less accomplished a naturalist than William Shakespeare described such a case of dawn anticipation in *Hamlet*, Act I, scene v: “*The glowworm shows the matin to be near And gins to pale his uneffectual fire.*” More recently, a fitness advantage conferred by the circadian clock has been demonstrated in cyanobacteria and *Chlamydomonas* (Johnson, 2001; McClung, 2001).

Organisms from cyanobacteria to humans endogenously measure time and temporally regulate aspects of their biology. There is no more obvious significance to the synchronization of the organism to local time than in the case of photosynthetic organisms that depend on sunlight for energy. Indeed, the endogenous nature of circadian rhythms was first established for the leaf movements of the “sensitive” heliotrope (probably *Mimosa pudica*) (de Mairan, 1729). Prior to 1900, plants were the primary system for the study of circadian rhythms and plant studies yielded the historical milestones of circadian rhythm research (summarized by McClung and Kay, 1994). Over the last decade, *Arabidopsis* has become the premiere system for the analysis of plant circadian rhythms and this review will focus on recent advances in our understanding of the molecular bases of *Arabidopsis* circadian rhythms.

Circadian (from the Latin *circa*, approximately, and *dies*, day) rhythms are endogenous rhythms with periods of approximately 24 hr. When placed in constant conditions and, thus, deprived of external time cues, circadian rhythms persist and “free-run” with an endogenous period that is close to but not exactly 24 hr. This establishes that these rhythms are truly endogenous and self-sustaining. In the real world, of course, organisms are exposed to environmental time cues such as light and temperature

cycles, and these cues serve to synchronize or “entrain” the endogenous organismal clock with local solar time. The period of a circadian rhythm remains relatively constant over the range of physiologically relevant temperatures, which is referred to as temperature compensation. This means that the circadian clock maintains its pace over a range of temperatures, but does not imply that temperature changes or cycles cannot serve as potent stimuli that can entrain the clock. These three properties: persistence in constant conditions with an approximately 24-hr period, entrainment and temperature compensation, are the diagnostic criteria of a circadian rhythm (Johnson et al., 1998; Sweeney, 1987).

AN OVERVIEW OF THE CIRCADIAN SYSTEM

Commonly, one considers the circadian system as consisting of three components: an input pathway (usually described as a photoreceptor) that entrains the clock, the central oscillator (clock), and output pathways to generate overt rhythms. These three components are arranged in a linear array with information flowing unidirectionally from input through the oscillator to the output (Figure 1). This review will deviate from the linear order of Figure 1 in its consideration of the *Arabidopsis* circadian system. In order to introduce the assays that feature in the analysis of plant clocks, output pathways will be considered first. Input pathways will be addressed second. Finally, the exciting recent progress in elucidating the oscillator mechanism in plants will be considered. We are frequently being reminded that the formal dissection of any circadian system into input, oscillator and output components is arbitrary and the lines between components are blurred. This will become particularly evident in the attempt to identify components of the central oscillator.

It has become increasingly clear that the simple linear of Figure 1 is inadequate and a more complex model has emerged (Figure 2). There are multiple photoreceptors that provide input to the clock (Devlin and Kay, 2001). In addition, there must be sensors for temperature and for other environmental stimuli that entrain the clock. Accumulating evidence suggests that there are multiple oscillators. These can represent central master oscillators as well as non-self-sustaining slave oscillators (Pittendrigh, 1960; Pittendrigh, 1981). In animals there are organ and tissue-specific oscillators (Plautz et al., 1997; Tosini and Menaker, 1998; Yamazaki et al., 2000). In the unicellular dinoflagellate, *Gonyaulax polyedra*, it has been established that there are at least two circadian oscillators within a single cell (Morse et al., 1994; Roenneberg and Morse, 1993).




Figure 1. Conceptual scheme illustrating simple linear information flow from input (entrainment) pathways through the central oscillator to output pathways. Modified from Eskin (Eskin, 1979)

Finally, some components of input pathways are themselves outputs and the circadian clock regulates its own sensitivity to environmental input (Devlin and Kay, 2001; Mellow et al., 1999; Mellow et al., 2001).

Rhythmic Outputs

Plants are richly rhythmic, presumably because of their dependence upon sunlight for energy and because they are sessile. Arabidopsis is no exception and displays myriad rhythmic outputs, or “hands” of the clock. Plant rhythmic processes have been reviewed in detail (Kreps and Kay, 1997; Lumsden and Millar, 1998; McClung, 2000; McClung, 2001; Millar, 1999; Somers, 1999; Sweeney, 1987). Here I will focus on those rhythms that have proven most useful in the study of the Arabidopsis clock.

Rhythms in mRNA Abundance

Differential gene expression underlies many overt rhythms in biochemistry, physiology and behavior. The cataloging of plant clock-controlled genes (CCG) began with Klopstech's (Klopstech, 1985) observation of a circadian oscillation in mRNA abundance of a chlorophyll *a/b* binding protein gene (*LHCB* or *CAB*). *LHCB* remains the best-studied clock-regulated plant gene, and circadian oscillation of *LHCB* mRNA abundance is widespread, if not universal, among angiosperms (Fejes and Nagy, 1998; Piechulla, 1999). Curiously, this does not extend into the gymnosperms (Piechulla, 1999).

The list of plant CCGs has grown to considerable length (Fejes and Nagy, 1998; Kreps et al., 2000; McClung, 2000; McClung, 2001; Somers, 1999). Happily, the genomic age has greatly facilitated this analysis with the application of both oligo-based and cDNA microarrays (Harmer et al., 2000; Schaffer et al., 2001). The cDNA microarray experiment suggested that ~11% of the 7800 genes sampled showed diurnal oscillations in mRNA abundance in a light–dark cycle and that oscillations for 2% persisted in continuous light and so were truly circadian (Schaffer et al., 2001). The oligo-based microarray, with better time resolution (4 hr versus 6 hr sampling intervals, and encompassing two versus a single circadian cycle), allowed the detection of statistically significant circadian (in continuous light) oscillations in mRNA abundance of 5–6% of the 8200 genes examined (Harmer et al., 2000). This latter meas-

urement suggests that there are at least 1275–1530 Arabidopsis CCGs, based on a current estimate of ~25,500 Arabidopsis (The Arabidopsis Genome Initiative, 2000). More recent cDNA microarray work, in which hybridization results were normalized by hybridization against genomic DNA, is more consistent with this higher estimate (Schaffer and Wisman, 2001). This is likely to represent a minimum estimate, limited to the detection of oscillating transcripts in those tissues, at that developmental stage, and under those specific growth conditions employed. It will take some time and many chips to exhaustively sample all possible developmental stages and environmental conditions. In addition, one must recall that transcript abundance per se does not distinguish between changes in transcription rates and mRNA stability. Nor will these experiments identify genes whose induction or repression in response to environmental or biological stimuli is gated by the clock. Nonetheless, 5–6% is considerably less than the situation encountered in the cyanobacterium *Synechococcus elongatus*, in which essentially the entire transcriptome is clock-regulated (Liu et al., 1995).

What genes are clock-regulated? Not unexpectedly, microarray analysis confirms that many genes associated with photosynthetic light harvesting oscillate synchronously with peaks in mRNA abundance after subjective dawn (Harmer et al., 2000; Schaffer et al., 2001). These include genes encoding LHCA and LHCB proteins as well as photosystem I and II reaction center proteins. A number of Arabidopsis genes involved in photosynthetic carbon assimilation (Pilgrim and McClung, 1993) as well as in photorespiration (McClung et al., 2000; Zhong and McClung, 1996; Zhong et al., 1994) are also clock-regulated. Microarray experiments have extended the list of clock-regulated genes involved in carbon metabolism. For example, mRNA abundance for a number of genes encoding enzymes of the glycolytic and oxidative pentose phosphate pathways, as well as genes encoding hexose transporters peak in the subjective afternoon. Genes encoding enzymes of starch mobilization, including starch kinase, β -amylase, fructose-bisphosphate aldolase, and sugar transporters peak at night (Harmer et al., 2000).

It has long been known that nitrogen assimilation is clock-regulated (Cohen and Cumming, 1974). Circadian oscillation in mRNA abundance of genes encoding nitrate reductase has been demonstrated, first in tobacco (Deng et al., 1990; Deng et al., 1989) and also in Arabidopsis (Cheng et al., 1991; Pilgrim et al., 1993). Microarray analysis has verified these observations (Schaffer et al., 2001) and extended them to include other genes with roles in nitrogen assimilation, including putative carboxypeptidases, amidase, glutamate dehydrogenase, and asparagine synthetase, as well as nitrate and ammonium transporters (Harmer et al., 2000). There is also a clock-regulated

group of genes involved in sulfur metabolism (Harmer et al., 2000).

Plant responses to biotic and abiotic stress responses are often gated by the circadian clock (Rikin, 1992; Rikin et al., 1993; Rikin et al., 1984). Both microarray experiments

identified oscillations in mRNA abundance of a number of genes involved in responses to stresses, including cold and pathogens (Harmer et al., 2000; Schaffer et al., 2001). Particularly noteworthy is the oscillation in mRNA abundance of the *DREB1a/CBF3* gene encoding a transcription




Figure 2. A more realistic model of a simple circadian system consisting of a set of input (entrainment) pathways, multiple central oscillators, and sets of output pathways. Entraining stimuli include light, mediated through phytochromes (PHY) and cryptochromes (CRY), temperature, and imbibition (not shown). Complexity in input pathways arises from multiple phytochromes and cryptochromes as well as interaction among them and their downstream signaling pathways. Each central oscillator is illustrated as a loop including positive and negative components that yields a self-sustaining oscillation with a period of approximately 24 hr. Coupling between oscillators is suggested by double-headed arrows. Multiple output pathways are drawn as each regulating an overt rhythm with a distinct phase. Some outputs may be driven by individual oscillators whereas others may receive input from more than one oscillator. Additionally, different oscillators may drive separate rhythms with distinct periods.

factor that plays a key role in cold tolerance (Thomashow et al., 2001). Rhythmic expression of this transcription factor could underlie a circadian rhythm in cold tolerance (Harmer et al., 2000). The downstream effector genes themselves would not necessarily exhibit circadian oscillations in mRNA abundance, but their induction might be gated by the clock through circadian regulation of key sensors or transcription factors.

One of the most interesting observations stemming from microarray analysis was that 23 genes encoding enzymes of phenylpropanoid biosynthesis are coordinately regulated, oscillating with mRNA peaks about 4 hr before subjective dawn (Harmer et al., 2000). Oscillating together with these genes is *PRODUCTION OF ANTHOCYANIN PIGMENT 1* (*PAP1*), which encodes a Myb domain transcription factor. *PAP1* overexpression up-regulates several enzymes of the pathway (Borevitz et al., 2000). This offers the likely scenario that the clock regulates *PAP1* expression which in turn regulates the entire phenylpropanoid biosynthetic pathway. This is a remarkable illustration of the power of genome-wide expression analysis to show not only that an entire pathway is co-regulated, but also to suggest the key transcription factor that mediates this regulation.

Although most genes exhibiting circadian oscillations are nuclear, there is accumulating evidence of circadian regulation of plastid-encoded genes of *Chlamydomonas* (Hwang et al., 1996; Salvador et al., 1993) and wheat (Nakahira et al., 1998). This oscillation of the wheat *psbD* gene is dependent on an atypical -35 promoter element and it was hypothesized that transcription of this gene requires a plastid-encoded RNA polymerase and a sigma factor encoded by a nuclear CCG (Nakahira et al., 1998). Indeed, circadian transcription of nuclear-encoded sigma factor genes has been shown in *Arabidopsis* and wheat (Kanamaru et al., 1999; Morikawa et al., 1999). This provides a novel mechanism for temporal coordination between the nuclear and plastid genomes.

Circadian Transcription

Circadian regulation of transcription underlies many of the rhythms in mRNA abundance described above. For example, both nuclear run-on experiments and transcriptional gene fusions have established a transcriptional component to the circadian regulation of *LHCB* genes in several angiosperms (Fejes and Nagy, 1998; Piechulla, 1999). The instability of luciferase (LUC) activity *in planta* means that light production requires translation of new LUC protein and provides a reliable assessment of *LUC* transcrip-

tion (Millar et al., 1992a). Moreover, the measurement of LUC activity is non-destructive and quantitative and allows both temporal and spatial resolution of gene expression in real time *in vivo*. Thus, LUC has become the reporter of choice for monitoring circadian gene expression (Figure 3; Millar et al., 1992a; Millar et al., 1992b).

Minimal nuclear promoters sufficient to confer maximal circadian transcription at a mid-morning phase have been identified for several *Arabidopsis* genes, including *LHCB* (Fejes and Nagy, 1998; Piechulla, 1999) and *RCA* (Liu et al., 1996). In both cases, transcription is well correlated with the mid-morning phase of peak mRNA abundance. The microarray experiments emphasize that the mRNA abundance of different genes occurs at distinct circadian phases (times of day) and a number of different phase angle markers are available (Harmer et al., 2000; Schaffer et al., 2001). For example, mRNA abundance of the *CAT2* and *CAT3* catalase genes of *Arabidopsis* peaks at dawn and dusk, respectively (Zhong and McClung, 1996) and we have defined minimal promoter sequences sufficient to confer dawn- and evening-specific circadian transcription on a *LUC* reporter (TP Michael, PA Salomé & CR McClung, unpublished data; see Figure 3). Evening-specific promoters have also been defined for the *Arabidopsis* genes encoding a glycine-rich RNA-binding protein (*ATGRP7/CCR2*) and a germin-like protein (*AtGER3*) (Staiger and Apel, 1999; Staiger et al., 1999; Strayer et al., 2000). Both mRNA abundance and transcription of *EARLY FLOWERING 3* (*ELF3*) peak about 4 hr after subjective dusk (Covington et al., 2001; Hicks et al., 2001). It is now pos-




Figure 3. Rhythmic gene transcription, monitored as LUC activity in seedlings carrying *promoter::LUC* transgenes to illustrate phase-specific transcription. Seedlings (Col ecotype) carrying either *CAT2::LUC* (red triangles), *LHCB::LUC* transgene (blue squares) or a *CAT3::LUC* transgene (cyan circles) were entrained to a light-dark (12:12) cycle and released into continuous light at T=0. The peak in *CAT2::LUC* activity occurs at subjective dawn, in *CAT2::LUC* activity at mid-day, and in *CAT3::LUC* activity at subjective dusk. Gray boxes indicate subjective night.

sible to target the expression of one's favorite gene to any particular time of day (Table 1).

The definition of minimal promoters both necessary and sufficient to confer phase-specific transcription represents a key step in the analysis of the mechanisms by which genes are transcribed at different times of day. The experiments have chiefly begun with progressive 5' deletion analysis, followed by site-directed mutagenesis to test the functional relevance of potential clock responsive elements. For example, *in vivo* functional analysis of progressively truncated *LHCB1*1* (*CAB2*) promoter fragments fused to *LUC* defined a 36 bp region sufficient to confer circadian transcription that was bound, *in vitro*, by multiple complexes (Carré and Kay, 1995). The *CIRCADIAN CLOCK ASSOCIATED 1* (*CCA1*) gene encodes a single Myb domain protein that shows circadian binding to an element (consensus AAAa/cAATCT) within the functionally defined region of the *LHCB1*3* (*CAB1*) promoter (Wang and Tobin, 1998). This *CCA1*-binding element is also found in the minimal *RCA* promoter (Liu et al., 1996), although the functional importance of *CCA1* binding to the *RCA* promoter has not been established. A related consensus, AAAATATCT, is also found in the functionally defined minimal evening-specific *AtGRP7/CCR2* (Staiger and Apel, 1999) and *CAT3* promoters (TP Michael & CR McClung, unpublished data), as well as in the putative pro-

motors of 31 evening-specific genes identified by microarray analysis (Harmer et al., 2000). Mutation of this element eliminates evening specific transcription of *CCR2* (Harmer et al., 2000) and *CAT3* (TP Michael & CR McClung, unpublished data). It seems surprising that as little as a single nucleotide change in a consensus element might result in transcription nearly 180° out of phase, although the functional significance of this T/A substitution has not yet been established. It seems quite likely that flanking sequences, perhaps themselves the targets of other DNA-binding proteins, may contribute to the establishment of phase. Certainly, it seems that the mechanism by which the phase of transcription is determined will not necessarily be the simple solution of a series of phase-specific transcriptional activators, as implied by the apparent co-regulation of the phenylpropanoid pathway by PAP1 (Borevitz et al., 2000; Harmer et al., 2000).

Movement and growth rhythms

The initial written description of a plant circadian rhythm, by Androstenes in the 4th century BC, described the diur-

Table 1. Clock regulated promoters drive circadian transcription at phases that span the circadian day. Shown are those promoters that have been experimentally shown to drive rhythmic transcription in transgenic plants. Phases are defined in Circadian Time (CT), which is a formalism to allow comparison of the circadian phasing of rhythms for which the period varies from 24 hr. $CT = 24 \times \text{phase/period}$.

Phase	Gene(s)	Reference
CT0	<i>CIRCADIAN CLOCK ASSOCIATED 1</i> (<i>CCA1</i>)	(Covington et al., 2001)
	<i>CATALASE 2</i> (<i>CAT2</i>)	PA Salomé & CR McClung, unpublished
CT4	<i>LHCB</i>	(Millar et al., 1992a)
	<i>RUBISCO ACTIVASE</i> (<i>RCA</i>)	(Liu et al., 1996)
	<i>PHYTOCHROME B</i> (<i>PHYB</i>)	(Bognár et al., 1999)
CT8	<i>COLD AND CIRCADIAN REGULATED 2</i> (<i>CCR2</i>)	(Strayer et al., 2000)
CT12	<i>TIMING OF CAB 1</i> (<i>TOC1</i>)	(Golden and Strayer, 2001)
	<i>CATALASE 3</i> (<i>CAT3</i>)	TP Michael & CR McClung, unpublished
CT16	<i>EARLY FLOWERING 3</i> (<i>ELF3</i>)	(Covington et al., 2001)
CT20	<i>CHALCONE SYNTHASE</i> (<i>CHS</i>)	(Thain et al., 2000)

nal leaf movements of the tamarind (*Tamarindus indicus*) observed during the marches of Alexander the Great (Moore-Ede et al., 1982). In this classic system of pulvinar leaf movements, cells in the extensor and flexor regions of the pulvinus swell in antiphase (180° out of phase) to drive a circadian oscillation in leaf position (Engelmann and Johnsson, 1998). These leaf movements may have adaptive value in regulating perception of photoperiodic light signals (Bünning, 1971).

Although *Arabidopsis* lacks the pulvinus, it displays rhythms in cell elongation and, thus, in growth rate. For example, there is a circadian rhythm in the elongation rate of the abaxial and adaxial cells of the petiole that confers an oscillation in position of cotyledons and leaves (Engelmann and Johnsson, 1998). Leaf movements of individual seedlings are easily monitored by video imaging (Engelmann et al., 1992; Millar et al., 1995a). Figure 4 illustrates the use of this leaf (more properly, cotyledon) movement assay to compare period lengths in wild type (ecotype C24) versus *timing of CAB expression 1 (toc1)* mutant seedlings, which exhibit a shortened period. The still images of a single C24 seedling presented in Figure 4B were derived from a time lapse video.

In addition to the growth rhythm in petioles, *Arabidopsis* also exhibits circadian rhythms in the rate of hypocotyl elongation (Dowson-Day and Millar, 1999). There is also a circadian rhythm in the elongation rate of inflorescence stem that is correlated with the level of indole-3-acetic acid (IAA) in rosette leaves, although IAA levels in the inflorescence stem do not oscillate (Jouve et al., 1998). There does not seem to be rhythmic synthesis of IAA in the shoot apex. Rather, there is either a rhythm either in polar transport of IAA or in the ability to elongate in response to IAA, because decapitation abolishes elongation that is restored by application of IAA (Jouve et al., 1999). Inhibition of IAA polar transport blocks elongation, but this does not distinguish between either rhythmic IAA transport or sensitivity as critical for the overt rhythm in elongation rate. Microarray analysis has shown circadian rhythms in mRNA abundance of two auxin transporter genes, *PIN3* and *PIN7* (Harmer et al., 2000), which would be consistent with rhythmic auxin transport. Additionally, transcript abundance for a putative expansin, a putative polygalacturonase and an aquaporin also oscillate, peaking towards the end of subjective day, with wall-synthesis genes coordinately peaking towards the end of the subjective night (Harmer et al., 2000). Collectively, these observations suggest that the movement of the hormone that signals elongation as well as the ability of the plant cell to expand in response to that signal may be regulated by the clock. However, this remains to be directly established.

The issue of possible circadian regulation of phytohormone production and responsiveness has been largely ignored. Circadian regulation of auxin transport is implied.

In addition, *AUXIN RESPONSE TRANSCRIPTION FACTOR 3 (ETTIN/ARF3)* exhibits mRNA oscillations in continuous light (Harmer et al., 2000), which suggests that the ability to respond to auxin will also prove to be gated by the clock. Ethylene production exhibits circadian rhythmicity in a number of species (Finlayson et al., 1998; Levinsh and Kreicbergs, 1992). In sorghum, 1-aminocyclopropane-1-carboxylic acid (ACC) oxidase mRNA abundance and activity are rhythmic (Finlayson et al., 1999). mRNA abundance oscillations have been established by microarray analysis for two *Arabidopsis* ACC oxidase genes (Harmer et al., 2000). There is also a diurnal oscillation in gibberellic acid levels in sorghum, although it has not been established that this rhythm persists in continuous conditions (Foster and Morgan, 1995).

Stomatal aperture, gas exchange and CO₂ assimilation

Like many plants, *Arabidopsis* exhibits a circadian rhythm in stomatal aperture (Webb, 1998). This is correlated with a circadian rhythm in the rate of CO₂ assimilation (P.A. Salomé, E.V. Kearns & C.R. McClung, unpublished). In beans there is circadian control of Calvin cycle reactions in addition to control of stomatal aperture and gas exchange (Hennessey and Field, 1991), but circadian regulation of the Calvin cycle has not been investigated in *Arabidopsis*. However, microarray analysis has indicated circadian oscillations in mRNA levels for a number of genes encoding Calvin cycle enzymes (Harmer et al., 2000). The molecular basis of the circadian rhythm of CO₂ assimilation in Crassulacean Acid Metabolism (CAM) plants is understood in considerable detail (Nimmo, 2000). Flux through PEP carboxylase (PEPc) is regulated by reversible phosphorylation in which the night (phosphorylated) form of PEPc is less sensitive to inhibition by malate. PEPc kinase from *Kalanchoë fedtschenkoi* is atypical in that it lacks regulatory domains and thus is not regulated by second messengers. Rather, rhythmic PEPc kinase activity reflects rhythmic transcript accumulation and *de novo* protein synthesis (Hartwell et al., 1999; Hartwell et al., 1996). The CAM CO₂ assimilation rhythm also entails metabolite compartmentalization in that there is a rhythm in the transport of malate across the tonoplast (Nimmo, 2000).




Figure 4. *toc1-1* shows a short period in leaf (cotyledon) movement. **(A)** Three images (initial, middle and last) of a video clip that can be reached by double-clicking on the image. Twelve *toc1-1* (red) and twelve isogenic wild type C24 ecotype (blue) seedlings were imaged for seven circadian cycles. **(B)** Images of a single C24 seedling [outlined by the blue box in the middle and right panels of **(A)**] captured at 90 min intervals across a single circadian cycle. **(C)** Quantification of cotyledon position from the images shown in **(A)** for *toc1-1* (red) and C24 (blue). The data represent the average for the two cotyledons of the seedlings indicated by the boxes in the middle and right panels of **(A)**. The shortened period of *toc1-1* is evident. Seedlings were entrained to a light-dark (12:12) cycle and released into continuous light at T=0. Gray boxes indicate subjective night as defined by the entraining cycle.

Calcium

Ca^{2+} is a ubiquitous second messenger in plant signaling pathways (Sanders et al., 1999). The relevance of Ca^{2+} signaling to circadian rhythmicity has been experimentally established; for example, application of either Ca^{2+} or a Ca^{2+} ionophore phase shifts the leaflet movement rhythm of *Robinia pseudoacacia* (Gómez and Simón, 1995). As monitored by aequorin luminescence in both tobacco and Arabidopsis, free cytosolic but not nuclear Ca^{2+} levels oscillate with a circadian rhythm (Johnson et al., 1995; Wood et al., 2001). Wood et al. (2001) looked at Ca^{2+} levels in different tissues by expression of apoaequorin from several tissue-specific promoters. Interestingly, the peaks of the Ca^{2+} oscillations occur at different phases in different tissues (Wood et al., 2001).

Microarray analysis shows circadian transcript oscillations for a number of genes associated with Ca^{2+} signaling, including genes encoding Calmodulin and a Calmodulin-like Ca^{2+} -binding protein, as well as a putative Ca^{2+} -binding EF-hand protein and a Ca^{2+} -transporting ATPase (Harmer et al., 2000). Ca^{2+} is important in guard cell signaling (Leckie et al., 1998; Schroeder et al., 2001) and is likely to be involved in the circadian regulation of stomatal aperture and gas exchange. Ca^{2+} is also implicated in red and blue light signal transduction (Barnes et al., 1997; Baum et al., 1999; Frohnmeyer et al., 1998; Long and Jenkins, 1998; Guo et al., 2001) and may play a role in the entrainment of the circadian oscillator. The light to dark transition stimulates a spike in chloroplast stromal Ca^{2+} levels (Johnson et al., 1995), although whether this signals the circadian clock is not known. Thus, Ca^{2+} is likely to play multiple roles in the circadian system, but none of these roles are yet well defined.

Photoperiodism

The timing of flowering in many species is photoperiodic (Lin, 2000a; Simpson et al., 1999). Bünning (Bünning, 1936) hypothesized that circadian timekeeping was essential for photoperiodic time measurement and many mutations that affect circadian rhythms in gene expression and leaf movement also affect flowering timing (Lin, 2000a; Simpson et al., 1999). Alteration of the activity of a number of components of the Arabidopsis circadian system through overexpression or through loss of function (whether by antisense or mutation), often alters flowering time (Table 2). For example, mutations in two components

of the photoperiodic pathway, *early flowering 3 (elf3)* and the late flowering *gigantea (gi)*, confer defects in the circadian timing and define components of the light input pathway. Overexpression of *LATE ELONGATED HYPOCOTYL (LHY)*, a third component of the photoperiodic pathway, results in arrhythmicity as well as late flowering (Schaffer et al., 1998). However, overexpression of *CONSTANS (CO)*, a CCG that accelerates flowering in response to long days, does not disrupt circadian rhythmicity (Súarez-López et al., 2001). Presumably *CO* lies distal to the circadian oscillator in the photoperiodic pathway. Nonetheless, flowering timing mutants constitute a reservoir of potential circadian clock mutants. For example, null alleles of *FLOWERING LOCUS C*, in the autonomous flowering pathway, confer early flowering and shorten the circadian period in leaf movement (Swarup et al., 1999).

Although *CO* may not be a component of the circadian oscillator, it plays a key role in the integration of circadian timekeeping and daylength perception (Súarez-López et al., 2001). *CO* mRNA abundance oscillates and the phase of the peak in *CO* mRNA varies with photoperiod. Only in long days does *CO* mRNA abundance peak in the light. *CO* encodes a Zn finger transcription factor (Putterill et al., 1995), and post-transcriptional regulation of *CO* activity by light is proposed to be critical to the transcriptional activation of *FLOWERING LOCUS T (FT)*, a *CO* target gene (Súarez-López et al., 2001). This provides a molecular explanation of the external coincidence model by which long days promote flowering (Bünning, 1936).

Entrainment (Input)

Circadian rhythms persist in the absence of external time cues but are entrainable to the environment. This entrainment is essential to allow the organism to use its circadian clock to anticipate environmental changes associated with dawn and dusk. The changes in light and temperature encountered at dawn and dusk serve as the chief environmental time cues used by most organisms in clock entrainment.

Light

Circadian clocks, without exception, respond to light (Roenneberg and Foster, 1997) and light is the most potent and best-characterized entraining stimulus in plants

Table 2. Putative components of the Arabidopsis clock. Perturbations of clock function are often accompanied by alterations in flowering timing and by altered inhibition of hypocotyl elongation.

Locus	GenBank (MIPS)	Expression Change	Clock Phenotype ^b	Clock Role	Flowering Time ^c	Hypocotyl Length ^d	Reference ^a
<i>TOC1/APRR1</i>	AF272039 (At5g61380)	loss of function overexpression	short nd ^e	oscillator	early (SD)	WT	(Somers et al., 1998)
<i>CCA1</i>	U79156 (At2g46830)	loss of function overexpression	short arrh	oscillator	late	long	(Green and Tobin, 1999) (Wang and Tobin, 1998)
<i>LHY</i>	AJ006404 (At1g01060)	loss of function overexpression	short arrh	oscillator?	nd late	nd long	(Song and Carré, 2001) (Schaffer et al., 1998)
<i>CKB3</i>	AF068318 (At3g60250)	loss of function overexpression	nd short	?	early	WT	(Sugano et al., 1999)
<i>ZTL</i>	AF254413 (At5g57360)	loss of function overexpression	long/arrh (R) nd	input?	late (LD) late	short long	(Somers et al., 2000) ^f (Nelson et al., 2000)
<i>FKF</i>	AF216523 (At1g68050)	loss of function overexpression	waveform nd	input?	late (LD)	short	(Nelson et al., 2000)
<i>LKP2</i>	AF252295 (At2g18910)	loss of function overexpression	nd arrh	oscillator?	late (LD)	long	(Schultz et al., 2001)
<i>ELF3</i>	AC004747.2 ^g (At2g25930)	loss of function overexpression	arrh (LL) fluence	input	early nd	long short	(Hicks et al., 1996) (Liu et al., 2001)
<i>GI</i>	AF076686 (At1g22770)	loss of function overexpression	period nd	input?	late	long	(Park et al., 1999) ^h
<i>FLC</i>	AF116527 (At5g10140)	loss of function overexpression	short nd	?	early late	nd nd	(Swarup et al., 1999)
<i>PHYB</i>	X17342 (At2g18790)	loss of function phase overexpression	fluence (R) fluence (R)	input	early late	long short	(Devlin and Kay, 2001) (Salomé et al., 2001) (Devlin and Kay, 2001)

^a A representative reference is listed for each locus, although more may be available. For more details, see text.

^b Clock phenotypes include arrhythmic (arrh), short period (short), long period (long), and altered fluence response in period length (fluence). R indicates phenotype seen in red light. LL indicates phenotype seen in continuous light.

^c SD, short days; LD, long days

^d WT, wild type

^e nd = not determined

^f see also (Jarillo et al., 2001)

^g protein_id="AAC31242.1

^h see also (Fowler et al., 1999; Huq et al., 2000)

(Devlin and Kay, 2001). Light perception in plants has been studied and reviewed in detail (see articles in this compilation by Briggs, Deng, and Liscum). The Arabidopsis genome encodes five red/far red-perceiving phytochromes (PHYA-PHYE) and two blue/UV-A-perceiving cryptochromes (CRY1 and CRY2) as well as several other blue light receptors, including the phototropins (PHOT1/NPH1 and PHOT2/NPL1) and zeaxanthin (Casal, 2000; Christie and Briggs, 2001; Lin, 2000b; Nagy and Schäfer, 2000). There is considerable experimental evidence demonstrating the roles of phytochromes and cryptochromes in providing light input to the clock (Devlin and Kay, 2001).

In plants and in light-active animals, period length is inversely related to light intensity; the circadian period shortens as light intensity increases (Aschoff, 1960). This mode of light input is called parametric entrainment, and can be quantified as a fluence response curve (Devlin and Kay, 2000; Somers et al., 1998). Genetic experiments with Arabidopsis mutants have established roles for PHYA, PHYB, PHYD, PHYE, CRY1 and CRY2 in the establishment of period length (Devlin and Kay, 2000; Millar et al., 1995b; Somers et al., 1998). Light-labile PHYA is the predominant photoreceptor for the clock at low intensities of red or blue light, whereas PHYB and CRY1 dominate at high intensities of red and blue light, respectively (Somers et al., 1998). Double mutant studies demonstrate a role for CRY2 in the establishment of period at intermediate intensities of blue light, although that role is redundantly specified by CRY1 (Devlin and Kay, 2000). *cry1 cry2* double mutants retain rhythmicity (Devlin and Kay, 2000), which eliminates the CRY proteins as non-redundantly-specified critical components of the plant circadian oscillator, unlike the case with the mouse clock (van der Horst et al., 1999). Moreover, the quadruple *phyA phyB cry1 cry2* mutant retains both rhythmicity (leaf movement) and the ability to be entrained to a light-dark cycle, making it clear that other photoreceptors (PHYC-PHYE, or others), can provide light input to the clock (Yanovsky et al., 2000b). Roles for PHYD and PHYE in clock input under high intensity red light are supported by period lengthening observed in triple *phyA phyB phyD* and *phyA phyB phyE* mutants versus the *phyA phyB* double mutant (Devlin and Kay, 2000). More triple and quadruple mutant combinations will be required to assess the role of PHYC.

Loss of PHYB or CRY1 (PA Salomé & CR McClung, unpublished data) function does not affect the period length in white light, but instead alters the phase angle of multiple rhythms, indicating that PHYB and CRY1 contribute to the establishment of circadian phase as well as period. PHYB and CRY1 signaling might establish phase through the induction of a critical clock component at dawn. For example, loss of *Neurospora vivid* function alters the phase of the rhythm in *FREQUENCY* (*FRQ*)

expression (Heintzen et al., 2001). In addition, mutants lacking the cyanobacterial phytochrome, CikA, show dramatic alterations in phase angle of multiple gene expression rhythms (Schmitz et al., 2000). CikA is thought to mediate light signaling to the clock proteins KaiA and KaiB (Schmitz et al., 2000).

PHOTOTROPIN 1 (PHOT1/NPH1; DE Somers, unpublished observations) mutants are unaffected in period length; neither PHOT2 (NPL1) nor zeaxanthin have been examined as components of the light input pathways to the clock. However, a novel family of putative photoreceptors, ZEITLUPE (ZTL/LKP1/ADO1) and FLAVIN-BINDING KELCH REPEAT F-BOX (FKF/ADO3) has recently been identified by the mutant phenotype of altered circadian rhythms (Jarillo et al., 2001; Nelson et al., 2000; Somers et al., 2000). A third family member, LOV DOMAIN KELCH PROTEIN 2 (LKP2/ADO2), was identified on the basis of the similarity among the PAS domains (Jarillo et al., 2001; Kiyosue and Wada, 2000). *FKF* mRNA abundance oscillates with an evening-specific maximum, but neither *ZTL* nor *LKP2* mRNAs oscillate (Nelson et al., 2000; Schultz et al., 2001; Somers et al., 2000). Recently, *LKP2* overexpressing plants were shown to be arrhythmic by leaf movement and gene expression in constant conditions, although a rhythm could still be driven by a light-dark cycle (Schultz et al., 2001). *ztl* mutants show long periods in multiple rhythms and the severity of the period lengthening displays fluence rate dependence (Somers et al., 2000), whereas *fkf* mutants exhibit altered waveform in *CCA1* and *LHCB* mRNA oscillations (Nelson et al., 2000). A T-DNA insertion allele of *ztl*, (called *adagio1* [*ado1*]), is arrhythmic in leaf movement in red light (Jarillo et al., 2001). Both *ztl* and *fkf* mutants are late flowering (Nelson et al., 2000; Somers et al., 2000).

These ZTL/FKF/LKP2 proteins possess novel combinations of known functional domains. Family members have a single PAS domain. PAS domains, which function in protein-protein interactions, are found in many clock proteins (Loros and Dunlap, 2001; Reppert and Weaver, 2001; Williams and Sehgal, 2001). The PAS domains of ZTL/FKF/LKP2 are most similar to those of PHOT1 (NPH1), the *Neurospora* White Collar clock proteins, and an unusual phytochrome from a fern (Christie and Briggs, 2001; Loros and Dunlap, 2001; Nozue et al., 1998). The PAS (also called LOV, for light oxygen voltage) domain of PHOT1 (NPH1) binds flavin and may be involved in blue light sensing (Christie et al., 1999), which suggests that ZTL/FKF/LKP2 may also be photoreceptors. ZTL, FKF and LKP2 each also contain Kelch repeats that function in protein-protein interactions (Adams et al., 2000). In addition, each contains an F-box, a domain that recruits target proteins to E3 ubiquitination complexes (Patton et al., 1998). Although a role in ubiquitination has not been functionally established for ZTL, FKF or LKP2, such evidence

has been collected for other plant F-box proteins that function in flower development (UFO; Samach et al., 1999) and auxin responses (TIR; Gray and Estelle, 2000). Proteolytic degradation has been shown to play important roles in other clock systems. For example, *Drosophila* TIM is degraded in the light by an ubiquitin-proteasome mechanism; TIM degradation is correlated with changes in phosphorylation (Naidoo et al., 1999), which is required for substrate recognition by F-box proteins (Craig and Tyers,

1999). As described below, Arabidopsis CCA1 is phosphorylated by casein kinase 2 (CK2) (Sugano et al., 1998; Sugano et al., 1999). Are phosphorylation and degradation of CCA1, LHY or another key target part of the circadian oscillator mechanism (Figure 5)?

There is considerable interaction among photoreceptors. PHYA and CRY1 directly interact at the molecular level, with CRY1 serving as a phosphorylation substrate for PHYA *in vitro* (Ahmad et al., 1998). *In vivo*, CRY1 is phos-




Figure 5. A speculative model of an Arabidopsis circadian clock. Light input via phytochromes and cryptochromes (for simplicity, only PHYB and CRY1 are shown, although others are certainly involved) is mediated through ZTL, ELF3 and GI, or through PIF3. ZTL/ADO1 is known to bind to PHYB and CRY1. PIF3 binds to CCA1 and LHY promoters and possibly to other targets in the clock. The pathway downstream of GI is not known. Although the input pathways are drawn as discrete linear pathways, there may be interaction among them. For simplicity, a single central oscillator is illustrated with a number of putative oscillator components indicated. Components on the circular arrows oscillate in mRNA or protein abundance. One should not infer causal relationships among putative components from the relative order of their placement on the circle as experimental proof is lacking. FKF/LKP2/ZTL are clustered, although there is no evidence that they form molecular complexes. Moreover, *LKP2* mRNA oscillates and overexpression results in arrhythmicity, making LKP2 a strong candidate as an oscillator component. CCA1 and LHY are phosphorylated by CK2, which may make them substrates for the F-box proteins (ZTL, FKF and LKP2) and target them for ubiquitination and degradation by the proteasome (trash can). Output pathways may emanate from any of the putative oscillator components. CCA1, LHY, RVEs and TOC1/APRR1 are DNA-binding proteins, and CCA1 is known to bind to *LHCB* promoters. Other outputs from the oscillator feed back to input components, such as PHYA, PHYB and CRY1, which are regulated by the clock at transcriptional and mRNA abundance levels

phorylated in response to red light in a far-red reversible manner (Ahmad et al., 1998). A *cry1* null mutant shows lengthened period in low intensity red or white light and there is no additivity seen in the double *phyA cry1* mutant (Devlin and Kay, 2000). This suggests that CRY1 acts as a signal transduction component downstream from PHYA in the low intensity light input pathway to the clock (Devlin and Kay, 2001). ZTL has also been shown in the yeast two-hybrid assay to interact physically with the photoreceptors PHYB and CRY1 (Jarillo et al., 2001). However, it is important to recall that the compartmentalization of these photoreceptors and their downstream signaling components is regulated (Nagy et al., 2001), so it is important to confirm the putative interaction *in vivo*. For example, PHYB and CRY2 have been shown to interact *in vivo* by Fluorescence Resonance Energy Transfer (Más et al., 2000).

Light pulses (non-parametric entrainment) can also shift the phase of the clock. Red light pulses phase shift the clock by a very low fluence PHY response (Kolar et al., 1998; Nagy et al., 1993) and far red light pulses phase shift in a PHYA-dependent fashion (Yanovsky et al., 2000a). It has long been clear that clock response to environmental stimuli varies over the circadian cycle. For example, a pulse of light administered prior to dawn advances the phase of the clock whereas the identical pulse applied after dusk delays clock phase. The phase response curve (PRC) is plot of the amplitude of the phase shift resulting from a stimulus applied at discrete phases spanning a circadian cycle (Johnson, 1990). Light phase response curves are available for a number of angiosperms, and recently phase response curves for pulses of red or blue light over a dark background were generated in Arabidopsis (Covington et al., 2001).

One mechanism by which the sensitivity of the oscillator to light might vary over the circadian cycle would be clock regulation of components of the light input pathway; input pathway components may themselves be encoded by CCGs. Microarray experiments indicate that *PHYB*, *CRY1*, *CRY2*, and *PHOT1* (*NPH1*) mRNAs oscillate (Harmer et al., 2000; Schaffer et al. 2001). *PHYB* transcription, as monitored with *PHYB::LUC* gene fusions, is rhythmic in tobacco and Arabidopsis, although bulk PHYB protein abundance does not oscillate (Bognár et al., 1999). *PHYA*, *PHYD*, *PHYE*, *CRY1* and *CRY2* show circadian oscillations both at mRNA abundance and transcriptional levels (Tóth et al., 2001). *PHYC* mRNA oscillates robustly, although transcription of a *PHYC::LUC* fusion is only weakly rhythmic. It seems that the light responsiveness of the *PHYC* promoter obscures any transcriptional rhythm and that the pronounced circadian oscillation in *PHYC* mRNA abundance reflects an important role of the clock in regulating *PHYC* mRNA stability (Tóth et al., 2001). The clear interpretation of these data is that the clock regulates its own

sensitivity to entraining stimuli through regulated expression of photoreceptors.

Signaling pathways downstream from PHY and CRY are under active study. Various signaling intermediates, such as cGMP and Ca^{2+} -calmodulin, as well as phosphorylation, are implicated in light signaling (Briggs and Huala, 1999; Casal, 2000; Christie and Briggs, 2001; Deng and Quail, 1999; Guo et al., 2001; Lin, 2000b). Many signaling components downstream from the photoreceptors have been identified. In particular, red-illuminated PHYB (PfrB) interacts with PIF3, a bHLH protein that binds directly to the G box in a number of phytochrome regulated promoters, including those of *CCA1* and *LATE ELONGATED HYPOCOTYL* (*LHY*) (Martínez-García et al., 2000), two putative oscillator components (see below). This G box (CACGTG) is notably similar to the animal E box (CANNTG) that is targeted by heterodimeric transcription factors of *Drosophila* and mammalian central oscillators (Reppert and Weaver, 2001; Williams and Sehgal, 2001). However, the G box has not been directly implicated in circadian transcription in plants.

Genetic studies have implicated two novel genes, *EARLY FLOWERING 3* (*ELF3*) and *GIGANTEA* (*GI*), in light signaling to the clock. *elf3* loss-of-function alleles yield early flowering, hypocotyl elongation, and conditional arrhythmicity in continuous light (Covington et al., 2001; Hicks et al., 1996; McWatters et al., 2000). *ELF3* is a CCG encoding a nuclear protein; both transcript and protein accumulation in the nucleus peak at dusk (Covington et al., 2001; Hicks et al., 2001; Liu et al., 2001). Genetic experiments suggest substantial redundancy in *ELF3* and *PHYB* function (Reed et al., 2000). *ELF3* interacts with *PHYB* and seems to act as a negative modulator of *PHYB* signaling to the clock, as *ELF3* overexpression both lengthens the circadian period and attenuates the resetting effects of red light pulses whereas loss of *ELF3* function renders the plant hypersensitive to light signals (Covington et al., 2001; Liu et al., 2001; McWatters et al., 2000).

GI is another CCG implicated in light input. *gi* mutants are altered in leaf movement and gene expression rhythms of multiple CCGs, including *GI* itself (Fowler et al., 1999; Park et al., 1999). The period shortening effect of *gi-1* on gene expression rhythms is less severe in extended dark than in continuous light and the extension of period length seen in light of decreasing fluence is less pronounced in *gi-1* than in wild type, which indicates that *GI* acts in light input (Park et al., 1999). *gi* exhibits a defect in inhibition of hypocotyl elongation in red but not in far red light, which implicates *GI* in *PHYB* signaling (Huq et al., 2000). *GI* is nuclear, which is consistent with a role in early *PHYB* signaling (Huq et al., 2000). However, *gi* phenotypes are complicated. In the null *gi-2* allele, the period of leaf movement is shortened but the period of gene expression rhythms gradually lengthens (Park et al., 1999). In addi-

tion, the effects of loss of GI function on hypocotyl elongation are the same as seen in *phyB* loss of function, which suggests that GI is a positive mediator of PHYB signaling. However, *gi* mutants are late flowering, which is opposite to the early flowering phenotype of *phyB* null alleles. Liu et al. (Liu et al., 2001) showed that the effects of ELF3 on flowering time are independent of PHYB. Perhaps GI and ELF3 mediate PHYB signaling to the clock but that their effects on flowering time are mediated through another signaling pathway (Huq et al., 2000; Liu et al., 2001). Alternatively, GI and ELF3 may play different roles at different developmental stages.

Earlier we discussed circadian regulation of photoreceptor expression as a mechanism to regulate sensitivity of the plant to light stimuli. There are other potential mechanisms to regulate sensitivity to light stimuli. For example,




Figure 6. Entrainment of the circadian rhythm in leaf (cotyledon) movement by temperature cycles. Upper panel shows individual traces (blue) of the relative cotyledon positions for the two cotyledons of a representative seedling entrained under continuous light to a temperature cycle of 12 hr at 22°C and 12 hr at 18°C and then released into continuous conditions of constant light at 22°C at T=0. The lower panel shows traces (red) for the two cotyledons of a seedling entrained to the antiphase (180° out-of-phase) temperature cycle of 12 hr at 18°C and 12 hr at 22°C. Gray bars indicate subjective night as defined by the cold period of the entraining cycle.

ELF3 is circadian regulated and serves as a negative regulator of PHYB signaling (Covington et al., 2001; Hicks et al., 1996; Liu et al., 2001; McWatters et al., 2000). *SUPPRESSOR OF PHYA* (*SPA1*) is a negative regulator of PHYA signaling, and microarray analysis indicates that *SPA1* mRNA is clock regulated (Harmer et al., 2000). Recent studies may have identified a circadian-regulated enhancer of light signaling. When plants are transferred into the dark, the circadian oscillations in light-inducible genes, such as *LHCB*, rapidly damp to low levels. This has been attributed to the depletion of the positive regulator, Pfr (Kay and Millar, 1993). A tobacco gene, *ZGT* (abbreviated from the Chinese phrase *zhong guang tiaokong*, meaning clock and light controlled) has been isolated (Xu and Johnson, 2001). As the name indicates, *ZGT* is light inducible and is rhythmically expressed with an mRNA abundance peak at mid-day. Although the Arabidopsis *ZGT* homolog has not been directly studied, overexpression of tobacco *ZGT* in Arabidopsis allows the oscillations in *LHCB::LUC* expression to persist in extended dark. In addition, *ZGT* overexpression renders the plant more sensitive to the phase-shifting effects of light pulses. Thus, it seems that *ZGT* enhances the sensitivity of the plant to light in a circadian pattern (Xu and Johnson, 2001), making it a positively acting counterpart to ELF3 and *SPA1*.

Temperature and Imbibition

The circadian oscillator is temperature compensated, which means that the pace of the oscillator is relatively constant over a range of temperatures. However, temperature pulses or temperature steps are potent entraining stimuli. Little is known of the mechanisms of temperature sensing by plants and even less is known about the specific pathways that convey temperature information to the clock. However, temperature pulse PRCs have been generated for several plants (Johnson, 1990). Temperature cycles entrain Arabidopsis rhythms in *LHCB* (Somers et al., 1998) and *CAT3* transcription (TP Michael & CR McClung, unpublished data). Figure 6 illustrates an experiment in which we used out-of-phase temperature cycles of 12 hr at 22°C and 12 hr at 18°C to entrain seedlings 180° out of phase, as measured by leaf movement. The ability of temperature to entrain the clock led us to hypothesize that the temperature step associated with release from stratification at 4°C to growth at 22°C would either initiate clock function or synchronize the clocks among a population of seedlings. Surprisingly, release from stratification was ineffective in resetting the phase of the rhythm in *CAT2* induction by light (Zhong et al., 1998), suggesting

a refractory period before temperature is capable of entraining the Arabidopsis oscillator. Similarly, a light-insensitive circadian oscillator is detected shortly after germination of tobacco and Arabidopsis (Kolar et al., 1995; Kolar et al., 1998). Although germinating seedlings are refractory to temperature and light input, the timing of imbibition (hydration) of the dried seed entrains the clocks within populations of seedlings (Zhong et al., 1998). This makes it clear that circadian clock is running from the time of imbibition.

THE OSCILLATOR: INTERLOCKED NEGATIVE AND POSITIVE FEEDBACK LOOPS

Genetic and molecular biological studies have identified a number of the components of circadian systems in cyanobacteria, Neurospora, Drosophila, mice and humans (Johnson, 2001; Loros and Dunlap, 2001; Reppert and Weaver, 2001; Williams and Sehgal, 2001). There is a great deal of conservation among the components of the fly and mammalian clocks but the PAS domain, a protein-protein interaction domain (Taylor and Zhulin, 1999), is the only element that has been found in all clock systems. With the determination of the complete sequence of the Arabidopsis genome (The Arabidopsis Genome Initiative, 2000), no obvious orthologs to most known clock proteins can be found, demonstrating that at least part of the Arabidopsis clock mechanism is novel. Nonetheless, studies in the other model systems have indicated that circadian oscillators are comprised of two interconnected feedback loops, one positive and one negative (Figure 7; Williams and Dunlap, 2001; Reppert and Weaver, 2001; Williams and Sehgal, 2001).

A number of loci have been implicated in the Arabidopsis circadian system and many putative clock genes have now been cloned (Table 2). Current efforts are directed towards testing the relevance of the interconnected feedback loop model and identifying the components of the Arabidopsis oscillator. Mutations in a central oscillator component should affect clock properties, such as period length or temperature compensation. The activity, although not necessarily the simple abundance, of an oscillator component should oscillate and interference with this oscillation, commonly through overexpression, should abolish rhythmicity. Loss of function of key clock components should abolish rhythmicity, but functional redundancy often complicates the interpretation of such experiments. In addition, it can be difficult to distinguish between components of input pathways versus of the central oscillator itself (Morrow et al., 1999; Morrow et al.,

2001). At present, a pseudo response regulator, TOC1/APRR1, and two myb transcription factors, CCA1 and LHY, are strong candidates as clock components.

A genetic screen based on alterations in rhythmic transcription of a *LUC* transgene driven by regulatory elements of *LHCB1*1* (*CAB2*) identified a series of *timing of CAB* (*toc*) mutations that disrupt clock function (Millar et al., 1995a). Mutations in *toc1* shorten the period of multiple rhythms, including transcription of a number of genes at distinct circadian phases (Krebs and Simon, 1997; Millar et al., 1995a), leaf movement, and stomatal conductance (Somers et al., 1998). The period-shortening effect of *toc1-1* is independent of light intensity and is seen in extended darkness, which argues that TOC1 does not act in light input. *TOC1* mRNA abundance oscillates in continuous light, peaking late in the day. TOC1 feeds back to control its own oscillation because the period of this oscillation is shortened in *toc1-1* seedlings (Strayer et al., 2000). Collectively, these data suggest that TOC1 is likely to be a component of an oscillator. However, *TOC1* mRNA oscillations damp rapidly in extended darkness (Strayer et al., 2000), yet the clock, as monitored by gene expression




Figure 7. A simple model of a circadian oscillator to illustrate interlocked positive and negative feedback loops. For detailed models of Neurospora, Drosophila and murine clocks, see (Heintzen et al., 2001; Reppert and Weaver, 2001; Williams and Sehgal, 2001).

(Kreps and Simon, 1997; Strayer et al., 2000; Zhong et al., 1997), persists robustly in the dark.

TOC1 encodes a pseudo-response regulator (Strayer et al., 2000) and was independently identified as *ARABIDOPSIS PSEUDO-RESPONSE-REGULATOR 1 (APRR1)* (Makino et al., 2000). *TOC1/APRR1* is the founding member of a 6-gene *APRR* family. mRNA abundance of 5 *APRR* family members (but not *APRR2*) oscillates with circadian period (Matsushika et al., 2000; Strayer et al., 2000). *APRR2* further differs from the other 5 *APRRs* in that it includes a single Myb domain related to that of *CCA1* instead of the *CONSTANS*-like domain found in the other 5 *APRRs* (Makino et al., 2000). Each of the 5 cycling *APRRs* shows a peak in mRNA abundance at a distinct time of the day, which suggests that the expression of each *APRR* in turn regulates the accumulation of the subsequent family members (Makino et al., 2001; Matsushika et al., 2000). However, this remains to be established experimentally. We eagerly await the analysis of loss-of-function alleles and of overexpression of the members of this family.

The identification of *TOC1/APRR1* implicates signal transduction through two-component systems (Sakakibara et al., 2000) in clock function. In typical bacterial two-component systems, a sensor histidine kinase responds to an environmental stimulus, autophosphorylates and transfers the phosphate to a response regulator which then effects a response. However, *TOC1* and the 5 other *APRRs* each lack the invariant phosphor-accepting Asp residue, as well as a second of three invariant residues, and are unlikely to function in a conventional His-Asp relay (Makino et al., 2000; Strayer et al., 2000). Might *TOC1* or another of the *APRRs* represent targets of atypical sensor kinases, perhaps *PHY*? Nonetheless, this suggests a mechanistic link to cyanobacterial clocks (Iwasaki and Kondo, 2000; Johnson, 2001), in which the sensory histidine kinase, *SasA*, interacts with the oscillator component *KaiC* (Iwasaki et al., 2000).

TOC1 also has a carboxy-terminal motif seen in the *CONSTANS* family of transcriptional activators (Putterill et al., 1995) as well as an acidic region often found in transcriptional activators. *TOC1/APRR1* interacts with a PIF3-like protein (Makino et al., 2000), which suggests a mechanism by which the clock might regulate acute induction by light or gate its sensitivity to light input (Millar and Kay, 1996; Zhong et al., 1998). Moreover, *TOC1/APRR1* was also identified as an *ABSCISIC ACID INSENSITIVE 3*-interacting protein (Kurup et al., 2000), which might indicate an interaction of the clock with abscisic acid (*ABA*) as an input or provide clock regulation of *ABA* responses.

Two *CONSTANS-LIKE* genes, *COL1* and *COL2*, are each clock-regulated and overexpression of *COL1* shortens the periods of leaf movement and of *LHCB::LUC* transcription (Ledger et al., 2001). However, these genes are

more likely to encode transcription factors necessary for clock output than to represent true oscillator components, as the plants retain rhythmicity. Two clock-regulated glycine-rich RNA binding protein genes, *AtGRP7 (CCR2)*, and the closely related *AtGRP8 (CCR1)*, exhibit evening-peaking oscillations in transcription and mRNA abundance (Heintzen et al., 1997). *AtGRP7* mRNA abundance declines as protein accumulates, suggesting negative autoregulation which was confirmed by overexpression in transgenic Arabidopsis. Although *AtGRP7* overexpression depresses endogenous *AtGRP7* expression and blocks the oscillations in mRNA abundance of *AtGRP7* and *AtGRP8*, it does not affect circadian oscillations in *LHCB* and *CAT* mRNAs. This suggests that *AtGRP7*, and presumably *AtGRP8*, are key components of a slave (non-self-sustaining) oscillator but not of a central oscillator (Heintzen et al., 1997).

LHY and *CCA1* are two single Myb domain transcription factors implicated in the circadian oscillator mechanism (Schaffer et al., 1998; Wang and Tobin, 1998). *CCA1* and *LHY* are both *CCGs* and both negatively autoregulate, consistent with roles as clock components (Covington et al., 2001; Schaffer et al., 1998; Wang and Tobin, 1998). Overexpression of either *LHY* or *CCA1* results in arrhythmicity of multiple clock outputs, including mRNA abundance of all clock-regulated genes tested to date (Fowler et al., 1999; Schaffer et al., 1998; Wang and Tobin, 1998), leaf movement (Strayer and Kay, 1999), and CO_2 assimilation (CR McClung & EM Tobin, unpublished observations). Loss of *CCA1* function in a T-DNA disruptant line shortens the period of mRNA oscillation in multiple clock-controlled genes, but the plants retain rhythmicity (Green and Tobin, 1999). Recently it has been reported that loss of *LHY* function results in a similar short period phenotype (Song and Carré, 2001). Possibly *CCA1* and *LHY* play redundant roles in the oscillator. Excitingly, the double *cca1 lhy* mutant is arrhythmic in leaf movement, although rapidly damping gene expression rhythms with abnormal phase could be detected with *LUC* activity (Song and Carré, 2001). This indicates that *CCA1* and *LHY* are necessary for sustained oscillator function.

The model of the oscillator as a feedback loop, or as interconnected loops (Figure 7; Loros and Dunlap, 2001; Reppert and Weaver, 2001; Williams and Sehgal, 2001) makes testable predictions, most obviously reciprocal regulation of negative feedback loop components. Recently, Alabadí and collaborators (Alabadí et al., 2001) have shown that *TOC1/APRR1* and *CCA1/LHY* comprise a feedback loop in which *LHY* and *CCA1* are negative regulators of *TOC1* and in which *TOC1* acts as a positive regulator of *CCA1* and *LHY*. Overexpression of either *LHY* or *CCA1* results in non-oscillating low-level accumulation of *TOC1* mRNA. Consistent with their roles as negative regulators of *TOC1* expression, *CCA1/LHY* binding sites have

been identified in the *TOC1* promoter. Conversely, in plants carrying a strong loss-of-function allele of *TOC1* (*toc1-2*), oscillations of *LHY* and *CCA1* mRNA exhibit both the short period characteristic of *toc1* mutations (Millar et al., 1995a; Somers et al., 1998) and greatly reduced mRNA abundance, consistent with a role of *TOC1* as a positive regulator. However, *TOC1* has not yet been shown to bind to either *CCA1* or *LHY* promoters, so it remains to be established that this positive regulation is direct. Nonetheless, the clear conclusion to these data is that *CCA1*, *LHY* and *TOC1/APRR1* participate in a negative feedback loop and represent true components of an Arabidopsis clock.

Additional members of the *CCA1/LHY* family, termed *REVEILLE* (*RVE*), have been identified (CR Andersson & SA Kay, personal communication). Like *CCA1* and *LHY*, at least some *RVEs* (CR Andersson & SA Kay, personal communication) are CCGs and oscillate at both mRNA and protein levels. *CCA1* binds in circadian fashion to a short element of the *LHCB1*3* (*CAB1*) promoter sufficient to confer phytochrome responsiveness and circadian transcription (Wang and Tobin, 1998). Thus *CCA1/LHY/RVE* may represent components of the central oscillator as well as components of the output pathway by which the clock regulates transcription (Schaffer et al., 1998; Wang and Tobin, 1998).

Phosphorylation by casein kinase I is critical in *Drosophila* and mammalian clocks (Lowrey et al., 2000; Reppert and Weaver, 2001; Williams and Sehgal, 2001). Phosphorylation of the FREQUENCY clock component is essential for *Neurospora* clock function (Loros and Dunlap, 2001) and autophosphorylation of KaiC is essential for cyanobacterial rhythmicity (Iwasaki and Kondo, 2000). Consistent with a critical role of phosphorylation in the Arabidopsis clock, *CCA1* binding to DNA is affected by phosphorylation by casein kinase II (CK2) (Sugano et al., 1998), which also phosphorylates *LHY* *in vitro* (Sugano et al., 1999). Overexpression of the regulatory CKB3 subunit increases CK2 activity, which would be presumed to enhance *CCA1* activity. However, CKB3 overexpression results in period shortening, similar to that seen in plants with reduced *CCA1* activity (Sugano et al., 1999). This apparent inconsistency probably indicates our incomplete understanding of the role of *CCA1/LHY/RVE* proteins in the circadian system.

Other genetic approaches have also identified loci with potential roles in the circadian system. There is considerable natural variation in period length, and Quantitative Trait analysis of lines derived from a cross of Cvi to Ler has identified a number of oscillator candidates in Arabidopsis (Swarup et al., 1999). Null mutations of *FLOWERING LOCUS C*, in the autonomous flowering pathway, confer early flowering and shorten the circadian period in leaf movement (Swarup et al., 1999). It is unlikely that we have

as yet identified all the potential components of the Arabidopsis oscillator, and further genetic analyses are likely to identify new candidates as well as new and informative alleles of current candidates.

ARE THERE MULTIPLE CIRCADIAN OSCILLATORS?

The unicellular dinoflagellate, *Gonyaulax polyedra* expresses two distinct circadian oscillators. First, it was established in long time courses that two distinct rhythms with different periods actually showed phase crossings (Roenneberg and Morse, 1993) and second, they showed that the two rhythms could be independently reset by a single stimulus (Morse et al., 1994). This has prompted efforts to demonstrate two or more clocks in multicellular organisms. However, one must be careful to distinguish between the existence of distinct clocks in different organs, tissues, or cell types from the existence of two molecular clocks within a single cell. There are a number of observations of multiple rhythms running with different periods within a plant (internal desynchronization). One recent example is in *Phaseolus vulgaris*, where rhythms in CO₂ assimilation and stomatal aperture exhibit a different period from the rhythm in leaf movement (Hennessey and Field, 1992). In Arabidopsis, the free-running periods in leaf movement and *LHCB* (*CAB*) expression are different, although both are shortened by the *toc1-1* mutation (Millar et al., 1995a). However, this type of data can readily be explained by slight differences in clock function in different tissues, as the leaf movement rhythm is expressed in cells of the petiole whereas the *LHCB* transcription rhythm is measured in leaf mesophyll cells. Explants of different organs retain rhythmicity in *LHCB*, *CHS*, and *PHYB* transcription in culture, which confirms that there are multiple self-sustaining and entrainable circadian oscillators (Thain et al., 2000). Do these clocks communicate to mutually entrain one another? Apparently not. The two cotyledons of intact Arabidopsis and tobacco seedlings can be entrained antiphase to one another. In tobacco, the cotyledons could be entrained to phases distinct from the initial phase retained by the shoot apex. Similar results were obtained with two primary tobacco leaves or with roots versus aerial tissues of Arabidopsis. Collectively, these experiments argue compellingly that there are autonomous clocks in different organs and tissues. Moreover, distal and proximal areas of a single primary tobacco leaf could be entrained to distinct phases (Thain et al., 2000). Thus, there are autonomous cellular clocks and these clocks can exhibit heterogeneity at the level of a single organ.

More suggestive of two distinct clocks within a single cell is the observation that, in extended darkness, the period of *LHCB* transcription lengthens to ~30 h whereas the oscillations in *AtGRP7/CCR2* and *AtGRP8/CCR1* (Carpenter et al., 1994; Strayer et al., 2000), and *CAT3* (Zhong et al., 1997; TP Michael and CR McClung, unpublished data) mRNA abundance and transcription retain 24 hr periods. In tobacco seedlings, cytosolic Ca^{2+} and *LHCB* transcription oscillate with different periods (Sai and Johnson, 1999). In each case, it remains to be established that the two rhythms are expressed in the same cells, but it is nonetheless clear that the rhythms are responding to distinct circadian oscillators (Sai and Johnson, 1999). The issue of one or more clocks at a cellular level represents more than an issue of clock esoterica. If we are to make sense of the complex input pathways, the profusion of putative oscillator components, and the myriad clock outputs described above, we must know if all the pieces fit into a single puzzle.

CONCLUSIONS

The preceding *Arabidopsis* monograph was published only seven years (~2500 circadian cycles) ago. At that time, the chief accomplishment of the *Arabidopsis* rhythms community had been to establish that *Arabidopsis* expressed circadian rhythms. We hope that this review demonstrates the great progress that has been made towards the understanding of the *Arabidopsis* circadian system. Like the rest of the *Arabidopsis* research community, circadian rhythms have embraced (or been embraced by) the genomics age. We have a real appreciation for the enormous influence of the circadian clock on the biology of the plant. There remains a great deal to be accomplished. Although we certainly have candidates for the central oscillator, we most certainly do not have a detailed molecular understanding of the mechanism of the *Arabidopsis* clock. We can be optimistic that this is coming. It is just a matter of time.

ACKNOWLEDGMENTS

Work in our laboratory on circadian rhythms is supported by grants IBN-9817603 and MCB-0091008 from the National Science Foundation to CRM.

REFERENCES

- Adams, J., Kelso, R., and Cooley, L. (2000). The kelch repeat superfamily of proteins: propellers of cell function. *Trends Cell Biol.* **10**, 17-24.
- Ahmad, M., Jarillo, J.A., Smirnova, O., and Cashmore, A.R. (1998). The CRY1 blue light photoreceptor of *Arabidopsis* interacts with phytochrome A in vitro. *Mol. Cell* **1**, 939-948.
- Alabadí, D., Oyama, T., Yanovsky, M.J., Harmon, F.G., Más, P., and Kay, S.A. (2001). Reciprocal regulation between *TOC1* and *LHY/CCA1* within the *Arabidopsis* circadian clock. *Science* **293**, 880-883.
- Aschoff, J. (1960). Exogenous and endogenous components in circadian rhythms. *Cold Spring Harbor Symposia on Quantitative Biology* **XXV**, 11-28.
- Barnes, S.A., McGrath, R.B., and Chua, N.-H. (1997). Light signal transduction in plants. *Trends Cell Biol.* **7**, 21-26.
- Baum, G., Long, J.C., Jenkins, G.I., and Trewavas, A.J. (1999). Stimulation of the blue light phototropic receptor NPH1 causes a transient increase in cytosolic Ca^{2+} . *Proc. Natl. Acad. Sci. U S A* **96**, 13554-13559.
- Beerling, D.J., Osborne, C.P., and Chaloner, W.G. (2001). Evolution of leaf-form in land plants linked to atmospheric CO_2 decline in the Late Palaeozoic era. *Nature* **410**, 352 - 354.
- Bognár, L.K., Hall, A., Ádám, É., Thain, S.C., Nagy, F., and Millar, A.J. (1999). The circadian clock controls the expression pattern of the circadian input photoreceptor, phytochrome B. *Proc. Natl. Acad. Sci. USA* **96**, 14652-14657.
- Borevitz, J.O., Xia, Y., Blount, J., Dixon, R.A., and Lamb, C. (2000). Activation tagging identifies a conserved MYB regulator of phenylpropanoid biosynthesis. *Plant Cell* **12**, 2383-2394.
- Briggs, W.R., and Huala, E. (1999). Blue-light photoreceptors in higher plants. *Annu. Rev. Cell Dev. Biol.* **15**, 33-62.
- Bünning, E. (1936). Die endogene Tagesrhythmik als Grundlage der photoperiodischen Reaktion. *Ber. dtsh. bot. Ges.* **54**, 590-607.
- Bünning, E. (1971). The adaptive value of circadian leaf movements. In *Biochronometry; proceedings of a symposium*, M. Menaker, ed (Washington: National Academy of Sciences), pp. 203-211.
- Carpenter, C.D., Kreps, J.A., and Simon, A.E. (1994). Genes encoding glycine-rich *Arabidopsis thaliana* proteins with RNA-binding motifs are influenced by cold treatment and an endogenous circadian rhythm. *Plant Physiol.* **104**, 1015-1025.
- Carré, I.A., and Kay, S.A. (1995). Multiple DNA-protein complexes at a circadian-regulated promoter element. *Plant Cell* **7**, 2039-2051.
- Casal, J.J. (2000). Phytochromes, cryptochromes, phototropin: photoreceptor interactions in plants. *Photochem. Photobiol.* **71**, 1-11.
- Cheng, C.-L., Acedo, G.N., Dewdney, J., Goodman, H.M., and Conkling, M.A. (1991). Differential expression of the two *Arabidopsis* nitrate reductase genes. *Plant Physiol.* **96**, 275-279.
- Christie, J.M., and Briggs, W.R. (2001). Blue light sensing in higher plants. *J. Biol. Chem.* **276**, 11457-11460.

- Christie, J.M., Salomon, M., Nozue, K., Wada, M., and Briggs, W.R.** (1999). LOV (light, oxygen, or voltage) domains of the blue-light photoreceptor phototropin (nph1): binding sites for the chromophore flavin mononucleotide. *Proc. Natl. Acad. Sci. U S A* **96**, 8779-8783.
- Cohen, A.S., and Cumming, B.G.** (1974). Endogenous rhythmic activity of nitrate reductase in a selection of *Chenopodium rubrum*. *Can. J. Bot.* **52**, 2351-2360.
- Covington, M.F., Panda, S., Liu, X.L., Strayer, C.A., Wagner, D.R., and Kay, S.A.** (2001). ELF3 modulates resetting of the circadian clock in Arabidopsis. *Plant Cell* **13**, 1305-1316.
- Craig, K.L., and Tyers, M.** (1999). The F-box: a new motif for ubiquitin dependent proteolysis in cell cycle regulation and signal transduction. *Prog. Biophys. Mol. Biol.* **72**, 299-328.
- de Mairan, J.** (1729). Observation botanique. *Hist. Acad. Roy. Sci.* 35-36.
- Deng, M.-D., Moureaux, T., Leydecker, M.-T., and Caboche, M.** (1990). Nitrate-reductase expression is under the control of a circadian rhythm and is light inducible in *Nicotiana tabacum* leaves. *Planta* **180**, 257-261.
- Deng, M.D., Moureaux, T., and Lamaze, T.** (1989). Diurnal and circadian fluctuation of malate levels and its close relationship to nitrate reductase activity in tobacco leaves. *Plant Sci.* **65**, 191-197.
- Deng, X.-W., and Quail, P.H.** (1999). Signalling in light-controlled development. *Sem. Cell Develop. Biol.* **10**, 121-129.
- Devlin, P.F., and Kay, S.A.** (2000). Cryptochromes are required for phytochrome signaling to the circadian clock but not for rhythmicity. *Plant Cell* **12**, 2499-2510.
- Devlin, P.F., and Kay, S.A.** (2001). Circadian photoperception. *Annu. Rev. Physiol.* **63**, 677-694.
- Dowson-Day, M.J., and Millar, A.J.** (1999). Circadian dysfunction causes aberrant hypocotyl elongation patterns in Arabidopsis. *Plant J.* **17**, 63-71.
- Engelmann, W., and Johnsson, A.** (1998). Rhythms in organ movement. In *Biological Rhythms and Photoperiodism in Plants*, P.J. Lumsden and A.J. Millar, eds (Oxford: BIOS Scientific Publishers, Ltd.), pp. 35-50.
- Engelmann, W., Simon, K., and Phen, C.Y.** (1992). Leaf movement rhythm in *Arabidopsis thaliana*. *Z. Naturforschung* **47c**, 925-928.
- Eskin, A.** (1979). Identification and physiology of circadian pacemakers. *Fed. Proc.* **38**, 2570-2572.
- Fejes, E., and Nagy, F.** (1998). Molecular analysis of circadian clock-regulated gene expression in plants: features of the 'output' pathways. In *Biological Rhythms and Photoperiodism in Plants*, P.J. Lumsden and A.J. Millar, eds (Oxford: BIOS Scientific Publishers, Ltd.), pp. 99-118.
- Finlayson, S.A., Lee, I.-J., and Morgan, P.W.** (1998). Phytochrome B and the regulation of circadian ethylene production in sorghum. *Plant Physiol.* **116**, 17-25.
- Finlayson, S.A., Lee, I.-J., Mullet, J.E., and Morgan, P.W.** (1999). The mechanism of rhythmic ethylene production in sorghum. The role of phytochrome B and simulated shading. *Plant Physiol.* **119**, 1083-1089.
- Foster, K.R., and Morgan, P.W.** (1995). Genetic regulation of development in *Sorghum bicolor*. IX. The *mag³* allele disrupts diurnal control of gibberellin biosynthesis. *Plant Physiol.* **108**, 337-343.
- Fowler, S., Lee, K., Onouchi, H., Samach, A., Richardson, K., Morris, B., Coupland, G., and Putterill, J.** (1999). *GIGANTEA*: a circadian clock-controlled gene that regulates photoperiodic flowering in *Arabidopsis* and encodes a protein with several membrane-spanning domains. *EMBO J.* **18**, 4679-4688.
- Frohnmeier, H., Bowler, C., Zhu, J.-K., Yamagata, H., Schäfer, E., and Chua, N.-H.** (1998). Different roles for calcium and calmodulin in phytochrome and UV-regulated expression of chalcone synthase. *Plant J.* **13**, 763-772.
- Golden, S.S., and Strayer, C.** (2001). Time for plants. *Progress in plant chronobiology. Plant Physiol.* **125**, 98-101.
- Gómez, L.A., and Simón, E.** (1995). Circadian rhythm of *Robinia pseudoacacia* leaflet movements: role of calcium and phytochrome. *Photochem. Photobiol.* **61**, 210-215.
- Gray, W.M., and Estelle, M.** (2000). Function of the ubiquitin-proteasome pathway in auxin response. *Trends Biochem. Sci.* **25**, 133-138.
- Green, R.M., and Tobin, E.M.** (1999). Loss of the circadian clock-associated protein 1 in *Arabidopsis* results in altered clock-regulated gene expression. *Proc. Natl. Acad. Sci. USA* **96**, 4176-4179.
- Guo, H., Mockler, T., Duong, H., and Lin, C.** (2001). SUB1, an Arabidopsis Ca^{2+} -binding protein involved in cryptochrome and phytochrome coaction. *Science* **291**, 487-490.
- Harmer, S.L., Hogenesch, J.B., Straume, M., Chang, H.-S., Han, B., Zhu, T., Wang, X., Kreps, J.A., and Kay, S.A.** (2000). Orchestrated transcription of key pathways in Arabidopsis by the circadian clock. *Science* **290**, 2110-2113.
- Hartwell, J., Gill, A., Nimmo, G.A., Wilkins, M.B., Jenkins, G.I., and Nimmo, H.G.** (1999). Phosphoenolpyruvate carboxylase kinase is a novel protein kinase regulated at the level of expression. *Plant J.* **20**, 333-342.
- Hartwell, J., Smith, L.H., Wilkins, M.B., Jenkins, G.I., and Nimmo, H.G.** (1996). Higher plant phosphoenolpyruvate carboxylase kinase is regulated at the level of translatable mRNA in response to light or a circadian rhythm. *Plant J.* **10**, 1071-1078.
- Heintzen, C., Loros, J.J., and Dunlap, J.C.** (2001). The PAS protein VIVID defines a clock-associated feedback loop that represses light input, modulates gating, and regulates clock resetting. *Cell* **104**, 453-464.
- Heintzen, C., Nater, M., Apel, K., and Staiger, D.** (1997). AtGRP7, a nuclear RNA-binding protein as a component of a circadian-regulated negative feedback loop in *Arabidopsis thaliana*. *Proc. Natl. Acad. Sci. USA* **94**, 8515-8520.
- Hennessey, T.L., and Field, C.B.** (1991). Oscillations in carbon assimilation and stomatal conductance under constant conditions. *Plant Physiol.* **96**, 831-836.
- Hennessey, T.L., and Field, C.B.** (1992). Evidence of multiple circadian oscillators in bean plants. *J. Biol. Rhythms* **7**, 105-113.
- Hicks, K.A., Albertson, T.M., and Wagner, D.R.** (2001). *EARLY FLOWERING3* encodes a novel protein that regulates circadian clock function and flowering in Arabidopsis. *Plant Cell* **13**, 1281-1292.
- Hicks, K.A., Millar, A.J., Carré, I.A., Somers, D.E., Straume, M., Meeks-Wagner, D.R., and Kay, S.A.** (1996). Conditional circadian dysfunction of the *Arabidopsis early-flowering 3* mutant. *Science* **274**, 790-792.

- Huq, E., Tepperman, J.M., and Quail, P.H. (2000). GIGANTEA is a nuclear protein involved in phytochrome signaling in *Arabidopsis*. *Proc. Natl. Acad. Sci. USA* **97**, 9654-9658.
- Hwang, S., Kawazoe, R., and Herrin, D.L. (1996). Transcription of *tufA* and other chloroplast-encoded genes is controlled by a circadian clock in *Chlamydomonas*. *Proc. Natl. Acad. Sci. USA* **93**, 996-1000.
- Levinsh, G., and Kreicbergs, O. (1992). Endogenous rhythmicity of ethylene production in growing intact cereal seedlings. *Plant Physiol.* **100**, 1389-1391.
- Iwasaki, H., and Kondo, T. (2000). The current state and problems of circadian clock studies in cyanobacteria. *Plant Cell Physiol.* **41**, 1013-1020.
- Iwasaki, H., Williams, S.B., Kitayama, Y., Ishiura, M., Golden, S.S., and Kondo, T. (2000). A KaiC-interacting sensory histidine kinase, SasA, necessary to sustain robust circadian oscillation in cyanobacteria. *Cell* **101**, 223-233.
- Jarillo, J.A., Capel, J., Tang, R.-H., Yang, H.-Q., Alonso, J.M., Ecker, J.R., and Cashmore, A.R. (2001). An *Arabidopsis* circadian clock component interacts with both CRY1 and phyB. *Nature* **410**, 487-490.
- Johnson, C.H. (1990). An atlas of phase response curves for circadian and circatidal rhythms. (Nashville: Department of Biology, Vanderbilt University).
- Johnson, C.H. (2001). Endogenous timekeepers in photosynthetic organisms. *Annu. Rev. Physiol.* **63**, 695-728.
- Johnson, C.H., Knight, M., Trewavas, A., and Kondo, T. (1998). A clockwork green: circadian programs in photosynthetic organisms. In *Biological Rhythms and Photoperiodism in Plants*, P.J. Lumsden and A.J. Millar, eds (Oxford: BIOS Scientific Publishers, Ltd.), pp. 1-34.
- Johnson, C.H., Knight, M.R., Kondo, T., Masson, P., Sedbrook, J., Haley, A., and Trewavas, A. (1995). Circadian oscillations of cytosolic and chloroplastic free calcium in plants. *Science* **269**, 1863-1865.
- Jouve, L., Gaspar, T., Kevers, C., Greppin, H., and Agosti, R.D. (1999). Involvement of indole-3-acetic acid in the circadian growth of the first internode of *Arabidopsis*. *Planta* **209**, 136-142.
- Jouve, L., Greppin, H., and Agosti, R.D. (1998). *Arabidopsis thaliana* floral stem elongation: evidence for an endogenous circadian rhythm. *Plant Physiol. Biochem.* **36**, 469-472.
- Kanamaru, K., Fujiwara, M., Seki, M., Katagiri, T., Nakamura, M., Mochizuki, N., Nagatani, A., Shinozaki, K., Tanaka, K., and Takahashi, H. (1999). Plastidic RNA polymerase sigma factors in *Arabidopsis*. *Plant Cell Physiol.* **40**, 832-842.
- Kay, S.A., and Millar, A.J. (1993). Circadian-regulated *cab* gene transcription in higher plants. In *The Molecular Genetics of Biological Rhythms*, M.W. Young, ed (New York: Marcel Dekker), pp. 73-89.
- Kiyosue, T., and Wada, M. (2000). LKP1 (LOV kelch protein 1): a factor involved in the regulation of flowering time in *Arabidopsis*. *Plant J.* **23**, 807-815.
- Kloppstech, K. (1985). Diurnal and circadian rhythmicity in the expression of light-induced nuclear messenger RNAs. *Planta* **165**, 502-506.
- Kolar, C., Ádám, É., Schäfer, E., and Nagy, F. (1995). Expression of tobacco genes for light-harvesting chlorophyll a/b binding proteins of photosystem II is controlled by two circadian oscillators in a developmentally regulated fashion. *Proc. Natl. Acad. Sci. USA* **92**, 2174-2178.
- Kolar, C., Fejes, E., Ádám, É., Schäfer, E., Kay, S., and Nagy, F. (1998). Transcription of *Arabidopsis* and wheat *Cab* genes in single tobacco transgenic seedlings exhibits independent rhythms in a developmentally regulated fashion. *Plant J.* **13**, 563-569.
- Kreps, J.A., and Kay, S.A. (1997). Coordination of plant metabolism and development by the circadian clock. *Plant Cell* **9**, 1235-1244.
- Kreps, J.A., Muramatsu, T., Furuya, M., and Kay, S.A. (2000). Fluorescent differential display identifies circadian clock-regulated genes in *Arabidopsis thaliana*. *J. Biol. Rhythms* **15**, 208-217.
- Kreps, J.A., and Simon, A.E. (1997). Environmental and genetic effects on circadian clock-regulated gene-expression in *Arabidopsis thaliana*. *Plant Cell* **9**, 297-304.
- Kurup, S., Jones, H.D., and Holdsworth, M.J. (2000). Interactions of the developmental regulator ABI3 with proteins identified from developing *Arabidopsis* seeds. *Plant J.* **21**, 143-155.
- Leckie, C.P., McAinsh, M.R., Montgomery, L., Priestley, A.J., Staxen, I., Webb, A.A.R., and Hetherington, A.M. (1998). Second messengers in guard cells. *J. Exp. Bot.* **49**, 339-349.
- Ledger, S., Strayer, C., Ashton, F., Kay, S.A., and Putterill, J. (2001). Analysis of the function of two circadian-regulated *CONSTANS-LIKE* genes. *Plant J.* **26**, 15-22.
- Lin, C. (2000a). Photoreceptors and regulation of flowering time. *Plant Physiol.* **123**, 39-50.
- Lin, C. (2000b). Plant blue-light receptors. *Trends Plant Sci.* **5**, 337-342.
- Liu, X.L., Covington, M.F., Fankhauser, C., Chory, J., and Wagner, D.R. (2001). *ELF3* encodes a circadian clock-regulated nuclear protein that functions in an *Arabidopsis* PHYB signal transduction pathway. *Plant Cell* **13**, 1293-1304.
- Liu, Y., Tsinoremas, N.F., Johnson, C.H., Golden, S.S., Ishiura, M., and Kondo, T. (1995). Circadian orchestration of gene expression in cyanobacteria. *Genes Dev.* **9**, 1469-1478.
- Liu, Z., Taub, C.C., and McClung, C.R. (1996). Identification of an *Arabidopsis* Rubisco Activase (*RCA*) minimal promoter regulated by phytochrome and the circadian clock. *Plant Physiol.* **112**, 43-51.
- Long, J.C., and Jenkins, G.I. (1998). Involvement of plasma membrane redox activity and calcium homeostasis in the UV-B and UV-A/blue light induction of gene expression in *Arabidopsis*. *Plant Cell* **10**, 2077-2086.
- Loros, J.J., and Dunlap, J.C. (2001). Genetic and molecular analysis of circadian rhythms in *Neurospora*. *Annu. Rev. Physiol.* **63**, 757-794.
- Lowrey, P.L., Shimomura, K., Antoch, M.P., Yamazaki, S., Zemenides, P.D., Ralph, M.R., Menaker, M., and Takahashi, J.S. (2000). Positional syntenic cloning and functional characterization of the mammalian circadian mutation *tau*. *Science* **288**, 483-491.
- Lumsden, P.J., and Millar, A.J. (1998). *Biological rhythms and photoperiodism in plants*. (Oxford: Bios Scientific Publishers).

- Makino, S., Kiba, T., Imamura, A., Hanaki, N., Nakamura, A., Suzuki, T., Taniguchi, M., Ueguchi, C., Sugiyama, T., and Mizuno, T.** (2000). Genes encoding pseudo-response regulators: insight into His-to-Asp phosphorelay and circadian rhythm in *Arabidopsis thaliana*. *Plant Cell Physiol.* **41**, 791-803.
- Makino, S., Matsushika, A., Kojima, M., Oda, Y., and Mizuno, T.** (2001). Light response of the circadian waves of the APRR1/TOC1 quintet: when does the quintet start singing rhythmically in *Arabidopsis*? *Plant Cell Physiol.* **42**, 334-339.
- Martínez-García, J.F., Huq, E., and Quail, P.H.** (2000). Direct targeting of light signals to a promoter element-bound transcription factor. *Science* **288**, 859-863.
- Más, P., Devlin, P.F., Panda, S., and Kay, S.A.** (2000). Functional interaction of phytochrome B and cryptochrome 2. *Nature* **408**, 207-211.
- Matsushika, A., Makino, S., Kojima, M., and Mizuno, T.** (2000). Circadian waves of expression of the APRR1/TOC1 family of pseudo-response regulators in *Arabidopsis thaliana*: insight into the plant circadian clock. *Plant Cell Physiol.* **41**, 1002-1012.
- McClung, C.R.** (2000). Plant circadian clocks: a millennial view. *Physiol. Plant.* **109**, 359-371.
- McClung, C.R.** (2001). Circadian rhythms in plants. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **52**, 139-162.
- McClung, C.R., Hsu, M., Painter, J.E., Gagne, J.M., Karlsberg, S.D., and Salomé, P.A.** (2000). Integrated temporal regulation of the photorespiratory pathway: circadian regulation of two *Arabidopsis* genes encoding serine hydroxymethyltransferase. *Plant Physiol.* **123**, 381-392.
- McClung, C.R., and Kay, S.A.** (1994). Circadian rhythms in *Arabidopsis thaliana*. In *Arabidopsis*, E.M. Meyerowitz and C.R. Somerville, eds (Plainview, NY: Cold Spring Harbor Laboratory Press), pp. 615-637.
- McWatters, H.G., Bastow, R.M., Hall, A., and Millar, A.J.** (2000). The *ELF3 zeitnehmer* regulates light signalling to the circadian clock. *Nature* **408**, 716-720.
- Morrow, M., Brunner, M., and Roenneberg, T.** (1999). Assignment of circadian function for the *Neurospora* clock gene *frequency*. *Nature* **399**, 584-586.
- Morrow, M., Franchi, L., Dragovic, Z., Görl, M., Johnson, J., Brunner, M., Macino, G., and Roenneberg, T.** (2001). Circadian regulation of the light input pathway in *Neurospora crassa*. *EMBO J.* **20**, 307-315.
- Millar, A.J.** (1999). Biological clocks in *Arabidopsis thaliana*. *New Phytol.* **141**, 175-197.
- Millar, A.J., Carré, I.A., Strayer, C.A., Chua, N.-H., and Kay, S.A.** (1995a). Circadian clock mutants in *Arabidopsis* identified by luciferase imaging. *Science* **267**, 1161-1163.
- Millar, A.J., and Kay, S.A.** (1996). Integration of circadian and phototransduction pathways in the network controlling *CAB* gene transcription in *Arabidopsis*. *Proc. Natl. Acad. Sci. USA* **93**, 15491-15496.
- Millar, A.J., Short, S.R., Chua, N.-H., and Kay, S.A.** (1992a). A novel circadian phenotype based on firefly luciferase expression in transgenic plants. *Plant Cell* **4**, 1075-1087.
- Millar, A.J., Short, S.R., Hiratsuka, K., Chua, N.-H., and Kay, S.A.** (1992b). Firefly luciferase as a reporter of regulated gene expression in higher plants. *Plant Mol. Biol. Reporter* **10**, 324-337.
- Millar, A.J., Straume, M., Chory, J., Chua, N.-H., and Kay, S.A.** (1995b). The regulation of circadian period by phototransduction pathways in *Arabidopsis*. *Science* **267**, 1163-1166.
- Moore-Ede, M.C., Sulzman, F.M., and Fuller, C.A.** (1982). The clocks that time us: physiology of the circadian timing system. (Cambridge, MA: Harvard University Press).
- Morikawa, K., Ito, S., Tsunoyama, Y., Nakahira, Y., Shiina, T., and Toyoshima, Y.** (1999). Circadian-regulated expression of a nuclear-encoded plastid *s* factor gene (*sigA*) in wheat seedlings. *FEBS Lett.* **451**, 275-278.
- Morse, D., Hastings, J.W., and Roenneberg, T.** (1994). Different phase responses of the two circadian oscillators in *Gonyaulax*. *J. Biol. Rhythms* **9**, 263-274.
- Nagy, F., Fejes, E., Wehmeyer, B., Dallman, G., and Schäfer, E.** (1993). The circadian oscillator is regulated by a very low fluence response of phytochrome in wheat. *Proc. Natl. Acad. Sci. USA* **90**, 6290-6294.
- Nagy, F., Kircher, S., and Schäfer, E.** (2001). Intracellular trafficking of photoreceptors during light-induced signal transduction in plants. *J. Cell Sci.* **114**, 475-480.
- Nagy, F., and Schäfer, E.** (2000). Nuclear and cytosolic events of light-induced, phytochrome-regulated signaling in higher plants. *EMBO J.* **19**, 157-163.
- Naidoo, N., Song, W., Hunter-Ensor, M., and Sehgal, A.** (1999). A role for the proteasome in the light response of the Timeless clock protein. *Science* **285**, 1737-1741.
- Nakahira, Y., Baba, K., Yoneda, A., Shiina, T., and Toyoshima, Y.** (1998). Circadian-regulated transcription of the *psbD* light-responsive promoter in wheat chloroplasts. *Plant Physiol.* **118**, 1079-1088.
- Nelson, D.C., Lasswell, J., Rogg, L.E., Cohen, M.A., and Bartel, B.** (2000). *FKF1*, a clock-controlled gene that regulates the transition to flowering in *Arabidopsis*. *Cell* **101**, 331-340.
- Nimmo, H.G.** (2000). The regulation of phosphoenolpyruvate carboxylase in CAM plants. *Trends Plant Sci.* **5**, 75-80.
- Nozue, K., Kanegae, T., Imaizumi, T., Fukuda, S., Okamoto, H., Yeh, K., Lagarias, J.C., and Wada, M.** (1998). A phytochrome from the fern *Adiantum* with features of the putative photoreceptor NPH1. *Proc. Natl. Acad. Sci. USA* **95**, 15826-15830.
- Park, D.H., Somers, D.E., Kim, Y.S., Choy, Y.H., Lim, H.K., Soh, M.S., Kim, H.J., Kay, S.A., and Nam, H.G.** (1999). Control of circadian rhythms and photoperiodic flowering by the *Arabidopsis* *GIGANTEA* gene. *Science* **285**, 1579-1582.
- Patton, E.E., Willems, A.R., and Tyers, M.** (1998). Combinatorial control in ubiquitin-dependent proteolysis: don't Skp the F-box hypothesis. *Trends Genet.* **14**, 236-243.
- Piechulla, B.** (1999). Circadian expression of the light-harvesting complex protein genes in plants. *Chronobiol. Intl.* **16**, 115-128.
- Pilgrim, M.L., Caspar, T., Quail, P.H., and McClung, C.R.** (1993). Circadian and light regulated expression of nitrate reductase in *Arabidopsis*. *Plant Mol. Biol.* **23**, 349-364.
- Pilgrim, M.L., and McClung, C.R.** (1993). Differential involvement of the circadian clock in the expression of genes required for Ribulose-1,5-bisphosphate carboxylase/oxygenase synthesis, assembly, and activation in *Arabidopsis thaliana*. *Plant Physiol.* **103**, 553-564.

- Pittendrigh, C.S.** (1960). Circadian rhythms and the circadian organization of living systems. Cold Spring Harbor Symposia on Quantitative Biology **XXV**, 159-184.
- Pittendrigh, C.S.** (1981). Circadian rhythms: general perspective. In Handbook of Behavioral Neurobiology. Biological Rhythms., J. Aschoff, ed (New York: Plenum Press), pp. 57-80.
- Plautz, J.D., Kaneko, M., Hall, J.C., and Kay, S.A.** (1997). Independent photoreceptive circadian clocks throughout *Drosophila*. Science **278**, 1632-1635.
- Putterill, J., Robson, F., Lee, K., Simon, R., and Coupland, G.** (1995). The *CONSTANS* gene of *Arabidopsis* promotes flowering and encodes a protein showing similarities to zinc finger transcription factors. Cell **80**, 847-857.
- Reed, J.W., Nagpal, P., Bastow, R.M., Solomon, K.S., Dowson-Day, M.J., Elumalai, R.P., and Millar, A.J.** (2000). Independent action of ELF3 and phyB to control hypocotyl elongation and flowering time. Plant Physiol. **122**, 1149-1160.
- Reppert, S.M., and Weaver, D.R.** (2001). Molecular analysis of mammalian circadian rhythms. Annu. Rev. Physiol. **63**, 647-676.
- Rikin, A.** (1992). Circadian-rhythm of heat-resistance in cotton seedlings - synthesis of heat-shock proteins. Eur. J. Cell Biol. **59**, 160-165.
- Rikin, A., Dillwith, J.W., and Bergman, D.K.** (1993). Correlation between the circadian-rhythm of resistance to extreme temperatures and changes in fatty-acid composition in cotton seedlings. Plant Physiol. **101**, 31-36.
- Rikin, A., St. John, J.B., Wergin, W.P., and Anderson, J.D.** (1984). Rhythmical changes in the sensitivity of cotton seedlings to herbicides. Plant Physiol. **76**, 297-300.
- Roenneberg, T., and Foster, R.G.** (1997). Twilight times: light and the circadian system. Photochem. Photobiol. **66**, 549-561.
- Roenneberg, T., and Morse, D.** (1993). Two circadian oscillators in one cell. Nature **362**, 362-364.
- Sai, J., and Johnson, C.H.** (1999). Different circadian oscillators control Ca(2+) fluxes and *Lhcb* gene expression. Proc. Natl. Acad. Sci. USA **96**, 11659-11663.
- Sakakibara, H., Taniguchi, M., and Sugiyama, T.** (2000). His-Asp phosphorelay signaling: a communication avenue between plants and their environment. Plant Mol. Biol. **42**, 273-278.
- Salvador, M.L., Klein, U., and Bogorad, L.** (1993). Light-regulated and endogenous fluctuations of chloroplast transcript levels in *Chlamydomonas*. Regulation by transcription and RNA degradation. Plant J. **3**, 213-219.
- Samach, A., Klenz, J.E., Kohalmi, S.E., Risseuw, E., Haughn, G.W., and Crosby, W.L.** (1999). The *UNUSUAL FLORAL ORGANS* gene of *Arabidopsis thaliana* is an F-box protein required for normal patterning and growth in the floral meristem. Plant J. **20**, 433-445.
- Sanders, D., Brownlee, C., and Harper, J.F.** (1999). Communicating with calcium. Plant Cell **11**, 691-706.
- Schaffer, R., Landgraf, J., Accerbi, M., Simon, V., Larson, M., and Wisman, E.** (2001). Microarray analysis of diurnal and circadian-regulated genes in *Arabidopsis*. Plant Cell **13**, 113-123.
- Schaffer, R., Ramsay, N., Samach, A., Corden, S., Putterill, J., Carré, I.A., and Coupland, G.** (1998). *LATE ELONGATED HYPOCOTYL*, an *Arabidopsis* gene encoding a MYB transcription factor, regulates circadian rhythmicity and photoperiodic responses. Cell **93**, 1219-1229.
- Schaffer, R., and Wisman, E.** (2001). Microarray analysis using genomic DNA as a common reference. 12th International Conference on Arabidopsis Research, Madison WI, June 23-27, 2001 Abstr. 111.
- Schmitz, O., Katayama, M., Williams, S.B., Kondo, T., and Golden, S.S.** (2000). CikA, a bacteriophytochrome that resets the cyanobacterial circadian clock. Science **289**, 765-768.
- Schroeder, J.I., Allen, G.J., Hugouvieux, V., Kwak, J.M., and Waner, D.** (2001). Guard cell signal transduction. Annu. Rev. Plant Physiol. Plant Mol. Biol. **52**, 627-658.
- Schultz, T.F., Kiyosue, T., Yanovsky, M., Wada, M., and Kay, S.** (2001). A role for LKP2 in the circadian clock of *Arabidopsis*. 12th International Conference on Arabidopsis Research, Madison WI, June 23-27, 2001 Abstr. 238.
- Simpson, G.G., Gendall, A.R., and Dean, C.** (1999). When to switch to flowering. Annu. Rev. Cell Dev. Biol. **15**, 519-550.
- Somers, D.** (1999). The physiology and molecular bases of the plant circadian clock. Plant Physiol. **121**, 9-19.
- Somers, D.E., Schultz, T.F., Milnamow, M., and Kay, S.A.** (2000). ZEITLUPE encodes a novel clock-associated PAS protein from *Arabidopsis*. Cell **101**, 319-329.
- Somers, D.E., Webb, A.A.R., Pearson, M., and Kay, S.A.** (1998). The short-period mutant, *toc1-1*, alters circadian clock regulation of multiple outputs throughout development in *Arabidopsis thaliana*. Development **125**, 485-494.
- Song, H.-R., and Carré, I.** (2001). Function of the *LHY* gene within the *Arabidopsis* circadian clock. 12th International Conference on Arabidopsis Research, Madison WI, June 23-27, 2001 Abstr. 240.
- Staiger, D., and Apel, K.** (1999). Circadian clock-regulated expression of an RNA-binding protein in *Arabidopsis*: characterisation of a minimal promoter element. Mol. Gen. Genet. **261**, 811-819.
- Staiger, D., Apel, K., and Trepp, G.** (1999). The *Atger3* promoter confers circadian clock-regulated transcription with peak expression at the beginning of night. Plant Mol. Biol. **40**, 873-882.
- Strayer, C., Oyama, T., Schultz, T.F., Raman, R., Somers, D.E., Más, P., Panda, S., Kreps, J.A., and Kay, S.A.** (2000). Cloning of the *Arabidopsis* clock gene *TOC1*, an autoregulatory response regulator homolog. Science **289**, 768-771.
- Strayer, C.A., and Kay, S.A.** (1999). The ins and outs of circadian regulated gene expression. Curr. Op. Plant Biol. **2**, 114-120.
- Suárez-López, P., Wheatley, K., Robson, F., Onouchi, H., Valverde, F., and Coupland, G.** (2001). *CONSTANS* mediates between the circadian clock and the control of flowering in *Arabidopsis*. Nature **410**, 1116-1120.
- Sugano, S., Andronis, C., Green, R.M., Wang, Z.-Y., and Tobin, E.M.** (1998). Protein kinase CK2 interacts with and phosphorylates the *Arabidopsis* circadian clock-associated 1 protein. Proc. Natl. Acad. Sci. USA **95**, 11020-11025.

- Sugano, S., Andronis, C., Ong, M.S., Green, R.M., and Tobin, E.M.** (1999). The protein kinase CK2 is involved in regulation of circadian rhythms in Arabidopsis. *Proc. Natl. Acad. Sci. USA* **96**, 12362-12366.
- Swarup, K., Alonso-Blanco, C., Lynn, J.R., Michaels, S.D., Amasino, R.M., Koornneef, M., and Millar, A.J.** (1999). Natural allelic variation identifies new genes in the Arabidopsis circadian system. *Plant J.* **20**, 67-77.
- Sweeney, B.M.** (1987). Rhythmic phenomena in plants. (New York: Academic Press).
- Taylor, B.L., and Zhulin, I.B.** (1999). PAS domains: internal sensors of oxygen, redox potential, and light. *Microbiol. Mol. Biol. Rev.* **63**, 479-506.
- Thain, S.C., Hall, A., and Millar, A.J.** (2000). Functional independence of multiple circadian clocks that regulate plant gene expression. *Curr. Biol.* **10**, 951-956.
- The Arabidopsis Genome Initiative** (2000). Analysis of the genome sequence of the flowering plant *Arabidopsis thaliana*. *Nature* **408**, 796-815.
- Thomashow, M.F., Gilmour, S.J., Stockinger, E.J., Jaglo-Ottosen, K.R., and Zarka, D.G.** (2001). Role of the Arabidopsis CBF transcriptional activators in cold acclimation. *Physiol. Plant.* **112**, 171-175.
- Tosini, G., and Menaker, M.** (1998). Multioscillatory circadian organization in a vertebrate, *Iguana iguana*. *J. Neurosci.* **18**, 1105-1114.
- Tóth, R., Kevei, É., Hall, A., Millar, A.J., Nagy, F., and Kozma-Bognár, L.** (2001). Circadian clock-regulated expression of phytochrome and cryptochrome genes in Arabidopsis. *Plant Physiol.* **127**, in press.
- van der Horst, G.T.J., Muijtjens, M., Kobayashi, K., Takano, R., Kanno, S.-I., Takao, M., De Wit, J., Verkerk, A., Eker, A.P.M., Van Leenen, D., Buijs, R., Bootsma, D., Hoeijmakers, J.H.J., and Yasui, A.** (1999). Mammalian Cry1 and Cry2 are essential for maintenance of circadian rhythms. *Nature* **398**, 627-630.
- Wang, Z.-Y., and Tobin, E.M.** (1998). Constitutive expression of the *CIRCADIAN CLOCK ASSOCIATED 1 (CCA1)* gene disrupts circadian rhythms and suppresses its own expression. *Cell* **93**, 1207-1217.
- Webb, A.A.R.** (1998). Stomatal rhythms. In *Biological Rhythms and Photoperiodism in Plants*, P.J. Lumsden and A.J. Millar, eds (Oxford: BIOS Scientific Publishers, Ltd.), pp. 69-79.
- Williams, J.A., and Sehgal, A.** (2001). Molecular components of the circadian system in *Drosophila*. *Annu. Rev. Physiol.* **63**, 729-755.
- Wood, N.T., Haley, A., Viry-Moussaïd, M., Johnson, C.H., van der Luit, A.H., and Trewavas, A.J.** (2001). The calcium rhythms of different cell types oscillate with different circadian phases. *Plant Physiol.* **125**, 787-796.
- Xu, Y., and Johnson, C.H.** (2001). A clock- and light-regulated gene that links the circadian oscillator to *LHCB* gene expression. *Plant Cell* **13**, 1411-1426.
- Yamazaki, S., Numano, R., Abe, M., Hida, A., Takahashi, R., Ueda, M., Block, G.D., Sakaki, Y., Menaker, M., and Tei, H.** (2000). Resetting central and peripheral circadian oscillators in transgenic rats. *Science* **288**, 682-685.
- Yanovsky, M.J., Izaguirre, M., Wagmaister, J.A., Gatz, C., Jackson, S.D., Thomas, B., and Casal, J.J.** (2000a). Phytochrome A resets the circadian clock and delays tuber formation under long days in potato. *Plant J.* **23**, 223-232.
- Yanovsky, M.J., Mazzella, M.A., and Casal, J.J.** (2000b). A quadruple photoreceptor mutant still keeps track of time. *Curr. Biol.* **10**, 1013-1015.
- Zhong, H.H., and McClung, C.R.** (1996). The circadian clock gates expression of two *Arabidopsis* catalase genes to distinct and opposite circadian phases. *Mol. Gen. Genet.* **251**, 196-203.
- Zhong, H.H., Painter, J.E., Salomé, P.A., Straume, M., and McClung, C.R.** (1998). Imbibition, but not release from stratification, sets the circadian clock in Arabidopsis seedlings. *Plant Cell* **10**, 2005-2017.
- Zhong, H.H., Resnick, A.S., Straume, M., and McClung, C.R.** (1997). Effects of synergistic signaling by phytochrome A and cryptochrome 1 on circadian clock-regulated catalase expression. *Plant Cell* **9**, 947-955.
- Zhong, H.H., Young, J.C., Pease, E.A., Hangarter, R.P., and McClung, C.R.** (1994). Interactions between light and the circadian clock in the regulation of *CAT2* expression in *Arabidopsis*. *Plant Physiol.* **104**, 889-898.