

## **FORTY-SECOND SUPPLEMENT TO THE AMERICAN ORNITHOLOGISTS' UNION CHECK-LIST OF NORTH AMERICAN BIRDS**

Source: The Auk, 117(3) : 847-858

Published By: American Ornithological Society

URL: [https://doi.org/10.1642/0004-8038\(2000\)117\[0847:FSSTTA\]2.0.CO;2](https://doi.org/10.1642/0004-8038(2000)117[0847:FSSTTA]2.0.CO;2)

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



## AOU Check-list Supplement

The Auk 117(3):847–858, 2000

### FORTY-SECOND SUPPLEMENT TO THE AMERICAN ORNITHOLOGISTS' UNION CHECK-LIST OF NORTH AMERICAN BIRDS

This first Supplement since publication of the 7th edition (1998) of the AOU *Check-list of North American Birds* summarizes changes made by the Committee on Classification and Nomenclature between its reconstitution in late 1998 and 31 January 2000. Because the makeup of the Committee has changed significantly since publication of the 7th edition, it seems appropriate to outline the way in which the current Committee operates. The philosophy of the Committee is to retain the present taxonomic or distributional status unless substantial and convincing evidence is published that a change should be made. The Committee maintains an extensive agenda of potential action items, including possible taxonomic changes and changes to the list of species included in the main text or the Appendix. Many of these are derived from statements of varying taxonomic treatments mentioned in notes in species accounts in the 7th edition. Each member has accepted primary responsibility for certain groups of birds or for particular distributional or other matters. When a suggestion for a change in taxonomic, nomenclatural, or distributional status is published, the member with responsibility for the affected group studies the situation and prepares a proposal for (or against) change. The Committee also considers proposals or suggestions by nonmembers if accompanied by adequate justification or evidence. Proposals, with recommendations, are circulated among the Committee and a period of discussion ensues, mainly by e-mail. Advice may be solicited from colleagues who are not members of the Committee. Eventually, a vote is taken. If approved, the proposal becomes an item for the next Supplement. If the proposal is not accepted, it returns to the agenda to await additional evidence. The Committee attempts to meet annually at the AOU meeting. Members of the Committee are also working toward a future edition of the *Check-list* that will include statements of geographic variation and a treatment at the subspecific level.

Changes in this Supplement fall into eight categories: (1) three species are added to the main list because of new distributional information (*Ardeola bacchus*, *Milvus migrans*, *Emberiza elegans*); (2) 11 species are added to the list because of splitting of species previously on the list (*Sula granti*, *Centrocercus minimus*, *Picoides arizonae*, *Dendroica subita*, *D. delicata*, *Spindalis nigricephala*, *S. dominicensis*, *S. portoricensis*,

*Icterus prothemelas*, *Lonchura cantans*, and *L. atricapilla*); (3) four species are changed (*Caracara cheriway*, *Glaucidium costaricanum*, *Myrmotherula pacifica*, *Pica hudsonia*) and one added (*Caracara lutosa*) by splits from now-extralimital forms; (4) four scientific names of species are changed because of generic reallocation (*Ibycter americanus*, *Stercorarius skua*, *S. maccormicki*, *Molothrus oryzivorus*); (5) one specific name is changed for nomenclatural reasons (*Baeolophus ridgwayi*); (6) the spelling of five species names is changed to make them grammatically correct relative to the generic name (*Jacamerops aureus*, *Poecile atricapilla*, *P. hudsonica*, *P. cincta*, *Buarremon brunneinucha*); (7) one English name is changed to conform to worldwide use (Long-tailed Duck), one is changed by removing an unneeded modifier (White Tern), and five are changed because of species splits (Greater Sage-Grouse, Western Spindalis, Greater Antillean Oriole, Indian Silverbill, Tricolored Munia); and (8) seven species are added to the Appendix (*Oceanodroma monorhis*, *Circus aeruginosus*, *Larus genei*, *L. novae-hollandiae*, *Phaethornis yaruqui*, *Tachycineta albiventer*, *Oryzoborus angolensis*). The 15 additions to the main list bring the number of species recognized as occurring in the *Check-list* area (main list) to 2,023. This Supplement also makes some technical nomenclatural and spelling changes that were not included with errata in the Notice from the Committee in *Auk* 116:282–283, 1999. Literature that provides the basis for the Committee's decisions is cited at the end of the Supplement, and citations not already in the Literature Cited of the 7th edition become additions to it. An updated list of species known from the *Check-list* area may be accessed from <<http://www.nmnh.si.edu/BIRDNET/index.html>> on the internet.

The following changes to the 7th edition (page numbers refer thereto) result from the Committee's actions:

pp. xvii–liv. In the list of bird species known from the *Check-list* area, change 2,008 to 2,023. In the list, insert the following species in the proper position as indicated by the text of this Supplement:

*Sula granti* Nazca Booby.

*Ardeola bacchus* Chinese Pond-Heron.

*Milvus migrans* Black Kite.

†*Caracara lutosa* Guadalupe Caracara.

*Centrocercus minimus* Gunnison Sage-Grouse.  
*Glaucidium costaricanum* Costa Rican Pygmy-Owl.  
*Picoides arizonae* Arizona Woodpecker.  
*Myrmotherula pacifica* Pacific Antwren.  
*Dendroica subita* Barbuda Warbler.  
*Dendroica delicata* St. Lucia Warbler.  
*Spindalis nigricephala* Jamaican Spindalis.  
*Spindalis dominicensis* Hispaniolan Spindalis.  
*Spindalis portoricensis* Puerto Rican Spindalis.  
*Emberiza elegans* Yellow-throated Bunting.  
*Icterus prothemelas* Black-cowled Oriole.  
*Lonchura cantans* African Silverbill (I).  
*Lonchura atricapilla* Chestnut Munia (I).

Change the following English names:

*Clangula hyemalis* Long-tailed Duck.  
*Centrocercus urophasianus* Greater Sage-Grouse.  
*Gygis alba* White Tern.  
*Spindalis zena* Western Spindalis.  
*Icterus dominicensis* Greater Antillean Oriole.  
*Lonchura malabarica* Indian Silverbill (I).  
*Lonchura malacca* Tricolored Munia (I).

Change the following scientific names, with no change in English names:

*Daptrius americanus* to *Ibycter americanus*  
*Caracara plancus* to *Caracara cheriway*  
*Catharacta skua* to *Stercorarius skua*  
*Catharacta maccormicki* to *Stercorarius maccormicki*  
*Jacamerops aurea* to *Jacamerops aureus*  
*Pica pica* to *Pica hudsonia*  
*Poecile atricapillus* to *Poecile atricapilla*  
*Poecile hudsonicus* to *Poecile hudsonica*  
*Poecile cinctus* to *Poecile cincta*  
*Baeolophus griseus* to *Baeolophus ridgwayi*  
*Buarremon brunneinuchus* to *Buarremon brunneinucha*  
*Scaphidura oryzivora* to *Molothrus oryzivorus*

Delete the following names:

*Glaucidium jardiinii* Andean Pygmy-Owl.  
*Myrmotherula surinamensis* Streaked Antwren.

Move *Icterus bullockii* to position following *I. pus-tulatus*

p. 28. *Sula granti* is recognized as a species distinct from *S. dactylatra*, as proposed by Pitman and Jehl (1998), on the basis of differences in bill color, size and proportions, some color characters in both juvenal and definitive plumages, preferred nesting habitat, and pelagic nonbreeding ranges. Modify the breeding distribution of *Sula dactylatra* by removing the phrase "and in the Galapagos" and the word "Ecuador." Add to the Notes under *S. dactylatra*: Some old sight reports may pertain to *S. granti* (e.g. Roberson 1998). Insert the following after the account for *S. dactylatra*:

*Sula granti* Rothschild. Nazca Booby.

*Sula granti* Rothschild, 1902, Bull. Brit. Ornithol. Club 13: 7. (Galapagos Archipelago = Culpepper Island, fide Hartert, Nov. Zool. 32: 274, 1925.)

**Habitat.**—Pelagic waters; nests on cliffs on islands.

**Distribution.**—Breeds in the eastern Pacific on the Galapagos Islands, Isla La Plata, Ecuador, and Malpelo Island off Colombia, and in small numbers on Clipperton Island and the Revillagigedos.

**Ranges** at sea off Middle America in the southern Gulf of California and from Colima, Mexico, to Ecuador.

**Notes.**—Formerly considered a subspecies of *S. dactylatra*, but separated by Pitman and Jehl (1998). See comments under *S. dactylatra*.

p. 32. In the Notes under the genus *Phalacrocorax*, the name *Hypoleucus* should be *Hypoleucos*.

p. 44. Before the genus *Butorides*, insert:

#### Genus *ARDEOLA* Boie

*Ardeola* Boie, 1822, Isis von Oken, col. 559. Type, by monotypy, *Ardea ralloides* Scopoli.

*Ardeola bacchus* (Bonaparte). Chinese Pond-Heron.

*Buphus bacchus* Bonaparte, 1855, Consp. Gen. Avium, 2:127. (Malay Peninsula.)

**Habitat.**—Marshland, rice fields, riverbanks, mangroves, tidepools, and margins of fish ponds.

**Distribution.**—Breeds from central Manchuria and Inner Mongolia south through eastern China and Taiwan to Assam, northern Indochina, and Hainan.

**Winters** in the southern part of the breeding range and to Thailand, the Malay Peninsula, Myanmar (Burma), Sumatra, Java, Borneo, and Sulawesi.

Wanders to Siberia, Korea, Japan, and Sri Lanka.

Accidental on St. Paul Island, Pribilofs, Alaska, 4–9 August 1996 (Gibson and Kessel 1997, Hoyer and Smith 1997).

p. 81. The Committee was petitioned by a group of biologists from the U.S. Fish and Wildlife Service in Alaska to change the English name of *Clangula hyemalis* from Oldsquaw to Long-tailed Duck, the name used for the species outside of North America. The basis for the petition was that the species is declining in numbers in Alaska, and conservation management plans require the help and cooperation of Native Americans. The biologists were concerned that the name Oldsquaw would offend the Native Americans. Requests to change the name had been made to the Committee in past years by some who consider the word "squaw" to be offensive. The Committee declines to consider political correctness alone in changing long-standing English names of birds but

is willing in this instance to adopt an alternative name that is in use in much of the world.

Change English name of *Clangula hyemalis* from Oldsquaw to Long-tailed Duck, to conform with English usage in other parts of the world. Change Notes to read: Formerly known as Oldsquaw in North America.

p. 91. After *Ictinia plumbea*, insert the following account:

Genus **MILVUS** Lacépède, 1799

*Milvus* Lacépède, 1799, Tab. Mamm. Ois., p. 4. Type, by tautonomy, *Falco milvus* Linnaeus.

***Milvus migrans*** (Boddaert). Black Kite.

*Falco migrans* Boddaert, 1783, Table Planches Enlum., p. 28. Based on "Le Milan noir" of Daubenton, Planches Enlum., pl. 472. (France.)

**Habitat.**—Riparian areas, open woodland, forest edge, coastal wetlands, farmland, garbage dumps, and cities.

**Distribution.**—Breeds (*migrans* group) in Eurasia from Finland to western Siberia and south to islands in the Mediterranean Sea and northwestern Africa, the Near East and Arabia, east through India and south to Sri Lanka and southwestern China and south through southeast Asia to Australia, and in much of Africa; and (*lineatus* group) in eastern Siberia and Japan south through China to northern India, Burma, and the Ryukyu Islands.

*Winters* (*migrans* group) from southern Eurasia to southern Africa and southern Australia; and (*lineatus* group) from southern Iraq to China and Japan, south to southern India, Sri Lanka, and southeast Asia.

Accidental (group unknown) in the Mariana Islands, and (*lineatus* group) in Hawaii on Sand Island, Midway, 1994–1995 (R. L. Pyle pers. comm.) and 1998 (Field Notes, 1998, pp. 147, 261, 272). Photographs documenting both Midway occurrences are on file in the Hawaii Rare Bird Documentary Photograph File at the B. P. Bishop Museum, Honolulu.

**Notes.**—Stresemann and Amadon in Mayr and Cottrell (1979) and Sibley and Monroe (1990) considered the two groups to be allopecies, *M. migrans* (Boddaert, 1783) [Black Kite] and *M. lineatus* (Gray, 1831) [Black-eared Kite]. Sibley and Monroe (1993: 40) further separated the birds in Arabia and eastern and southern Africa as a group *aegyptius* (Gmelin, 1788) [Yellow-billed Kite] within *M. migrans*.

p. 105. In the Notes under Family Falconidae, change Griffiths (1994) to Griffiths (1994a).

p. 106. The Red-throated Caracara is moved from the genus *Daptrius* to the genus *Ibycter*. Members of

these genera differ in syringeal morphology and mitochondrial DNA (Griffiths 1994b, 1999) and in ecology (Brown and Amadon 1968). The genetic studies show that *Daptrius* is paraphyletic if *americanus* is included. Replace the heading and citation for Genus *Daptrius* with:

Genus **IBYCTER** Vieillot

*Ibycter* Vieillot, 1816, Analyse, p. 22. Type, by monotypy, *Falco americanus* Boddaert.

Change the species heading from *Daptrius americanus* (Boddaert) to *Ibycter americanus* (Boddaert). Change the Notes at the end of the species account to read: Formerly placed in the genus *Daptrius* Vieillot, 1816, but separated on the suggestions of Brown and Amadon (1968) and Griffiths (1994b, 1999).

p. 106. The three groups in *Caracara plancus* are recognized as distinct species following an analysis of plumage, morphology, and reported hybridization by Dove and Banks (1999). Replace the account for now extralimital *C. plancus* with the following two accounts:

***Caracara cheriway*** (Jacquin). Crested Caracara.

*Falco cheriway* Jacquin, 1784, Beytr. Gesch. Vögel., p. 17, pl. 4. (Aruba.)

**Habitat.**—Arid Lowland Scrub, Arid Montane Scrub, Low Seasonally Wet Grassland, Second-growth Scrub, lowland pine savanna (0–3,000 m; Tropical to Temperate zones).

**Distribution.**—As *cheriway* group in *plancus* account.

**Notes.**—Formerly combined with the South American *Caracara plancus* (Miller, 1777) [Southern Caracara] and *C. lutosus* as Crested Caracara, but separated by Dove and Banks (1999).

†***Caracara lutosus*** (Ridgway). Guadalupe Caracara.

*Polyborus lutosus* Ridgway, 1876, Bull. U. S. Geol. Geogr. Surv. Terr. 1:459. (Guadalupe Island, Lower California.)

**Habitat.**—Arid Lowland Scrub.

**Distribution.**—as *lutosus* group in *plancus* account.

**Notes.**—See notes under *C. cheriway*.

p. 119. On the basis of genetic differences (Kahn et al. 1999, Oyler-McCance et al. 1999) and differences in size (Hupp and Braun 1991), courtship behavior (calls, strut rate; Young et al. 1994), tail pattern, and modified body plumes (Young et al. 2000), the Gunnison Sage-Grouse, *Centrocercus minimus*, is recognized as a species distinct from the Greater Sage-Grouse, *C. urophasianus*. Change the English name of *C. urophasianus* to Greater Sage-Grouse and modify the Distribution and Notes sections of the account for *C. urophasianus* as follows:

**Distribution.**—*Resident* locally (formerly more widespread) in eastern (Mono County) and north-eastern California, and from eastern Oregon, central Washington, southeastern Alberta, southwestern Saskatchewan, Montana, and southwestern North Dakota south to central Nevada, northern Utah, Wyoming, northern Colorado, and southwestern South Dakota. Formerly north to southern British Columbia (Okanagan Valley) and east to western Nebraska.

**Notes.**—Formerly included *C. minimus* and known as the Sage Grouse.

Insert the following new account for *C. minimus*:

***Centrocercus minimus*** Bradbury and Vehrencamp.  
Gunnison Sage-Grouse.

*Centrocercus minimus* Bradbury and Vehrencamp, 1998, Principles of Animal Communication, cover, iv, 770. (Gunnison, Colorado.)

**Habitat.**—Sagebrush, sagebrush dominated shrub-steppe.

**Distribution.**—*Resident* locally in the Gunnison Basin and southwestern Colorado south of the Eagle and Colorado rivers, and in adjacent southeastern Utah east of the Colorado River. Formerly widespread but discontinuous in southern Colorado and extreme eastern Utah. Early records of sage-grouse from northeastern Arizona, southwestern Kansas, northern New Mexico, and western Oklahoma are presumed to have been of this species rather than *C. urophasianus* (Young et al. 2000).

**Notes.**—Formerly considered part of *C. urophasianus* but shown to be distinct by Hupp and Braun (1991), Young et al. (1994), Kahn et al. (1999), Oyler-McCance et al. (1999), and Young et al. (2000).

p. 120. In the citation for *Lagopus mutus*, change the date of publication to 1781 (fide Tyrberg 1998).

p. 181. A series of studies on molecular genetics (Cohen et al. 1997, Andersson 1999, Braun and Brumfield 1998) have shown that relationships of the skuas and jaegers are best expressed by placing all of the species in a single genus. This supports earlier suggestions based on studies of behavior (Andersson 1973) and parasite faunas (Cohen et al. 1997). The genus *Catharacta* is merged into the genus *Stercorarius*. Delete the words “genera and” from the Notes under Subfamily Stercorariinae. Move the generic heading for the genus *Stercorarius* on p. 182 to replace the generic name *Catharacta* Brünnich, and add the citations for *Catharacta* Brünnich and *Megalestris* Bonaparte to the synonymy of *Stercorarius* in the order *Stercorarius*, *Catharacta*, *Coprotheres*, *Megalestris*. Under the generic heading and synonymy, insert:

**Notes.**—We follow Andersson (1999) in merging *Catharacta* into *Stercorarius* on the basis of studies of molecular genetics and reconsideration of other data.

Replace the species heading *Catharacta skua* Brünnich with *Stercorarius skua* (Brünnich). Throughout the account, change group name *antarctica* to *antarcticus* and replace the generic initial *C.* with *S.*

p. 182. Replace the species heading *Catharacta maccormicki* (Saunders) with *Stercorarius maccormicki* Saunders. In the Notes for that species, change the generic initial *C.* to *S.* and add: Formerly placed in the genus *Catharacta*.

In the Notes under *Stercorarius pomarinus*, delete “(*Catharacta*)” and add: Braun and Brumfield (1998) suggested that *pomarinus* be placed in the monotypic genus *Coprotheres*, between *Catharacta* and *Stercorarius*. The complex phylogenetic relationships are best expressed by considering all species in a single genus (Andersson 1999).

p. 207. Change the English name of *Gygis alba*, Common White-Tern, to White Tern.

pp. 257–258. The distribution of *Bubo virginianus* is restated to recognize existence of a gap in Central America (Olson 1997). Modify the paragraph as follows:

**Distribution.**—*Breeds* from . . . and Newfoundland south to Honduras, rarely in Guatemala and El Salvador, and perhaps in north-central Nicaragua, and from northern Colombia and Venezuela south to Tierra del Fuego (except for most of Amazonia). Absent from the West Indies, most other islands, and apparently much of Central America.

p. 259. Pygmy-owls in Costa Rica and Panama previously assigned to *Glaucidium jardinii* are recognized as a distinct species, *G. costaricanum*, following Robbins and Stiles (1999). Vocalizations and biochemical studies indicate a closer relationship to members of the *G. gnoma* complex than to *G. jardinii* (König 1991, Heidrich et al. 1995, Robbins and Stiles 1999). However, *G. costaricanum* is separated from *G. gnoma* on the basis of differences in plumage color, morphology, voice, habitat, and mtDNA (Robbins and Stiles 1999). Replace the account for *G. jardinii*, now extralimital, with the following:

***Glaucidium costaricanum*** L. Kelso. Costa Rican Pygmy-Owl.

*Glaucidium jardinii costaricanum* L. Kelso, 1937, Auk 54:304. (Costa Rica.)

**Habitat.**—Montane Evergreen Forest (2000–3400 m; Subtropical and Temperate zones).

**Distribution.**—*Resident* in the mountains of central Costa Rica south and east to western Panama (Chiriquí and Veraguas).

**Notes.**—Formerly considered a subspecies of *G. jardinii* (Bonaparte, 1855) [Andean Pygmy-Owl], but

shown to be more closely allied to the *G. gnoma* complex (Robbins and Stiles 1999).

p. 292. Place a dagger, to indicate extinct status, before the name *Chlorostilbon bracei*.

p. 328. Change the specific name of the Great Jacamar to *aureus*, to agree in gender with the masculine generic name *Jacameroops*.

p. 340. *Picoides arizonae* is split from the disjunct *P. stricklandi* on the basis of differences in morphology, behavior, and habitat (Davis 1965, Ligon 1968, Johnson et al. 1999). Insert the following account before that of *P. stricklandi*:

***Picoides arizonae*** (Hargitt). Arizona Woodpecker.

*Picus arizonae* Hargitt, 1886, Ibis, p. 115. (Santa Rita Mts., Arizona.)

**Habitat.**—Oak and pine-oak woodland and riparian vegetation (1,200–2,400 m; upper Subtropical zone).

**Distribution.**—That of the *arizonae* group in 7th ed. account of *P. stricklandi*.

**Notes.**—Often considered conspecific with *P. stricklandi* (Davis 1965, Short 1982), but see Ligon (1968) and Johnson et al. (1999). Also called Brown-backed Woodpecker, but that name should be restricted to the African *Dendrocopos obsoletus* (Wagler, 1829).

In *Picoides stricklandi* account, change habitat to read “Coniferous forest (2,500–4,100 m; Temperate Zone).” Remove *arizonae* group from Distribution. Change Notes to read: “See comments under *P. arizonae*.”

p. 364. *Myrmotherula pacifica* is separated as a species distinct from *M. surinamensis*, now extralimital, on the basis of differences in vocalizations and in color patterns of females (Isler et al. 1999). Replace the account of *M. surinamensis* with the following:

***Myrmotherula pacifica*** Hellmayr. Pacific Antwren.

*Myrmotherula surinamensis pacifica* Hellmayr, 1911, Proc. Zool. Soc. London, p. 1159 (Buenaventura, Chocó, [depto. Valle], W. Colombia.)

**Habitat.**—River-edge Forest, Tropical Lowland Evergreen Forest Edge, Secondary Forest (0–600 m; Tropical zone).

**Distribution.**—*Resident* from Panama (entire Caribbean slope, and Pacific drainage west to western Panamá province) south on the west side of the Andes to southwestern Ecuador (northwestern Azuay), and east in the northern Colombian lowlands to the Rio Magdalena Valley (Santander).

**Notes.**—Formerly considered a subspecies of *Myrmotherula surinamensis* (Gmelin, 1788) [Streaked Ant-

wren] but separated by Isler et al. (1999). The form *multostriata* Sclater, 1858 [Amazonian Streaked-Antwren] of southern and western Amazonia also is split from *M. surinamensis* [Guianan Streaked-Antwren] of northeastern Amazonia by those authors.

p. 368. In the citation for the genus *Myrmornis*, the page number 180 should be 188.

p. 448. North American *Pica hudsonia* is treated as a species distinct from Old World *P. pica*, which becomes extralimital, on the basis of a number of morphologic, behavioral, and genetic characters (Birkhead 1991, Enggist-Dublin and Birkhead 1992, Zink et al. 1995).

Replace *P. pica* account with account of *P. hudsonia*.

***Pica hudsonia*** (Sabine). Black-billed Magpie.

*Corvus hudsonius* Sabine, 1823, in Franklin, Narr. Journ. Polar Sea, 1823, p. 671. (Cumberland House, Hudson Bay [=Cumberland House, Saskatchewan].)

**Habitat.**—Open country with scattered trees, riparian and open woodland, forest edge, and farmlands.

**Distribution.**—*Resident* from south-coastal Alaska (west to the Alaska Peninsula and Shumagin Islands), southern Yukon, northern Alberta, central Saskatchewan, central Manitoba, extreme southwestern Ontario, and northern Minnesota south to northeastern and east-central California (to Inyo County), south-central Nevada, Utah, extreme northeastern Arizona (Apache County, formerly more widespread), northern New Mexico, western (casually northeastern) Oklahoma, central Kansas, and Nebraska (except southeastern). Absent from coastal areas and regions from southeastern Alaska southward and west of the Cascade range and Sierra Nevada.

Wanders as in paragraph at top of p. 449.

**Notes.**—Formerly considered a subspecies of Old World *Pica pica* (Linnaeus, 1758) [Eurasian Magpie], but separated on the basis of differences summarized by Birkhead (1991) and Enggist-Dublin and Birkhead (1992). Vocal and behavioral data suggest that *P. hudsonia* is more closely related to *P. nuttalli* than to *P. pica*. All taxa in *Pica* were considered conspecific by Phillips (1986).

p. 449. Change Notes for *P. nuttalli* to read: “See comments under *P. hudsonia*, and Verbeek (1972).”

p. 455. Add to the distribution of *Progne cryptoleuca* a statement “Accidental in Florida (Key West, 9 May 1895).” Add to the Notes: Other Florida records mentioned in earlier *Check-lists* are of *P. subis* (Banks 2000).

pp. 461–462. In the headings for *Petrochelidon pyr-*

*rhonota* and *P. fulva*, the name Vieillot should be in parentheses, (Vieillot).

In the account of *Petrochelidon fulva*, the group name *pelodoma* should be changed to *pallida* because the latter is not preoccupied by the former in the genus *Petrochelidon*. The sentence to that effect should be deleted from the Notes. In the Notes, the group name “*P. pelodoma* Brooke, 1974” should be changed to “*P. pallida* Nelson, 1902.” *P. rufocollaris* Peale, 1848, should be *P. rufocollaris* (Peale, 1848).

p. 463. In the Notes at the top of the page, for *Hirundo rustica*, change *H. dumicola* to *H. domicola*.

pp. 463–465. The generic name *Poecile* is feminine, and adjectival species names must agree in gender. Change the specific names *atricapillus*, *hudsonicus*, and *cinctus* to *atricapilla*, *hudsonica*, and *cincta*. In the Notes under *P. atricapilla*, *P. montanus* becomes *P. montana*.

p. 466. Change *Baeolophus griseus* (Ridgway) to *Baeolophus ridgwayi* (Richmond). The name *griseus* is permanently invalidated because it was replaced as a junior secondary homonym before 1961 (ICZN 1999, Art. 59.3).

p. 472. In the Notes under *Campylorhynchus zonatus*, at the top of the page, the citation for *C. fasciatus* should be (Swainson, 1838). In the Notes under *Campylorhynchus chiapensis*, the citation for *C. griseus* should be (Swainson, 1838).

p. 517. In the citation for *Oreoscoptes montanus*, change C. K. Townsend to J. K. Townsend.

p. 546. On the basis of genetic differences, combined with differences in plumage and morphology and probably song (Curson et al. 1994, Lovette et al. 1998, Lovette and Bermingham 1999), *Dendroica adelaidae* is divided into three species. In the account for *D. adelaidae*, change Distribution to read: *Resident* on Puerto Rico (including Vieques Island). Change Notes to: Formerly included *D. subita* and *D. delicata*, now considered distinct species (Lovette et al. 1998, Lovette and Bermingham 1999). Lowery and Monroe in Paynter (1968) proposed that *D. adelaidae* (including *subita* and *delicata*) and *D. graciae* were each other's closest relatives, but mitochondrial DNA data (Lovette and Bermingham 1999) do not support this relationship.

Insert the following two accounts after *Dendroica adelaidae*:

***Dendroica subita*** Riley. Barbuda Warbler.

*Dendroica subita* Riley, 1904, Smiths. Misc. Coll. 47: 289. (Barbuda.)

**Habitat.**—Arid Lowland Scrub, Riparian Thickets.

**Distribution.**—Resident on Barbuda in the Lesser Antilles.

**Notes.**—See comments under *D. adelaidae*.

***Dendroica delicata*** Ridgway. St. Lucia Warbler.

*Dendroica adelaidae delicata* Ridgway, 1883, Proc. U.S. Nat. Mus. 5 (1882): 525. (St. Lucia.)

**Habitat.**—Arid Lowland Scrub, Tropical Lowland Evergreen Forest (0–700 m).

**Distribution.**—Resident on St. Lucia in the Lesser Antilles.

**Notes.**—See comments under *D. adelaidae*.

p. 581. *Spindalis zena* is divided into four species, following suggestions by Garrido et al. (1997), based on differences in mensural data, coloration, body mass, and vocalizations. This treatment has been followed by Raffaele et al. (1998). In the citation for the genus *Spindalis*, delete “= *Fringilla zena* Linnaeus.” Insert the following after the heading and citation for the genus *Spindalis*:

**Notes.**—To avoid long hyphenated compound English names, we revert to the English group name *Spindalis*, used (e.g. Bond 1936) before the populations were merged (without comment) by Bond (1947).

Change the English name of *Spindalis zena* to Western *Spindalis*. Change the Distribution of *S. zena* to: *Resident* in the Bahama Islands (Grand Bahama, Great Abaco, Little Abaco, and Green Turtle Cay in the northern Bahamas, from the Berry Islands south to Great Inagua in the southern Bahamas), Providenciales in the Turks and Caicos, Cuba (including the Isle of Pines and numerous keys), Grand Cayman Island, and Cozumel Island off Quintana Roo, Mexico. Remove “[*zena* group]” from sentence beginning “Ranges . . .”

Change Notes under *S. zena* to read: Formerly included *S. nigricephala*, *S. dominicensis*, and *S. portoricensis*, with the English name Stripe-headed Tanager, but the complex is treated as four allospecies of a superspecies following Garrido et al. (1997).

After the account of *Spindalis zena*, insert the following three accounts:

***Spindalis nigricephala*** (Jameson). Jamaican *Spindalis*.

*Tanagra nigricephala* Jameson, 1835, Edinburgh New Philos. Journ. 19: 213. (Jamaica.)

**Habitat.**—Tropical Montane Forest, Tropical Lowland Evergreen Forest, Secondary Forest (0–1,800 m).

**Distribution.**—*Resident* on Jamaica.

**Notes.**—Formerly considered part of *S. zena*, but separated by Garrido et al. (1997).

***Spindalis dominicensis*** (Bryant). Hispaniolan *Spindalis*.

*Tanagra dominicensis* Bryant, 1867, Proc. Boston Soc. Nat. Hist. 11: 92. (southeast Haiti.)

**Habitat.**—Tropical Montane Forest, Pine Forest, Tropical Lowland Evergreen Forest, Secondary Forest (0–2,300 m).

**Distribution.**—*Resident* on the island of Hispaniola, and on Gonave Island.

**Notes.**—Formerly considered part of *S. zena*, but separated by Garrido et al. (1997).

*Spindalis portoricensis* (Bryant). Puerto Rican Spindalis.

*Tanagra portoricensis* Bryant, 1866, Proc. Boston Soc. Nat. Hist. 10:252. (Puerto Rico.)

**Habitat.**—Tropical Montane Forest, Tropical Lowland Evergreen Forest, Secondary Forest (0–1,050 m).

**Distribution.**—*Resident* on Puerto Rico.

**Notes.**—Formerly considered part of *S. zena*, but separated by Garrido et al. (1997).

p. 601. Change *Buarremon brunneinuchus* to *Buarremon brunneinucha*; this is a noun used in apposition, and its gender does not change. In the Notes under that species, *B. apertus* Wetmore, 1942 should be *B. apertus* (Wetmore, 1942).

p. 629, after the account for *Emberiza rustica*, insert:

*Emberiza elegans* Temminck. Yellow-throated Bunting.

*Emberiza elegans* Temminck, 1835, Planches Color., livr. 98, pl. 583, fig. 1. (Japan.)

**Habitat.**—Open dry deciduous forest on hills and ridges.

**Distribution.**—*Breeds* from southern Siberia, Manchuria, and northern Korea south to southern China.

*Winters* from eastern China, southern Korea, and Japan south to southern China and Burma.

Accidental in Alaska (Attu, in the Aleutian Islands, 25 May 1998; Sykes 1998).

p. 649. *Scaphidura* is merged into *Molothrus* on the basis of several genetic studies (Lanyon 1994, Johnson and Lanyon 1999, Lanyon and Omland 1999). Delete the heading for the genus *Scaphidura* and the Notes under it. Move the citations for the generic names *Scaphidura* and *Psomocolax* to proper chronological positions under the genus *Molothrus*. Change the species heading *Scaphidura oryzivora* (Gmelin) to *Molothrus oryzivorus* (Gmelin).

Change the Notes after the species account to read: Also known as Rice Grackle. Formerly placed in the monotypic genus *Scaphidura*, but shown by genetic data (Johnson and Lanyon 1999, Lanyon and Omland 1999) to be the sister species to the other species of

*Molothrus* in our area and to be more closely related to them than to extralimital *M. rufoaxillaris* Cassin.

p. 649. *Icterus prothemelas* is recognized as a species distinct from *I. dominicensis* and is moved to a position next to *I. spurius* on the basis of genetic data presented by Omland et al. (1999). The analysis by Omland et al. (1999) indicates that the *I. dominicensis* complex may consist of up to four species. However, the exclusion of *I. d. dominicensis* from part of the genetic data set, and the lack of an analysis of vocal and plumage differences among the four island taxa, make it impossible to determine at this time how many biological species should be recognized. Replace the account for *I. dominicensis* with the following:

*Icterus dominicensis* (Linnaeus). Greater Antillean Oriole.

The citation for the species is unchanged.

**Habitat.**—Tropical Lowland Evergreen Forest Edge, Secondary Forest (0–1,000 m; Tropical Zone).

**Distribution.**—*Resident* [*dominicensis* group] on Hispaniola; [*portoricensis* group] on Puerto Rico; [*northropi* group] on Andros, Great Abaco, and Little Abaco in the northern Bahamas; and [*melanopsis* group] on Cuba and the Isle of Pines.

**Notes.**—Groups: *I. dominicensis* (Linnaeus, 1766) [Hispaniolan Oriole]; *I. portoricensis* Bryant, 1866 [Puerto Rican Oriole]; *I. northropi* Allen, 1890 [Bahaman Oriole]; *I. melanopsis* (Wagler, 1829) [Cuban Oriole]. Genetic analysis by Omland et al. (1999) suggests that some or all of the groups may merit specific rank, but further study is needed to clarify their relationships.

p. 650. Insert the following new account before the account for *Icterus spurius*:

*Icterus prothemelas* (Strickland). Black-cowled Oriole.

*Xanthornus prothemelas* Strickland, 1850, in Jardine's Contrib. Ornith., 2, p. 120, pl. 62. (Guatemala.)

**Habitat.**—Tropical Lowland Evergreen Forest Edge, Secondary Forest (0–1,200 m; Tropical Zone).

**Distribution.**—*Resident* from southern Veracruz, northern Oaxaca, Tabasco, Chiapas, and the Yucatan Peninsula south on the Caribbean slope of Central America to extreme western Panama (western Bocas del Toro).

**Notes.**—*Icterus prothemelas* was considered a distinct species until Bond (1947) included it in *I. dominicensis*, without comment but apparently on the basis of its similarity in plumage to *I. d. northropi*. Subsequently treated as a subspecies of *I. dominicensis* but shown by genetic analysis (Omland et al. 1999) to be more closely related to *I. spurius*.



p. 653. So that the sequence of species more closely reflects relationships as determined by molecular genetics (Freeman and Zink 1995, Omland et al. 1999), move the account of *Icterus bullockii* from p. 655 to a position following the account of *I. pustulatus*. Change the Notes under *I. bullockii* to read: See notes under *I. galbula* and *I. abeillei*, with which this species was formerly combined.

p. 682. *Lonchura cantans*, African Silverbill, is separated as a species distinct from *L. malabarica*, which becomes Indian Silverbill, following Restall (1996).

Remove “=*Loxia malabarica* Linnaeus” from citation of *Euodice* in synonymy of *Lonchura*.

Change English name of *Lonchura malabarica* to Indian Silverbill. Replace the account for *L. malabarica* with the following:

**Habitat.**—Dry, grassy brush and scrub.

**Distribution.**—*Resident* from eastern Saudi Arabia and Oman east to Bangladesh and eastern India, and south to Sri Lanka.

Introduced and established on Puerto Rico; reported on St. Croix, Virgin Islands.

**Notes.**—Formerly included *L. cantans*, now considered a distinct species (Harrison 1964, Kakizawa and Watada 1985, Restall 1996), with the name Warbling Silverbill. Also known as White-throated Silverbill or White-throated Munia.

After the account for *Lonchura malabarica*, insert the following new account:

***Lonchura cantans*** (Gmelin). African Silverbill.

*Loxia cantans* Gmelin, 1789, Syst. Nat. 1(2): 859. (Africa. Restricted to Dakar, Senegal, by Sclater and Mackworth-Praed, 1918, Ibis, p. 440.)

**Habitat.**—Savanna, arid scrub and brush, grassland, and around human habitation.

**Distribution.**—*Resident* in Africa south of the Sahara and north of the equatorial Congo Basin from Senegal east to Oman on the Arabian Peninsula and south in eastern Africa to northern Tanzania.

Introduced and established in the Hawaiian Islands (originally on Hawaii, recently spreading to Maui, Lanai, and Molokai, with sight reports from Kauai, Oahu, and Kahoolawe). A pair successfully bred on Merritt Island, Florida, in 1965 (1965, Aud. Field Notes 19: 537), but the species did not become established.

**Notes.**—Formerly included with *Lonchura malabarica* and together called Warbling Silverbill. Hawaiian records were erroneously assigned to *L. malabarica* in 7th edition, but see Falkenmayer (1988).

p. 683. *Lonchura atricapilla* is recognized as a species distinct from *L. malacca*, with the latter species called Tricolored Munia, following Restall (1995).

Change the English name of *Lonchura malacca* to Tricolored Munia and replace the account with the following:

**Habitat.**—Wet and marshy areas with long grasses; rice fields.

**Distribution.**—*Resident* in central and southern India and Sri Lanka.

Introduced and established in Puerto Rico, Jamaica, Hawaiian Islands (Oahu), Venezuela, and Japan. Reportedly introduced or observed in Cuba, Hispaniola, and Martinique (Raffaele et al. 1998), but in the absence of voucher specimens some of these reports may be of *L. atricapilla*. Reported breeding on Merritt Island, Florida (1965, Aud. Field Notes 19: 537), but this record also may refer to *L. atricapilla*.

**Notes.**—Formerly included *L. atricapilla* and known as Chestnut Mannikin, but separated by Restall (1995).

Insert after the account of *L. malacca*:

***Lonchura atricapilla*** (Vieillot). Chestnut Munia.

*Loxia atricapilla* Vieillot, 1807, Ois. Chant., p. 84, pl. 53. (Les Grandes-Indes, restricted to Lower Bengal by Robinson and Kloss, 1924, Jour. Nat. Hist. Soc. Siam 5: 362.)

**Habitat.**—Grassy areas, marshes; rice fields.

**Distribution.**—*Resident* in northern and eastern India, Nepal, Southeast Asia, southern China, Hainan, and Taiwan south to Sri Lanka, the Greater Sund Islands, and the Philippines.

Introduced and established in Puerto Rico, Jamaica, Hawaiian Islands (Oahu, Kauai), Guam, and Palau (Pratt et al. 1987, as *L. malacca*; see Restall 1996).

**Notes.**—Formerly merged with *L. malacca* and known as Chestnut Mannikin, but separated by Restall (1995). See notes and distribution statement under *L. malacca*.

p. 688. *Oceanodroma monorhis* is added to the Appendix. Before the account for *Oceanodroma hornbyi*, insert:

***Oceanodroma monorhis*** (Swinhoe). Swinhoe's Storm-Petrel.

*Thalassidroma monorhis* Swinhoe, 1867, Ibis, p. 386. (near Amoy, China.)

This species, which breeds in the North Pacific and ranges in the Indian Ocean and Arabian Sea, was reported in the western North Atlantic Ocean, south-east of Hatteras, North Carolina, 8 August 1998 (photographs; O'Brien et al. 1999) and perhaps on previous occasions (Brinkley 1995). It has been reported occasionally (since 1983) in the eastern North Atlantic (Cubitt 1995). This species is not well known, and identification from photographs is considered tenuous.

p. 690. *Circus aeruginosus* is added to the Appendix. After the account for *Aythya nyroca*, insert:

***Circus aeruginosus*** (Linnaeus). Western Marsh-Harrier.

*Falco aeruginosus* Linnaeus, 1758, Syst. Nat. (ed. 10) 1: 91. (Europe = Sweden.)

This species of Eurasia and northern Africa was reportedly seen at Chincoteague National Wildlife Refuge, Accomack County, Virginia, on 4 December 1994. Photographs were reportedly obtained but were not published (Shedd et al. 1998).

p. 692. Change *Catharacta chilensis* (Bonaparte) to *Stercorarius chilensis* Bonaparte.

p. 692. *Larus genei* is added to the Appendix. After the account for *Stercorarius chilensis*, insert:

***Larus genei*** Brème. Slender-billed Gull.

*Larus Genei* Brème, 1839, Rev. Zool., p. 321. (Sar-dinia.)

This species of the Mediterranean and Indian Ocean coasts was reportedly seen on Antigua, 24 April 1976 (Holland and Williams 1978, Raffaele et al. 1998:451).

p. 692. *Larus novaehollandiae* is added to Appendix, after *Larus genei*.

***Larus novaehollandiae*** Stevens. Silver Gull.

*Larus Novae-Hollandiae* Stevens, 1826, in Shaw's General Zoology 13, pt. 1, p. 196. (New South Wales.)

This Southern Hemisphere species, also known as Red-billed Gull, is frequently kept in zoos in the United States. A specimen (August 1947) from the mouth of the Genesee River in New York (Beardslee and Mitchell 1965) was thought to be a wanderer but is now considered to have been an escapee (Bull 1974). A bird photographed (NAS Field Notes 51:33, 1997) in Salem County, New Jersey, in autumn 1996, was assumed to have escaped from captivity.

p. 694. *Phaethornis yaruqui* is added to the Appendix. Insert between *Coccyzus lansbergi* and *Anthracothorax viridigula*:

***Phaethornis yaruqui*** (Bourcier). White-whiskered Hermit.

*Trochilus Yaruqui* Bourcier, 1851, Compt. Rend. Acad. Sci. Paris 32:187. (Vicinity of Yaruqui, Ecuador.)

This species of the Chocó region of western Colombia and northwestern Ecuador was reported at Manané, Darién, Panama, 10 July 1996 (Seutin 1998).

p. 696. *Tachycineta albiventer* is added to the Ap-

pendix. Insert between *Melanocorypha calandra* and *Parus major*:

***Tachycineta albiventer*** (Boddaert). White-winged Swallow.

*Hirundo albiventer* Boddaert, 1783, Table Planches Enlum., p. 32. Based on Daubenton, Planches Enlum., pl. 546. (Cayenne.)

This species, widespread in tropical South America, was reported from the Tuira River downstream from Unión Chocó, Darién, Panama, 6 July 1996 (Seutin 1998). There is also a sight report of this species at Schoelcher, Martinique, 10 August 1993 (Feldmann et al. 1999).

p. 698. *Oryzoborus angolensis* is added to the Appendix. Insert before *Icterus nigrogularis*:

***Oryzoborus angolensis*** (Linnaeus). Chestnut-bellied Seed-Finch.

*Loxia angolensis* Linnaeus, 1766, Syst. Nat. (ed. 12) 1:303. Based on "The Black Gros-Beak" Edwards, Glean. Nat. Hist. 3, p. 296, pl. 352. (Angola, error, eastern Brazil suggested by Hellmayr, 1906, Novit. Zool. 13:19.)

Caged birds of this South American species escaped on Martinique and established a small wild breeding population by 1984. Breeding has been reported in 1995 and 1996 (Feldmann et al. 1999), but the population is still small and localized. Photographs have been deposited in VIREO.

pp. 705–730. In the list of French Names for North American Birds:

Change the following scientific names, retaining the French names:

*Daptrius americanus* to *Ibycter americanus*  
*Caracara plancus* to *Caracara cheriway*  
*Catharacta skua* to *Stercorarius skua*  
*Catharacta maccormicki* to *Stercorarius maccormicki*  
*Jacamerops aurea* to *Jacamerops aureus*  
*Pica pica* to *Pica hudsonia*  
*Poecile atricapillus* to *Poecile atricapilla*  
*Poecile hudsonicus* to *Poecile hudsonica*  
*Poecile cinctus* to *Poecile cincta*  
*Baeolophus griseus* to *Baeolophus ridgwayi*  
*Buarremon brunneinuchus* to *Buarremon brunneinucha*  
*Scaphidura oryzivora* to *Molothrus oryzivorus*

Change the French name that accompanies the following scientific name:

*Spindalis zena* to *Zéna à tête rayée*

Insert the following in the appropriate places, as indicated by preceding text:

*Sula granti* Fou de Grant  
*Ardeola bacchus* Crabier chinois

*Milvus migrans* Milan noir  
*Caracara lutosa* Caracara de Guadalupe  
*Centrocercus minimus* Tétraz du Gunnison  
*Glaucidium costaricanum* Chevêchette du Costa Rica  
*Picoides arizonae* Pic d'Arizona  
*Myrmotherula pacifica* Myrmidon du Pacifique  
*Dendroica subita* Paruline de Barbuda  
*Dendroica delicata* Paruline de Sainte-Lucie  
*Spindalis nigricephala* Zéna de Jamaïque  
*Spindalis dominicensis* Zéna d'Hispaniola  
*Spindalis portoricensis* Zéna de Porto Rico  
*Emberiza elegans* Bruant élégant  
*Icterus prothemelas* Oriole monacal  
*Lonchura cantans* Capucin bec-d'argent  
*Lonchura atricapilla* Capucin à tête noire  
*Oceanodroma monorhis* Océanite de Swinhoe.  
*Circus aeruginosus* Busard des roseaux  
*Larus genei* Goéland railleur  
*Larus novaehollandiae* Mouette argentée  
*Phaethornis yaruqui* Ermite yaruqui  
*Tachycineta albiventer* Hirondelle à ailes blanches  
*Oryzoborus angolensis* Sporophile curio

Move *Icterus bullockii* to position following *I. pustulatus*

Delete the entries for the following:

*Glaucidium jordinii*  
*Myrmotherula surinamensis*

p. 742. Replace GRIFFITHS, C. 1994 with GRIFFITHS, C. S. 1994a.

The Committee has discussed most agenda items that have accumulated since the 7th edition. Changes in treatment were delayed on some items with the hope that more convincing evidence would be forthcoming. Proposals considered but not yet accepted by the Committee include the following: separation of *Pterodroma heraldica* from *P. arminjoniana*; separation of *Puffinus newelli* from *P. auricularis*; separation of *Numenius hudsonicus* from *N. phaeopus*; separation of *Cuculus optatus* from *C. saturatus*; separation of the extralimital population *magellanicus* from *Bubo virginianus*; separation of *Glaucidium gnoma* into two or more species; division of *Corvus palmarum* into two species; division of *Chasiempis sandwichensis* into three species; removal of *Troglodytes troglodytes* to the genus *Nannus*; merger of *Myadestes woahensis* into *M. lanaiensis*; separation of *Turdus graysoni* from *T. rufo-palliatu*; separation of *Spizella taverneri* from *S. breweri*; revision of generic relationships in the Emberizidae; and separation of *Loxia megaplaga* from *L. leucoptera*. Several other matters published late in 1999 have been added to the agenda for consideration in the next two years.

**Acknowledgments.**—Michel Gosselin is serving the Committee as its authority for French names, and Normand David is serving as authority for classical languages, especially relative to gender of generic

names. We thank all those who have called our attention to errors in the 7th edition and those who have helped in the preparation of this Supplement. This actually is almost everyone we have spoken with in the past two years, but we particularly thank D. G. Ainley, C. L. Braun, M. B. Braun, J. Choe, W. S. Clark, R. A. Erickson, D. D. Gibson, M. J. Iliff, M. L. Isler, P. R. Isler, J. R. Jehl, Jr., A. R. Keith, N. K. Klein, A. Knox, I. J. Lovette, S. L. Olson, M. A. Patten, R. B. Payne, J. N. Penhallurick, N. J. Pharris, H. D. Pratt, P. Pyle, R. L. Pyle, R. Restall, R. Righter, M. B. Robbins, P. W. Smith, S. O. Williams III, and J. R. Young.

#### LITERATURE CITED

- ANDERSSON, M. 1973. Behaviour of the Pomarine Skua *Stercorarius pomarinus* Temm. with comparative remarks on Stercorariinae. *Ornis Scandinavica* 4:1–16.
- ANDERSSON, M. 1999. Phylogeny, behaviour, plumage evolution and neoteny in skuas Stercorariidae. *Journal of Avian Biology* 30:205–215.
- BANKS, R. C. 2000. The Cuban Martin in Florida. *Florida Field Naturalist* 28: in press.
- BEARDSLEE, C. S., AND H. D. MITCHELL. 1965. Birds of the Niagara Frontier Region. *Bulletin of the Buffalo Society of Natural Science* Vol. 22.
- BIRKHEAD, T. R. 1991. The magpies. T & A D Poyser, London.
- BOND, J. 1936. Birds of the West Indies. Academy of Natural Sciences, Philadelphia.
- BOND, J. 1947. Field guide to birds of the West Indies. Macmillan, New York.
- BRAUN, M. J., AND R. T. BRUMFIELD. 1998. Enigmatic phylogeny of skuas: An alternative hypothesis. *Proceedings of the Royal Society of London Series B* 265:995–999.
- BRINKLEY, E. S. 1995. Dark-rumped storm-petrels in the North Atlantic. *Birding* 27:95–97.
- BROWN, L., AND D. AMADON. 1968. Eagles, hawks, and falcons of the world. Country Life Books, Feltham, United Kingdom.
- BULL, J. 1974. Birds of New York State. Doubleday/Natural History Press. Reissued in 1985, Cornell University Press, Ithaca, New York.
- COHEN, B. L., A. J. BAKER, K. BLECHSCHMIDT, D. L. DITTMANN, R. W. FURNESS, J. A. GERWIN, A. J. HELBIG, J. DE KORTE, H. D. MARSHALL, R. L. PALMA, H.-U. PETER, R. RAMLI, I. SIEBOLD, M. S. WILLCOX, R. H. WILSON, AND R. M. ZINK. 1997. Enigmatic phylogeny of skuas (Aves: Stercorariidae). *Proceedings of the Royal Society of London Series B* 264:181–190.
- CUBITT, M. G. 1995. Swinhoe's Storm-Petrels at Tyne-mouth: New to Britain and Ireland. *British Birds* 88:342–348.
- CURSON, J., D. QUINN, AND D. BEADLE. 1994. Warblers of the Americas. Houghton Mifflin, Boston.
- DAVIS, J. 1965. Natural history, variation, and distri-

- bution of the Strickland's Woodpecker. *Auk* 82: 537-590.
- DOVE, C. J., AND R. C. BANKS. 1999. A taxonomic study of Crested Caracaras (Falconidae). *Wilson Bulletin* 111:330-339.
- ENGGIST-DUBLIN, P., AND T. R. BIRKHEAD. 1992. Differences in the calls of European and North American Black-billed Magpies and the Yellow-billed Magpie. *Bioacoustics* 4:185-194.
- FALKENMAYER, K. 1988. Problems of nomenclature and identification of introduced birds in Hawaii: A case study of two estrildid species. *Elepaio* 48: 91-93.
- FELDMANN, P., E. BENITO-ESPINAL, AND A. R. KEITH. 1999. New bird records from Guadeloupe and Martinique, West Indies. *Journal of Field Ornithology* 70:80-94.
- FREEMAN, S., AND R. M. ZINK. 1995. Phylogenetic study of the blackbirds based on variation in mitochondrial DNA restriction sites. *Systematic Biology* 44:409-420.
- GARRIDO, O. H., K. C. PARKES, G. B. REYNARD, A. KIRKCONNELL, AND R. SUTTON. 1997. Taxonomy of the Stripe-headed Tanager, genus *Spindalis* (Aves: Thraupidae) of the West Indies. *Wilson Bulletin* 109:561-594.
- GIBSON, D. D., AND B. KESSEL. 1997. Inventory of the species and subspecies of Alaska birds. *Western Birds* 28:45-95.
- GRIFFITHS, C. S. 1994a. Monophyly of the Falconiformes based on syringeal morphology. *Auk* 111: 787-805.
- GRIFFITHS, C. S. 1994b. Syringeal morphology and the phylogeny of the Falconidae. *Condor* 96:127-140.
- GRIFFITHS, C. S. 1999. Phylogeny of the Falconidae inferred from molecular and morphological data. *Auk* 116:116-130.
- HARRISON, C. J. O. 1964. The taxonomic status of the African Silverbill *Lonchura cantans* and the Indian Silverbill *Lonchura malabarica*. *Ibis* 106:462-468.
- HEIDRICH, P., C. KÖNIG, AND M. WINK. 1995. Bioakustik, Taxonomie, und molekulare Systematik amerikanischer Sperlingskäuze (Strigidae: *Glaucidium* spp.). *Stuttgarter Beiträge zur Naturkunde Serie A (Biologie)* 534:1-47.
- HOLLAND, C. S., AND J. M. WILLIAMS. 1978. Observations on the birds of Antigua. *American Birds* 32:1095-1105.
- HOYER, R. C., AND S. D. SMITH. 1997. Chinese Pond-Heron in Alaska. *Field Notes* 51:953-956.
- HUPP, J. W., AND C. E. BRAUN. 1991. Geographic variation among Sage Grouse in Colorado. *Wilson Bulletin* 103:255-261.
- INTERNATIONAL COMMISSION ON ZOOLOGICAL NOMENCLATURE. 1999. International Code of Zoological Nomenclature, 4th ed. London.
- ISLER, M. L., R. ISLER, AND B. M. WHITNEY. 1999. Species limits in antbirds (Passeriformes: *Thamnophilidae*): The *Myrmotherula surinamensis* complex. *Auk* 116:83-96.
- JOHNSON, K. P., AND S. M. LANYON. 1999. Molecular systematics of the grackles and allies, and the effect of additional sequence (cyt *B* and ND2). *Auk* 116:759-768.
- JOHNSON, R. R., L. T. HAIGHT, AND J. D. LIGON. 1999. Strickland's Woodpecker (*Picoides stricklandi*). In *The birds of North America*, no. 474 (A. Poole and F. Gill, Eds.). Academy of Natural Sciences, Philadelphia, and American Ornithologists' Union, Washington, D.C.
- KAHN, N. W., C. E. BRAUN, J. R. YOUNG, S. WOOD, D. R. MATA, AND T. W. QUINN. 1999. Molecular analysis of genetic variation among large- and small-bodied Sage Grouse using mitochondrial control-region sequences. *Auk* 116:819-824.
- KAKIZAWA, R., AND R. WATADA. 1985. The evolutionary genetics of the Estrildidae. *Journal of the Yamashina Institute for Ornithology* 17:143-158.
- KÖNIG, C. 1991. Zur Taxonomie und Ökologie der Sperlingskäuze (Strigidae: *Glaucidium* spp.) des Andenraumes. *Ökologie der Vögel* 13:15-76.
- LANYON, S. M. 1994. Polyphyly of the blackbird genus *Agelaius* and the importance of assumptions of monophyly in comparative studies. *Evolution* 48:679-693.
- LANYON, S. M., AND K. E. OMLAND. 1999. A molecular phylogeny of the blackbirds (Icteridae): Five lineages revealed by cytochrome-*B* sequence data. *Auk* 116:629-639.
- LIGON, J. D. 1968. Observations on Strickland's Woodpecker, *Dendrocopos stricklandi*. *Condor* 70: 83-84.
- LOVETTE, I. J., AND E. BERMINGHAM. 1999. Explosive speciation in the New World *Dendroica* warblers. *Proceedings of the Royal Society of London Series B* 266:1629-1636.
- LOVETTE, I. J., E. BERMINGHAM, G. SEUTIN, AND R. E. RICKLEFS. 1998. Evolutionary differentiation in three endemic West Indian warblers. *Auk* 115: 890-903.
- MAYR, E., AND G. W. COTTRELL (Eds.). 1979. Checklist of birds of the world, vol. 1, 2nd ed. Museum of Comparative Zoology, Cambridge, Massachusetts.
- O'BRIEN, M., J. B. PATTESON, G. L. ARMISTEAD, AND G. B. PEARCE. 1999. Swinhoe's Storm-Petrel. *North American Birds* 53:6-10.
- OLSON, S. L. 1997. Avian biogeography in the islands of the Pacific coast of western Panama. Pages 69-82 in *The era of Allan R. Phillips: A Festschrift* (R. W. Dickerman, compiler). Albuquerque, New Mexico.
- OMLAND, K. E., S. M. LANYON, AND S. J. FRITZ. 1999. A molecular phylogeny of the New World Orioles (*Icterus*): The importance of dense taxon

- sampling. *Molecular Phylogeny and Evolution* 12:224–239.
- OYLER-MCCANCE, S. J., N. W. KAHN, K. P. BURNHAM, C. E. BRAUN, AND T. W. QUINN. 1999. A population genetic comparison of large- and small-bodied Sage Grouse in Colorado using microsatellite and mitochondrial DNA markers. *Molecular Ecology* 8:1457–1466.
- PAYNTER, R. J., JR. (Ed.). 1968. Check-list of birds of the world, vol. 14. Museum of Comparative Zoology, Cambridge, Massachusetts.
- PHILLIPS, A. R. 1986. The known birds of North and Middle America. Part 1. Published by the author, Denver, Colorado.
- PITMAN, R. L., AND J. R. JEHL, JR. 1998. Geographic variation and reassessment of species limits in the “Masked” Boobies of the eastern Pacific Ocean. *Wilson Bulletin* 110:155–170.
- PRATT, H. D., P. L. BRUNER, AND D. G. BERRETT. 1987. A field guide to the birds of Hawaii and the tropical Pacific. Princeton University Press, Princeton, New Jersey.
- RAFFAELE, H., J. WILEY, O. GARRIDO, A. KEITH, AND J. RAFFAELE. 1998. A guide to the birds of the West Indies. Princeton University Press, Princeton, New Jersey.
- RESTALL, R. 1995. Proposed additions to the genus *Lonchura* (Estrildinae). *Bulletin of the British Ornithologists' Club* 115:140–157.
- RESTALL, R. 1996. Munias and mannikins. Yale University Press, New Haven, Connecticut.
- ROBBINS, M. B., AND F. G. STILES. 1999. A new species of pygmy-owl (Strigidae: *Glaucidium*) from the Pacific slope of the northern Andes. *Auk* 116:305–315.
- ROBERSON, D. 1998. Sulids unmasked: Which large booby reaches California? *Field Notes* 52:276–297.
- SEUTIN, G. 1998. Two bird species new for Panama and Central America: White-whiskered Hermit *Phaethornis yaruqui* and White-winged Swallow *Tachycineta albiventer*. *Cotinga* 9:22–23.
- SHEDD, D. H., R. D. GETTINGER, B. L. SHEDD, AND F. R. SCOTT. 1998. First record of a Western Marsh Harrier (*Circus aeruginosus*) [sic] in Virginia. *Raven* 69:56.
- SHORT, L. L., JR. 1982. Woodpeckers of the world. Delaware Museum of Natural History, Greenville.
- SIBLEY, C. G., AND B. L. MONROE, JR. 1990. Distribution and taxonomy of birds of the world. Yale University Press, New Haven, Connecticut.
- SIBLEY, C. G., AND B. L. MONROE, JR. 1993. A supplement to distribution and taxonomy of birds of the world. Yale University Press, New Haven, Connecticut.
- SYKES, P. W., JR. 1998. Yellow-throated Bunting at Attu. *Field Notes* 52:398–403.
- TYRBERG, T. 1998. The date of publication of Montin's description of *Lagopus mutus*. *Bulletin of the British Ornithologists' Club* 118:56–57.
- YOUNG, J. R., C. E. BRAUN, S. J. OYLER-MCCANCE, T. W. QUINN, AND J. W. HUPP. 2000. A new species of Sage Grouse (Phasianidae: *Centrocercus*) from southwestern Colorado, USA. *Wilson Bulletin* 112: in press.
- YOUNG, J. R., J. W. HUPP, J. W. BRADBURY, AND C. E. BRAUN. 1994. Phenotypic divergence of secondary sexual traits among Sage Grouse, *Centrocercus urophasianus*, populations. *Animal Behaviour* 47:1353–1362.
- ZINK, R. M., S. ROHWER, A. V. ANDREEV, AND D. L. DITTMANN. 1995. Trans-Beringia comparisons of mitochondrial DNA differentiation in birds. *Condor* 97:639–649.
- Committee:*  
 RICHARD C. BANKS, *Chairman*  
 CARLA CICERO  
 JON L. DUNN  
 ANDREW W. KRATTER  
 HENRI OUELLET (deceased)  
 PAMELA C. RASMUSSEN  
 J. V. REMSEN, JR.  
 JAMES A. RISING  
 DOUGLAS F. STOTZ
- Preferred citation: American Ornithologists' Union. 2000. Forty-second supplement to the American Ornithologists' Union *Check-list of North American Birds*. *Auk* 117:847–858.