

## **Altitudinal bird migration in North America**

Author: Boyle, W. Alice

Source: The Auk, 134(2) : 443-465

Published By: American Ornithological Society

URL: <https://doi.org/10.1642/AUK-16-228.1>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



REVIEW

## Altitudinal bird migration in North America

W. Alice Boyle

Division of Biology, 116 Ackert Hall, Kansas State University, Manhattan, Kansas, USA  
[aboyle@ksu.edu](mailto:aboyle@ksu.edu)

Submitted November 4, 2016; Accepted January 18, 2017; Published March 29, 2017

### ABSTRACT

Altitudinal bird migration involves annual seasonal movements up and down elevational gradients. Despite the fact that species from montane avifaunas worldwide engage in altitudinal migration, the patterns, causes, and prevalence of these movements are poorly understood. This is particularly true in North America where the overwhelming majority of avian migration research has focused on obligate, long-distance, temperate-tropical movements. Elsewhere in the world, most altitudinal migrants are partial migrants, making downhill movements to nonbreeding areas. However, spatial and temporal patterns, the prevalence and predictability of migration at individual and population levels, and the ultimate ecological factors selecting for movement behavior vary considerably among taxa and regions. I conducted a systematic survey of the evidence for altitudinal migration to fill gaps in our understanding of this behavior among the landbirds of North America and Hawaii. Altitudinal migration was as prevalent as in other avifaunas, occurring in >20% of continental North American and nearly 30% of Hawaiian species. Of the species wintering within the USA and Canada, ~30% engage in altitudinal migrations. Altitudinal migrants are far more common in the West, are taxonomically and ecologically diverse, and North American species exhibit patterns similar to altitudinal migrants elsewhere in the world. Because altitudinal migration systems are relatively tractable, they present excellent opportunities for testing hypotheses regarding migration generally. Altitudinal migration has likely been overlooked in North America due to contingency in the history of ornithological research. Our need to understand the patterns and causes of altitudinal migrations has never been greater due to emerging environmental threats to montane systems.

*Keywords:* alpine, altitudinal movements, elevational migration, facultative migration, high elevation, partial migration, post-breeding dispersal, nomadism

### Migración altitudinal de aves en América del Norte

La migración altitudinal refiere a movimientos estacionales a través de gradientes elevacionales. Aunque aves de altura por todo el mundo hacen migraciones altitudinales, los patrones, causas, y la prevalencia son poco conocidas comparando con la migración latitudinal de larga distancia. Se desconoce los patrones en particular para la avifauna de Norteamérica, donde la gran mayoría de investigaciones han enfocado en movimientos entre zonas templadas y tropicales. En otras partes del mundo, migraciones altitudinales tienden de ser parciales, y las aves bajan desde zonas altas donde anidan hacia zonas bajas donde pasan la época non-reproductiva. Sin embargo, existe mucha variación en los patrones espaciales y temporales, la prevalencia y previsibilidad de migración al nivel individual y poblacional, y los factores ecológicos que afecta a este comportamiento. Realicé una encuesta sistemática de la evidencia por la migración altitudinal en las aves terrestres de Norteamérica y Hawaii. La prevalencia de migración altitudinal era parecido con otros avifaunas, ocurriendo en más de 20% de las especies continentales y 30% de las especies de Hawái. De las especies que invernán completamente o parcialmente en los EEUU o Canadá, ~30% migran altitudinalmente. Los migrantes son mucho más común en el oeste, son diversos en cuanto la taxonomía y ecología, y muestran patrones similares a migrantes altitudinales en otras partes del mundo. Como son relativamente fáciles de estudiar, son sistemas excelentes para probar hipótesis para la migración en general. Lo más posible es que la migración altitudinal se ha pasado por alto por azar o por idiosincrasias de la historia de investigaciones ornitológicas. Nos urge entender los patrones y causas de migración altitudinal en Norteamérica como en otras avifaunas frente los cambios ambientales que están amenazando a zonas de altura mundiales.

*Palabras clave:* alpina, movimiento altitudinales, movimiento elevacional, migración facultativa, migración parcial, dispersión, nomadismo

## INTRODUCTION

Birds are the most mobile of animal taxa. Their mobility has profoundly influenced every other aspect of their ecology and evolution. To paraphrase Dobzhansky's (1973) famous quote, nothing in ornithology makes sense except in light of movement strategies. The study of avian movements has focused on migration, a term which (in ornithological literature) refers to predictable, annual, return movements between breeding and nonbreeding ranges. Classic or "regular" bird migration involves long-distance movements of all individuals in a population between discrete breeding and wintering areas (Terrill and Able 1988) via movements that are often innately controlled (Berthold 1991). The overwhelming majority of avian migration studies have focused on obligate, innate, long-distance movements, from macro-ecological scales (e.g., Dingle 2008, Somveille et al. 2013) to field-intensive, population studies (e.g., Marra et al. 1998, Ydenberg et al. 2002, Stanley et al. 2012). These and hundreds of other studies have taught us a great deal about the implications of being migratory, especially for physiology, demography, and conservation.

While long-distance, obligate migrations may be "classic" examples, they may not be representative or even typical. Most species of avian migrants are probably partial migrants rather than complete migrants (Berthold 2001) meaning that not all individuals migrate. Differences in migratory tendency can occur at the population level (Berthold 1998, Jahn et al. 2012) or at the individual level (Ogonowski and Conway 2009, Hegemann et al. 2015). Many species exhibiting behavior at this "fuzzier" end of the spectrum of migratory behavior also migrate facultatively over shorter distances and exhibit less predictability in tendency, timing, and distance (Newton 2012). One such type of migratory behavior involves movements over elevational gradients. These *altitudinal migrations* often involve annual, return movements between breeding and nonbreeding ranges, justifying application of the term migration in an ornithological context. They are poorly understood, particularly within temperate avifaunas.

Ignoring altitudinal migrants and other facultative, short-distance movements has several consequences. First, our world view of migration is inherently biased, meaning that scientific perceptions of "normal" are flawed. Second, we miss the opportunity to learn a lot about the selective pressures that result in migration by ignoring comparatively tractable systems. Third, an incomplete understanding of avian movement interferes with our ability to protect species. This review first describes major patterns and the prevalence of altitudinal migration in birds globally, briefly outlines explanations and evidence for altitudinal migrations, then explores the patterns and


prevalence of altitudinal migration within the North American avifauna via a systematic survey.

### The What, Where, and How of Altitudinal Migration

Unlike obligate, latitudinal migrations, altitudinal migrations typically (1) involve short distances, (2) are controlled facultatively, and often (3) consist of partially migratory populations. These attributes make biological sense; ecological gradients change quickly over short distances with elevational change (Körner 2007), meaning that birds may not need to move as far as do latitudinal migrants to experience similar fitness payoffs from migration (e.g., escaping unfavorable climatic conditions, exploiting food resources in areas with different phenological patterns). Lower movement costs likely reduces selection for morphological and physiological adaptations required to complete those movements. Thus, the expression of migration can be more flexible, varying among individuals (e.g., Gillis et al. 2008) or within individuals under different circumstances during their lifetimes (Norbu et al. 2013).

Altitudinal migrant birds live in all major mountain ranges of the Earth (Figure 1). Asia may be home to the greatest number of altitudinal migrants in the world where, purportedly, up to 65% of high-elevation–breeding species in the Himalayas engage in such movements. Altitudinal migrants are distributed throughout the avian tree of life with representatives in orders as morphologically and ecologically distinct as the Charadriiformes (Ferrari et al. 2008) and Psittaciformes (Chassot and Monge-Arias 2012). Altitudinal migrants also represent a broad range of foraging guilds. Neotropical nectarivores and frugivores are well-known altitudinal migrants (Stiles 1980, Strewé 2006), but nectarivores elsewhere (Grant and Grant 1967), insectivores (Nocedal 1994), carnivores (Bildstein 2004), piscivores (Mackas et al. 2010), granivores (Horvath and Sullivan 1988, Borrás et al. 2010), and birds of other dietary guilds (e.g., Martínez del Río et al. 2009) also migrate altitudinally. They range in body size from under a few grams to several kilograms (Boyle 2011). Altitudinal migrants live in forests, deserts, shrublands, and terrestrial aquatic environments. They dwell at all levels from the ground (e.g., Dark-eyed Juncos; Rabenold and Rabenold 1985) up to the canopy (e.g., Great Green Macaws; Chassot and Monge-Arias 2012), are flocking (e.g., Mountain Quail; Brennan et al. 1987) or solitary (e.g., Andean Condors; Sick 1993), and represent diverse mating systems.

Although species differ in the elevations they occupy, many migrants breed at higher elevations than where they spend their nonbreeding season in both tropical and temperate regions (Dixon and Gilbert 1964, Ramos-Olmos 1983, Norment and Green 2004, Powell and Bjork 2004). Many exceptions do occur, however. For example, the endangered Nēnē or Hawaiian Goose (*Branta sandvicensis*)



**FIGURE 1.** Altitudinal bird migration around the world. Numbers in red ovals correspond to references listed below documenting altitudinal migration in most major mountain ranges globally. Tallies of community- or guild-level prevalence of altitudinal migration (reported in dark boxes) provide a comparison to the results of this study in North America (inset). [1] Rabenold and Rabenold 1985; [2–4] Cade and Hoffman 1993, Connelly 1988, Dixon and Gilbert 1964; [5, 6] Dunning and Bowers 1984, Horvath and Sullivan 1988; [7–11] Garwood et al. 2009, Gillis et al. 2008, Grinnell and Miller 1944, Laymon 1989, Lorenz and Sullivan 2009; [12, 13] Des Granges 1979, Solórzano et al. 2000; [14–20] Boyle 2011, Chassot and Monge-Arias 2012, Chaves-Campos 2004, Chaves-Campos et al. 2003, Fraser et al. 2008, Loiselle and Blake 1991, Stiles 1980; [21–27] Newsome et al. 2015, Tinoco et al. 2009, Strewe and Navarro 2003, Hobson et al. 2003, Hilty 1997, Hughes 1980, 1984; [28] Beebe 1947; [29–31] Galetti 2001, Bencke and Kindel 1999, Areta and Bodrati 2010; [28, 32–34] Beebe 1947, Bildstein 2004, Alves 2007, Capllonch 2007; [35, 36] Borrás et al. 2010, Klemp 2003; [37, 38] Brambilla and Rubolini 2009, De La Hera et al. 2014; [39] Burgess and Mlingwa 2000; [40, 41] Brown 2006, Johnson and Maclean 1994; [42] Lu et al. 2010; [43, 44] Kimura et al. 2001, Ryan 2012; [45, 46] Dingle 2004, 2008; [47] Nosedal 1994; [48] Stiles and Skutch 1989; [32] Bildstein 2004; [39] Burgess and Mlingwa 2000; [41] Johnson and Maclean 1994; [49] Barve et al. 2016; [45] Dingle 2004; [50] Tidemann et al. 1988; [51] Dixit et al. 2016.

breeds and molts at low elevations, migrating to high elevations during the nonbreeding season (Hess et al. 2012). A few species apparently make both uphill and downhill movements post-breeding; in the New England region of New South Wales, Australia, Eastern Spinebills (*Acanthorhynchus tenuirostris*) migrate uphill in winter (Ford and Pursey 1982), whereas farther south in the same mountain range, this species migrates to lower elevations in winter (Tidemann et al. 1988). In partially migratory populations, the elevation where nonmigrant individuals reside year-round can be at either the upper or lower end of the elevational gradient. In species where nonmigrants spend the nonbreeding season at high elevation, the migrants and residents generally breed together but spend the nonbreeding season apart (e.g., Resplendent Quetzals; Solórzano et al. 2000). In contrast, where nonmigrants spend the nonbreeding season at low elevations, then

migrants and residents breed separately, meaning that the potential for assortative mating is strong (e.g., American Dippers; Middleton et al. 2006).

We know little regarding how birds make altitudinal migratory movements. Mountain Quail apparently traverse >2,000 m elevational gradients primarily on foot (Gutiérrez and Delehanty 1999). Whether small songbirds make altitudinal migrations in nocturnal flights as do many of their long-distance counterparts is unknown. Likewise, for most species, we do not know whether migrations occur in a single directed flight or many short flights. In Central American Three-wattled Bellbirds (*Procnias tricaunculata*), radio-tagged birds typically departed their high-elevation, Pacific-slope breeding areas in a single sustained flight to the Caribbean lowlands. A few months later, they recrossed the Continental Divide in multiple flights to the Pacific lowlands before returning to

high-elevation breeding areas (Powell and Bjork 2004). Satellite-tagged Nēnē exhibit considerable inter-individual flexibility in the duration and distance of migratory journeys, with some individuals making nonstop flights while others make multiple stopovers (Leopold and Hess 2014). In some tropical passerines, it seems likely that migratory flights can be fast and directed given the rapid increase in capture rates of altitudinal migrants following storm events (Ramos-Olmos 1983, Boyle et al. 2010). In this way, some altitudinal migrations parallel the unpredictable, short-term elevational movements of birds escaping from inclement weather (Hahn et al. 2004, O'Neill and Parker 1978); a key distinction is that, once in the lowlands, tropical altitudinal migrants remain for the rest of the nonbreeding season.

### Why Do Birds Migrate Altitudinally?

The causes of altitudinal migration have only been studied in a few systems and species. Soon after altitudinal migration was described for tropical birds (Beebe 1947, Slud 1964), the idea that such movements were driven by spatial and temporal variation in food resources gained traction (Stiles 1980, 1988; Wheelwright 1983). This hypothesis made strong intuitive sense, especially given broad species-level associations between short-distance intra-tropical migration and dependence on nectar or fruit (Levey and Stiles 1992). Several studies tested predictions of this hypothesis to explain community-level (Loiselle and Blake 1991) and species-level patterns of abundance and movement (e.g., Rosselli 1994, Chaves-Campos 2004). Frequently, while part of the annual movement cycle was consistent with birds seeking elevations with high food abundance, the timing and direction of return movements did not match food-based predictions. For example, Bare-necked Umbrellabirds (*Cephalopterus glabricollis*) bred in locations and seasons of high food abundance, but migrated downhill to elevations having lower food availability (Chaves-Campos et al. 2003). Species-pairs of frugivorous migrants and nonmigrants revealed that neither dietary specialization nor competitive asymmetries could explain differences in migration, but even within genera, migrants consumed more fruit relative to arthropod prey (Boyle et al. 2011a). This finding suggested that energetic constraints driven by dependence on nutritionally dilute foods may underlie migratory tendency in some tropical species.

Evidence exists for alternative processes driving altitudinal migration including predation risk (Fretwell 1980, Greenberg 1980) and climatic constraints (Cox 1985). For instance, differences in nest predation risk could potentially explain the movements of birds between middle and high elevations (Boyle 2008a). However, in many species, the temporal and spatial patterns predicted by multiple ecological processes overlap (Ornelas and Arizmendi

1995). Climatic constraints have received relatively little attention in tropical regions despite Skutch (1969: 85) writing, "The Bellbird's wanderings appear to be caused partly by changes in weather..." However, Nosedal (1994) conjectured that in western Mexico, interannual variation in migratory propensity may well reflect individual differences in climatic tolerances and interannual variation in weather.

A mechanistic link between dietary guild, climate, and migration exists; dependence on low-protein, low-fat, dilute foods means that frugivores and nectarivores must forage frequently. Intense rain storms or cooler weather that increase thermogenic costs will affect small birds most acutely as they have narrow energetic safety margins and require high food intake rates. Storms can potentially limit the number of foraging hours to such an extent that such birds with high metabolic rates risk starvation, not because there is not enough food, but because the climate constrains their ability to forage (Boyle 2008b). Evidence in support of this hypothesis comes from Costa Rican White-ruffed Manakins (*Corapipo altera*); heavy rainfall is perceived by birds as a stressor and leads to increased fuel storage and dependence on stored energy for survival (Boyle et al. 2010). Additionally, in a diverse set of species, more altitudinal migrants reach the lowlands in years with wetter wet seasons (Boyle 2011). Thus, it seems likely that, at least in some tropical systems, the relationship between diet and migratory strategy are not driven by simple food availability, but via interactions with climate and energetics to shape individual and species-level migratory behavior.

In temperate North America, interest in altitudinal migration dates back farther than in tropical systems (e.g., Presnall 1935, Grinnell and Miller 1944, Dixon and Gilbert 1964). Climate has always made intuitive sense in explaining why birds might "escape" from temperate high elevations during winter. However, for most dietary guilds, seasons of climatic harshness coincide with food scarcity, and few authors have attempted to determine whether direct physiological challenges imposed by weather or insufficient food is more important in shaping movement. Carolina Dark-eyed Juncos engage in female-biased downhill migration away from high-elevation breeding areas (Rabenold and Rabenold 1985), as would be predicted by the *dominance*, *body-size*, and *arrival time* hypotheses proposed to explain differential and partial latitudinal migration (Ketterson and Nolan 1983). Inter-annual variation in the identity and abundance of high-elevation residents implicates climate in mediating some costs of residency and that, likely, competition for scarce winter food explains why many individuals depart breeding grounds. In other well-studied temperate systems, competition also appears to be important in shaping migratory strategies. American Dippers breed at both high and low elevations in southwestern Canada and winter together

along low-elevation rivers. Snow and ice restricts access to food during winter at high elevations. Because fecundity is lower in migrants, and survival is only a little higher, the benefits of breeding at high elevation are less clear (Green et al. 2015) and, possibly, altitudinal migration in this system is driven by intraspecific competition for food-rich low-elevation territories. In contrast, Yellow-eyed Juncos (*Junco phaeonotus*) breeding at high elevations of Southern Arizona's sky islands appear to be limited in their ability to remain at high elevations year-round due to direct and indirect energetic consequences of cold temperatures and snow limiting foraging opportunities during winter (Lundblad 2014).

Generalization regarding the causes of altitudinal migration within major biogeographic regions or among them is extremely difficult due to the few species that have been the subject of detailed investigations. Given the diversity of patterns exhibited by altitudinal migrants, multiple ecological processes almost certainly shape this behavior. One important emergent conclusion is that altitudinal migrations are often facultative, at least within the Americas. Thus, the fitness costs and benefits of alternate strategies can be interpreted in the context of current ecological drivers, making them excellent systems in which to understand the evolutionary pressures that have shaped less labile movement strategies.

### ALTITUDINAL MIGRATION: A GAP IN NORTH AMERICAN ORNITHOLOGY?

Perhaps because they represent less spectacular feats of navigation and endurance, altitudinal migration is understudied in North American ornithology. In a recent search of the ISI Web of Knowledge database, the number of articles (all years, all journals) returned for "altitudinal migration" or "elevational migration" in combination with "bird" resulted in only 45 studies. Recent volumes (Jan 2011–Aug 2016) of *The Auk: Ornithological Advances* and *The Condor: Ornithological Applications* tell a similar story. A search on the word "migration" within abstracts resulted in 80 and 82 articles, respectively. Forty (*Auk*) and 45 (*Condor*) of those studies focused on topics relating to annual, return movements of birds, and all focused on long-distance latitudinal migration. This bias is reflected in pedagogical materials; in a commonly used 750-page ornithology textbook, altitudinal migration was mentioned in only 2 sentences that referred to tropical frugivores and nectarivores wandering unpredictably in search of food resources (Gill 2007).

Why do we know so little about altitudinal and other short-distance migrations of North American birds? One possibility is that there are few of them in our avifauna. Therefore, focusing on such movements lacks local relevance. An alternative is that understanding such

movements does not advance fundamental understanding of animal movement generally, and therefore the topic has been ignored for a reason. A third hypothesis is that we have overlooked these movements due to artifacts of the cultural history of ornithology in North America—because of geography or random chance, research traditions have taken the discipline elsewhere. I reviewed the prevalence and patterns of altitudinal migration in North American birds, then used those results to frame an evaluation of these 3 hypotheses.

### A Systematic Survey

I systematically assessed the prevalence of altitudinal migration in the landbirds breeding in North America north of the Mexico border (i.e. Canada and the USA) and Hawaii. I used the *Birds of North America* (Rodewald 2015) as it represents the most comprehensive set of life histories available for this avifauna presented in a common format. I accessed this source and tabulated data during the summer and fall of 2014. I included all 607 extant, native landbirds breeding in this region, excluding the orders Pelecaniformes, Procellariiformes, Gaviiformes, Podicipediformes, Phaethontiformes, Suliformes, and the families Stercorariidae and Alcidae as these groups are almost exclusively aquatic with limited opportunities for altitudinal movements. Nomenclature and species concepts follow the *American Ornithologists' Union Check-list* and supplements (American Ornithologists' Union 1998, Chesser et al. 2013).

I read the Introduction, Migration, Distribution, and Habitat sections for each species. I tabulated information for all species for which any populations were mentioned migrating altitudinally, referencing original sources. Use of the word "migration" varied among authors of life history accounts. Accordingly, I included species referenced as undertaking "elevational movements," "downhill dispersal," or other similar terms when they referred to seasonal movements between breeding and nonbreeding areas. I coded the strength of evidence qualitatively. I considered evidence to be "strong" when accounts reported results of movement-focused studies or if reports from multiple populations reported altitudinal migration with no ambiguity in the language describing altitudinal movements. I considered the strength of evidence to be "intermediate" when references consisted of more anecdotal information, but different authors working in different populations reported such movements and authors did not use ambiguous wording. I considered the evidence to be "weak" if movement descriptions were accompanied by words denoting uncertainty (e.g., "In montane areas, Pileated Woodpeckers *may* move to lower elevations in winter (Burleigh 1972; Simpson and Pratt 1992)"). I coded each putative altitudinal migrant species as completely or partially migratory, and if partially migrant, whether that

variation occurred among or within populations. For partially migratory populations, I coded individuals as either breeding or wintering together. I noted species that included latitudinal migrant populations, and whether any population overwinters within the continental USA and Canada. I coded species for geographic overlap in breeding and wintering range and, where present, whether that overlap occurred within the continental USA and Canada.

I classified species into one of eight broad geographic distributions: “western” and “eastern” for species whose ranges primarily lie west or east of the 100th meridian, “widespread” for species occurring commonly in both eastern and western regions, “northern” for species occurring wholly or primarily in boreal or more northerly regions, “southwestern” for species only occurring in North America in deserts or sky islands of the Southwest or extreme southern Texas, “Hawaii” for species not found elsewhere in North America, and “other” for species not easily classifiable into any of these categories. I excluded extinct or introduced birds in the summaries below, but read and tabulated data for such species’ accounts when they were available. Although there were few missing species accounts, life histories varied in the date of initial publication and subsequent revision, and it is possible that new studies may have been published since tabulation. Given constraints of study design and the relative paucity of studies on altitudinal migration in North America, the estimates presented below should be viewed as extremely conservative minima.

## RESULTS

Of 607 native, extant breeding landbirds covered in the *Birds of North America* series, accounts for 163 species (26.8%) reported altitudinal migration in at least one population somewhere in their range (Table 1). Of those, 10 species only occur in Hawaii; altitudinal migrants represent 28.6% of the 32 extant landbird species restricted to Hawaii. Ninety-eight (64.1%) of the 153 continental altitudinal migrants belong to species that also engage in latitudinal migrations. The majority (111 species; 72.5%) of continental altitudinal migrants were partially migratory at the population level (87 species), the individual level (51 species), or both (27 species). In species where migratory behavior varies among individuals, 70.1% of species breed together and spend the nonbreeding season at different elevations (e.g., Lesser Goldfinch, *Spinus psaltria*). Likewise, 75.2% of continental partial migrants breed at higher elevations than where they winter.

One hundred and sixteen continental species undertake altitudinal migrations within the USA or Canada (75.8% of continental altitudinal migrants or 20.2% of all continental species in the dataset). This represents 30.3% of the 383 continental landbirds that overwinter within the USA and

Canada. Evidence for altitudinal migration was strong for 45 species, intermediate for 45 species, and weak for 26 species. One hundred and twelve (36.3%) of the 309 species with overlapping North American breeding and wintering distributions migrate altitudinally. The West was home to half (58) of the species that engage in altitudinal migrations within North America. An additional 40 altitudinal migrant species were distributed on both sides of the 100th meridian. Thirteen of the 68 southwestern species (19.1%) contained references documenting altitudinal migration within the region. Only one species coded as eastern (Northern Bobwhite, *Colinus virginianus*; Rosene 1969) was noted undertaking seasonal altitudinal migrations.

Within orders and families, the prevalence of altitudinal migration varied considerably (Table 2). Only 5 orders contained no altitudinal migrants, and these all are represented in North America by  $\leq 6$  breeding species. In 6 orders, and 8 families within Passeriformes,  $>25\%$  of species migrate altitudinally somewhere within their range (Table 2).

Several species displayed complex movement patterns that challenged categorization. For instance, Black Rosy-Finches (*Leucosticte atrata*) generally winter at lower elevations, but can make multiple uphill and downhill movements at any time of year, responding to short-term weather events (King and Wales 1964, Hendricks and Swenson 1983, Johnson 2002). In Arizona, Phainopeplas (*Phainopepla nitens*) breed in low-elevation Sonoran deserts in February–April, then move upslope, breeding again in oak–sycamore canyons in May–July, then move even higher during August–September (Swarth 1904, Crouch 1943, Walsberg 1977). In October, Phainopeplas migrate back down to the lowest elevations for the remainder of the nonbreeding season, and where they can be quite mobile (Chu and Walsberg 1999). Uphill post-breeding movements to higher elevations, followed by downhill movements to wintering areas below the breeding elevation, was described for several species including Greater Sage-Grouse (Schroeder et al. 1999), Prairie Falcon (Steenhof 2013), Anna’s Hummingbird (Clark and Russell 2012), Clark’s Nutcracker (Tomback 1998), and Orange-crowned Warbler (Gilbert et al. 2010). Individuals and populations of many other species included in Table 1 combine both latitudinal and altitudinal movements, a pattern common to altitudinal migrant bats (McGuire and Boyle 2013). For example, Golden-crowned Kinglets exhibit “downslope altitudinal migration in winter in western mountains” (Swanson et al. 2012). However, what is not clear is the extent to which high-elevation breeders also shift south like their boreal-breeding conspecifics.

Species accounts for some extinct species referenced altitudinal migration including the Hawaiian ‘Ō‘ō (*Moho*

**TABLE 1.** All 163 extant species described as making altitudinal migrations or whose life history accounts suggested they make regular movements along elevational gradients anywhere within their range. See text for basis of categorical assignment of the strength of evidence (weak, intermediate, strong) and region (eastern, western, widespread, northern, southwest, Hawaii, other). Additionally, I coded each species based on whether or not descriptions of altitudinal migrations pertained to movements made within North America (i.e. continental USA and Canada; "Alt mig in NA?"), whether or not the species was partially migratory and, if so, whether the variation in migratory behavior partitioned at individual and/or population levels ("Partially migratory?"), whether altitudinal migrations involved downward movements from higher elevation breeding areas or vice versa ("Breeds high?"), and whether (in partially migratory populations) migrants and residents breed at the same elevations ("Breeds together?"). Original sources for migration descriptions referenced in the *Birds of North America* accounts are listed by number below the table.

Family	Scientific name	Common name	Evidence	Alt mig in NA?	Region	Partial migrant?	Breeds high?	Breeds together?	References	
Anatidae	<i>Branta sandvicensis</i>	Hawaiian Goose	strong		Hawaii	individ		x	[1–7]	
	<i>Cygnus buccinator</i>	Trumpeter Swan	intermediate	x	widespread	both	x	x	[8]	
	<i>Anas wyvilliana</i>	Hawaiian Duck	weak		Hawaii	individ	x	x	[9]	
	<i>Anas cyanoptera</i>	Cinnamon Teal	weak		western				[10]	
	<i>Histrionicus histrionicus</i>	Harlequin Duck	intermediate	x	widespread	individ	x	x	[11, 12]	
	<i>Bucephala albeola</i>	Bufflehead	strong	x	widespread		x		[13]	
	<i>Bucephala islandica</i>	Barrow's Goldeneye	strong	x	widespread		x		[14]	
	<i>Lophodytes cucullatus</i>	Hooded Merganser	weak	x	widespread		x		[14]	
	<i>Mergus merganser</i>	Common Merganser	weak	x	widespread		x		[15, 16]	
	<i>Oxyura jamaicensis</i>	Ruddy Duck	intermediate	x	widespread		x		[17]	
	Odontophoridae	<i>Oreortyx pictus</i>	Mountain Quail	strong	x	western		x		[18–24]
		<i>Callipepla californica</i>	California Quail	weak	x	western	individ	x		[8]
		<i>Callipepla gambelii</i>	Gambel's Quail	weak	x	western	individ	x		[25, 26]
<i>Colinus virginianus</i>		Northern Bobwhite	intermediate	x	eastern	pop	x		[27]	
Phasianidae	<i>Centrocercus urophasianus</i>	Greater Sage-Grouse	strong	x	western	both		x	[28–35]	
	<i>Falcipternis canadensis</i>	Spruce Grouse	strong	x	widespread	individ			[36–38]	
	<i>Lagopus lagopus</i>	Willow Ptarmigan	strong	x	northern	both	x	x	[39–42]	
	<i>Lagopus leucura</i>	White-tailed Ptarmigan	strong	x	western	both	x	x	[36, 43–47]	
	<i>Dendragapus obscurus</i>	Dusky Grouse	strong	x	western	both	x	x	[48, 49]	
	<i>Dendragapus fuliginosus</i>	Sooty Grouse	strong	x	western	both	x	x	[48, 49]	
	<i>Tympanuchus phasianellus</i>	Sharp-tailed Grouse	intermediate	x	western	both	x	x	[50–52]	
Cathartidae	<i>Coragyps atratus</i>	Black Vulture	weak	x	widespread	both	x		[53, 54]	
Accipitridae	<i>Accipiter gentilis</i>	Northern Goshawk	weak	x	widespread				[8]	
	<i>Buteo lineatus</i>	Red-shouldered Hawk	intermediate	x	widespread		x		[55]	
Falconidae	<i>Buteo albonotatus</i>	Zone-tailed Hawk	intermediate		western	both	x	x	[56]	
	<i>Falco mexicanus</i>	Prairie Falcon	strong	x	western	both			[57]	
Rallidae	<i>Gallinula galeata</i>	Common Gallinule	intermediate	x	widespread	pop	x		[58]	
	<i>Fulica americana</i>	American Coot	intermediate	x	widespread	pop	x		[59, 60]	
Scolopacidae	<i>Calidris virgata</i>	Surfbird	strong	x	northern	pop	x		[8]	
	<i>Calidris ptilocnemis</i>	Rock Sandpiper	strong	x	northern	pop	x		[8]	



TABLE 1. Continued.

Family	Scientific name	Common name	Evidence	Alt mig in NA?	Region	Partial migrant?	Breeds high?	Breeds together?	References	
Laridae	<i>Pagophila eburnea</i>	Ivory Gull	strong		northern	pop	x		[61, 62]	
	<i>Larus californicus</i>	California Gull	strong	x	western	pop	x		[8, 63]	
Columbidae	<i>Patagioenas leucocephala</i>	White-crowned Pigeon	strong		other	both		x	[64]	
	<i>Patagioenas fasciata</i>	Band-tailed Pigeon	weak	x	western	both	x	x	[8]	
	<i>Columbina passerina</i>	Common Ground-Dove	weak		other	both	x	x	[8]	
	<i>Megascops trichopsis</i>	Whiskered Screech-Owl	weak	x	southwest	individ	x		[8]	
Strigidae	<i>Glaucidium gnoma</i>	Northern Pygmy-Owl	strong	x	western	x	x		[65–70]	
	<i>Strix occidentalis</i>	Spotted Owl	strong	x	western	both	x	x	[71–76]	
	<i>Strix nebulosa</i>	Great Gray Owl	intermediate	x	widespread	pop	x		[77–79]	
	<i>Aegolius acadicus</i>	Northern Saw-whet Owl	strong	x	widespread	pop	x		[80–82]	
	<i>Phalaenoptilus nuttallii</i>	Common Poorwill	intermediate	x	western	both	x		[83]	
	<i>Anrostomus ridgwayi</i>	Buff-collared Nightjar	weak		southwest	pop	x		[56]	
Apodidae	<i>Anrostomus arizonae</i>	Mexican Whip-poor-will	weak		southwest	pop	x		[84]	
	<i>Aeronautes saxatalis</i>	White-throated Swift	intermediate	x	western		x		[17, 82, 84–86]	
Trochilidae	<i>Lampornis clemenciae</i>	Blue-throated Hummingbird	intermediate		southwest	pop	x		[87]	
	<i>Eugenes fulgens</i>	Magnificent Hummingbird	strong		southwest		x		[88–91]	
	<i>Calothorax lucifer</i>	Lucifer Hummingbird	weak		southwest				[92, 93]	
	<i>Archilochus alexandri</i>	Black-chinned Hummingbird	weak	x	western	individ		x	[8]	
	<i>Calypte anna</i>	Anna's Hummingbird	strong	x	western	both		x	[94–99]	
	<i>Calypte costae</i>	Costa's Hummingbird	weak		western	pop			[100–102]	
	<i>Selasphorus rufus</i>	Rufous Hummingbird	intermediate		western					
	<i>Melanerpes lewis</i>	Lewis's Woodpecker	intermediate	x	western				[17, 103–106]	
	<i>Melanerpes uropygialis</i>	Gila Woodpecker	intermediate	x	southwest	both		x	[107, 108]	
	<i>Sphyrapicus thyroideus</i>	Williamson's Sapsucker	intermediate	x	western	pop	x		[108, 109]	
Picidae	<i>Sphyrapicus nuchalis</i>	Red-naped Sapsucker	intermediate	x	western	both			[109]	
	<i>Sphyrapicus ruber</i>	Red-breasted Sapsucker	strong	x	western		x		[70, 110, 111]	
	<i>Picoides nuttallii</i>	Nuttall's Woodpecker	intermediate	x	western	individ		x	[17]	
	<i>Picoides villosus</i>	Hairy Woodpecker	intermediate	x	widespread	individ	x	x	[17, 70]	
	<i>Picoides arizonae</i>	Arizona Woodpecker	intermediate	x	southwest	individ	x	x	[86, 108]	
	<i>Picoides albolarvatus</i>	White-headed Woodpecker	weak	x	western	individ	x		[112, 113]	
	<i>Colaptes auratus</i>	Northern Flicker	intermediate	x	widespread		x		[17, 114, 115]	
	<i>Dryocopus pileatus</i>	Pileated Woodpecker	weak	x	widespread	individ	x	x	[116, 117]	
	Tyrannidae	<i>Campostoma imberbe</i>	Northern Beardless-Tyrannulet	weak	x	southwest	individ	x		[108]
		<i>Contopus pertinax</i>	Greater Pewee	strong		southwest	pop	x		[84, 118–122]
<i>Empidonax difficilis</i>		Pacific-slope Flycatcher	weak		western		x		[8]	
<i>Empidonax occidentalis</i>		Cordilleran Flycatcher	weak		western		x		[8]	
<i>Empidonax fulvifrons</i>		Buff-breasted Flycatcher	intermediate		southwest		x		[8]	

TABLE 1. Continued.

Family	Scientific name	Common name	Evidence	Alt mig in NA?	Region	Partial migrant? high?	Breeds together?	References
Vireonidae	<i>Sayornis nigricans</i>	Black Phoebe	intermediate	x	western	pop		[17, 91]
	<i>Pyrocephalus rubinus</i>	Vermilion Flycatcher	intermediate	x	southwest		x	[8]
	<i>Myiarchus tuberculifer</i>	Dusky-capped Flycatcher	intermediate		western		x	[84]
Corvidae	<i>Vireo plumbeus</i>	Plumbeous Vireo	weak		western		x	[86, 123, 124]
	<i>Vireo huttoni</i>	Hutton's Vireo	intermediate	x	western		x	[94, 108, 125, 126]
	<i>Vireo gilvus</i>	Warbling Vireo	weak		western			[8]
	<i>Vireo altiloquus</i>	Black-whiskered Vireo	weak		other	pop		[127]
Alaudidae	<i>Perisoreus canadensis</i>	Gray Jay	weak	x	widespread	individ	x	[128–130]
	<i>Cyanocitta stelleri</i>	Steller's Jay	strong	x	western	pop	x	[108, 131, 132]
	<i>Aphelocoma californica/ woodhouseii</i>	California/ Woodhouse's Scrub-Jay	strong	x	western		x	[132–136]
	<i>Nucifraga columbiana</i>	Clark's Nutcracker	strong	x	western	pop		[137–143]
	<i>Pica hudsonia</i>	Black-billed Magpie	intermediate	x	western	both	x	[8, 144]
	<i>Corvus brachyrhynchos</i>	American Crow	weak	x	widespread		x	[17, 126, 134, 145]
	<i>Corvus hawaiiensis</i>	Hawaiian Crow	strong	x	Hawaii		x	[2, 146]
Alaudidae	<i>Eremophila alpestris</i>	Horned Lark	intermediate	x	widespread	pop		[147, 148]
	<i>Tachycineta thalassina</i>	Violet-green Swallow	strong		western	pop	x	[147]
Paridae	<i>Stelgidopteryx serripennis</i>	Northern Rough-winged Swallow	weak		widespread	pop		[8]
	<i>Poecile gambeli</i>	Mountain Chickadee	strong	x	western	pop	x	[149, 150]
Aegithalidae	<i>Poecile rufescens</i>	Chestnut-backed Chickadee	intermediate	x	western			[126, 144]
	<i>Baeolophus wollweberi</i>	Bridled Titmouse	intermediate	x	western	pop	x	[108, 151, 152]
Sittidae	<i>Psittiparus minimus</i>	Bush-tit	intermediate	x	western	pop	x	[124, 153, 154]
	<i>Sitta canadensis</i>	Red-breasted Nuthatch	strong	x	widespread	pop	x	[144, 155, 156]
Certhiidae	<i>Sitta carolinensis</i>	White-breasted Nuthatch	weak	x	widespread	individ	x	[82]
	<i>Sitta pygmaea</i>	Pygmy Nuthatch	strong	x	western	pop	x	[123, 126, 157–166]
Troglodytidae	<i>Certhia americana</i>	Brown Creeper	weak	x	widespread	pop		[108, 144, 167]
	<i>Salpinctes obsoletus</i>	Rock Wren	weak	x	western	pop	x	[17, 108, 168]
Cinclidae	<i>Catherpes mexicanus</i>	Canyon Wren	weak	x	western	individ	x	[169]
	<i>Troglodytes aedon</i>	House Wren	weak	x	widespread	pop		[17]
Regulidae	<i>Troglodytes pacificus</i>	Pacific Wren	strong	x	western	both	x	[126, 144, 170]
	<i>Cinclus mexicanus</i>	American Dipper	strong	x	western	individ	x	[171–174]
Turdidae	<i>Regulus satrapa</i>	Golden-crowned Kinglet	strong	x	widespread	pop	x	[144, 175–178]
	<i>Regulus calendula</i>	Ruby-crowned Kinglet	intermediate	x	widespread	pop	x	[144, 177, 178]
Turdidae	<i>Sialia mexicana</i>	Western Bluebird	strong	x	western	pop	x	[102, 179–181]

TABLE 1. Continued.

Family	Scientific name	Common name	Evidence	Alt mig in NA?	Region	Partial migrant?	Breeds high?	Breeds together?	References
	<i>Sialia currucoides</i>	Mountain Bluebird	weak	x	western	pop	x		[8]
	<i>Myadestes townsendi</i>	Townsend's Solitaire	strong	x	western	pop	x		[8]
	<i>Catharus ustulatus</i>	Swainson's Thrush	weak		widespread				[182]
	<i>Catharus guttatus</i>	Hermit Thrush	intermediate	x	widespread	pop	x		[179]
	<i>Turdus migratorius</i>	American Robin	weak	x	widespread	pop	x		[8]
	<i>Ixoreus naevius</i>	Varied Thrush	weak	x	western	pop	x		[8]
Mimidae									
	<i>Toxostoma crissale</i>	Crissal Thrasher	intermediate	x	southwest	pop	x		[98, 183–188]
Motacillidae									
	<i>Anthus rubescens</i>	American Pipit	strong	x	western	pop	x		[84, 144, 189–192]
Ptilonotidae									
	<i>Phainopepla nitens</i>	Phainopepla	strong	x	southwest	pop			[98, 193–196]
Peucedramidae									
	<i>Peucedramus taeniatus</i>	Olive Warbler	intermediate	x	southwest	indiv	x		[86, 197]
Calcaridae									
	<i>Plectrophenax nivalis</i>	Snow Bunting	weak		northern	indiv	x		[198, 199]
Parulidae									
	<i>Protonotaria citrea</i>	Prothonotary Warbler	weak		eastern				[182, 200]
	<i>Oreothlypis celata</i>	Orange-crowned Warbler	intermediate	x	widespread				[8]
	<i>Setophaga pitiayumi</i>	Tropical Parula	intermediate		southwest	pop	x		[84]
	<i>Setophaga caerulescens</i>	Black-throated Blue Warbler	weak		eastern	indiv			[201]
	<i>Setophaga coronata</i>	Yellow-rumped Warbler	weak	x	widespread	pop	x		[202, 203]
	<i>Setophaga graciae</i>	Grace's Warbler	intermediate		southwest	pop	x		[151, 204]
	<i>Setophaga occidentalis</i>	Hermit Warbler	intermediate	x	western				[205]
	<i>Myioborus pictus</i>	Painted Redstart	intermediate	x	southwest	indiv	x		[8]
Emberizidae									
	<i>Pipilo chlorurus</i>	Green-tailed Towhee	strong	x	western		x		[17, 108, 176, 206, 207]
	<i>Pipilo maculatus</i>	Spotted Towhee	intermediate	x	western		x		[108, 124, 156]
	<i>Aimophila ruficeps</i>	Rufous-crowned Sparrow	intermediate	x	western	indiv			[94, 108, 181, 208–211]
	<i>Melospiza fusca</i>	Canyon Towhee	intermediate	x	southwest	pop	x		[134, 206]
	<i>Melospiza crissalis</i>	California Towhee	intermediate	x	western		x		[17, 212, 213]
	<i>Spizella atrogularis</i>	Black-chinned Sparrow	intermediate	x	southwest	both	x		[84, 108, 188, 214, 215]
	<i>Amphispiza bilineata</i>	Black-throated Sparrow	intermediate	x	western	pop	x		[108, 156, 214]
	<i>Artemisospiza nevadensis/belli</i>	Sage Sparrow	strong	x	western		x		[216]
	<i>Calamospiza melanocorys</i>	Lark Bunting	intermediate	x	other				[217, 218]
	<i>Passerculus sandwichensis</i>	Savannah Sparrow	weak	x	widespread	pop	x		[8]
	<i>Passerella iliaca</i>	Fox Sparrow	strong	x	widespread	pop	x		[219, 220]
	<i>Melospiza melodia</i>	Song Sparrow	intermediate	x	widespread	pop	x		[221–223]
	<i>Melospiza lincolni</i>	Lincoln's Sparrow	weak	x	widespread	indiv			[8]
	<i>Zonotrichia leucophrys</i>	White-crowned Sparrow	weak		widespread				[224]
	<i>Zonotrichia atricapilla</i>	Golden-crowned Sparrow	intermediate	x	western				[17, 126, 225, 226]
	<i>Junco hyemalis</i>	Dark-eyed Junco	strong	x	widespread	both	x		[227]
	<i>Junco phaeonotus</i>	Yellow-eyed Junco	strong	x	southwest	both	x		[228, 229]
Cardinalidae									
	<i>Piranga flava</i>	Hepatic Tanager	intermediate		southwest	pop	x		[90, 122, 230–232]

TABLE 1. Continued.

Family	Scientific name	Common name	Evidence	Alt mig in NA?	Region	Partial migrant?	Breeds high?	Breeds together?	References
Icteridae	<i>Piranga ludoviciana</i>	Western Tanager	intermediate		western				[230, 233]
	<i>Cardinalis sinuatus</i>	Pyrrhuloxia	weak	x	southwest	pop	x		[234]
	<i>Pheucticus melanocephalus</i>	Black-headed Grosbeak	weak		western	pop	x		[84, 195]
Fringillidae	<i>Agelaius tricolor</i>	Tricolored Blackbird	strong		western	pop			[235–237]
	<i>Sturnella neglecta</i>	Western Meadowlark	intermediate	x	western	pop	x		[233, 238]
	<i>Euphagus cyanocephalus</i>	Brewer's Blackbird	strong	x	western	pop	x		[239–241]
	<i>Molothrus bonariensis</i>	Shiny Cowbird	strong		other	individ	x		[242]
	<i>Molothrus aeneus</i>	Bronzed Cowbird	intermediate		southwest	pop	x		[233]
		<i>Leucosticte tephrocotis</i>	Gray-crowned Rosy-Finch	strong	x	western	both	x	
	<i>Leucosticte atrata</i>	Black Rosy-Finch	strong	x	western	pop			[243, 244]
	<i>Leucosticte australis</i>	Brown-capped Rosy-Finch	strong	x	western	pop			[245–247]
	<i>Pipilo maculatus</i>	Pine Grosbeak	intermediate	x	widespread	individ	x		[17, 128, 248, 249]
	<i>Haemorhous purpureus</i>	Purple Finch	strong	x	widespread	pop	x		[250–253]
	<i>Haemorhous cassinii</i>	Cassin's Finch	strong	x	western	both	x		[254–258]
	<i>Haemorhous mexicanus</i>	House Finch	strong	x	widespread	pop	x		[259, 260]
	<i>Spinus pinus</i>	Pine Siskin	weak		widespread	both	x		[8]
	<i>Spinus psaltria</i>	Lesser Goldfinch	strong	x	western	both	x		[17, 98, 156, 160, 261, 262]
	<i>Spinus lawrencei</i>	Lawrence's Goldfinch	weak		southwest	pop			[125, 263]
	<i>Coccothraustes vespertina</i>	Evening Grosbeak	strong	x	widespread	pop	x		[264–266]
	<i>Loxoides bailleui</i>	Pallid	strong		Hawaii	pop	x		[267, 268]
	<i>Chlorodrepanis virens</i>	Hawaii Amakihi	intermediate		Hawaii	individ	x		[269–272]
	<i>Chlorodrepanis flavus</i>	Oahu Amakihi	weak		Hawaii	individ			[8]
	<i>Chlorodrepanis stejnegeri</i>	Kauai Amakihi	weak		Hawaii	individ			[270]
	<i>Drepanis coccinea</i>	Iiwi	strong		Hawaii	individ			[2, 269, 273, 274]
	<i>Palmeria dolei</i>	Akohekohe	intermediate		Hawaii	individ	x		[275]
	<i>Himatione sanguinea</i>	Apapane	intermediate		Hawaii				[2, 269, 273]

- Henshaw, H.W., *Birds of the Hawaiian Islands: Being a Complete List of the Birds of the Hawaiian Possessions, With Notes on their Habits*. 1902, Honolulu, HI: T.G. Thrum.
- Perkins, R.C.L., *Vertebrata*, in *Fauna Hawaiensis*, D. Sharp, Editor. 1903, Cambridge University Press: Cambridge, England, pp. 365–466. **3.** Munro, G.C., *Birds of Hawaii*. 1944, Honolulu, HI: Tongg Publishing Company. **4.** Baldwin, P.H., *The Hawaiian Goose, its distribution and reduction in numbers*. The Condor, 1945. 47:27–37. **5.** Banko, W.E., and W.H. Elder, *History of endemic Hawaiian birds: Population histories, species accounts: Scrub-grassland birds: Nene-Hawaiian Goose*, in *Avian History Report*. 1990, Cooperative National Park Resources Studies Unit, University of Hawaii at Manoa, Department of Botany: Manoa, HI. **6.** Black, J.M., et al., *Survival, movements, and breeding of released Hawaiian Geese: An assessment of the reintroduction program*. The Journal of Wildlife Management, 1997. 61:1161–1173. **7.** Rojek, N.A., *The development of feeding behavior in nene (Branta sandvicensis) goslings: Implications for captive propagation and release*. 1994, University of Hawaii. **8.** *Unpublished data, personal communication, inference from summer/winter distribution, or other case of no published reference*. **9.** Engilis, A., Jr, and T.K. Pratt, *Status and population trends of Hawaii's native waterbirds, 1977–1987*. The Wilson Bulletin, 1993. 105:142–158. **10.** Snyder, L.L., and H.G. Lumsden, *Variation in Anas cyanoptera*. 1951: University of Toronto Press. **11.** Dzinbal, K.A., *Coastal Feeding Ecology of Harlequin Ducks in Prince William Sound, Alaska, During Summer in Marine Birds: Their Feeding Ecology and Commercial Fisheries Relationships*, D.N. Nettleship, G.A. Sanger, P.F. Springer, Editors. 1982, Canadian Wildlife Service, pp. 6–10. **12.** Crowley, D.W., *Breeding Habitat of Harlequin Ducks in Prince William Sound, Alaska*. 1994, Oregon State University. **13.** Erskine, A.J., *Buffleheads*. 1972, Ottawa Canada: Environment Canada Wildlife Service. **14.** Campbell, R.W., N.K. McTaggart-Cowan, and I. Dawe, *Birds of British Columbia, Volume 1: Nonpasserines: Introduction, Loons through Waterfowl*. 1990, Vancouver, BC, Canada: University of British Columbia Press. **15.** Cramp, S., and K.E.L. Simmons, *The Birds of the Western Palearctic, Vol. 1: Ostrich to Ducks*. Handbook of the Birds of Europe, the Middle East, and North Africa. Vol. 1. 1977, Oxford, England: Oxford University Press. **16.** Bellrose, F.C., *Ducks, Geese, and Swans of North America*. Rev. ed. 1980, Harrisburg, PA: Stackpole Books. **17.** Small, A., *California Birds: Their Status and Distribution*. 1994, Vista, CA: Ibis Publishing Co. **18.** Barlow, C., and W.W. Price, *A List of the land birds of the Placerville-Lake Tahoe stage road, central Sierra Nevada Mountains, California*. The Condor, 1901. 3:151–184. **19.** Belding, L., *The Fall Migration of Oreortyx pictus plumiferus*.

TABLE 1. Continued.

- The Condor, 1903. 5:18. **20.** McLean, D.D., *The quail of California*. Game bulletin ; no. 2. 1930, Sacramento, CA: California Division of Fish and Game. **21.** Ormiston, J.H., *The food habits, habitat and movements of Mountain Quail in Idaho*. 1966, University of Idaho: Moscow. ID. **22.** Vogt, J., *Bird notes from Lassen Volcanic National Park*. The Condor, 1941. 43:161–162. **23.** Miller, E.V., *Life history and management of Mountain Quail in California*. 1950, California. Division of Fish and Game: Sacramento, CA. **24.** Heekin, P.E., C.A. Vogel, and K.P. Reese, *Uncovering the elusive habits of Mountain Quail in Idaho*. Quail Unlimited, 1994. 13(3):14–16. **25.** Gullion, G.W., *Organization and movements of coveys of a Gambel Quail population*. The Condor, 1962. 64:402–415. **26.** Brown, D.E., *Arizona Game Birds*. 1989. **27.** Rosene, W., *The Bobwhite Quail, Its Life and Management*. 1969, New Brunswick, N.J.: Rutgers University Press. **28.** Pyrah, D.B., *A preliminary study toward sage grouse management in Clark and Fremont counties based on seasonal movements*. 1954, University of Idaho: Moscow. ID. **29.** Crawford, J.E., Jr, *The movements, productivity, and management of Sage Grouse in Clark and Fremont Counties, Idaho*. 1960, University of Idaho: Moscow, ID. **30.** Gill, R.B., and F.A. Glover, *Daily and seasonal movements of Sage Grouse*. 1965, Colorado Cooperative Wildlife Research Unit: Fort Collins, CO. **31.** Wallestad, R.O., *Summer movements and habitat use by Sage Grouse broods in central Montana*. The Journal of Wildlife Management, 1971. 35:129–136. **32.** Connelly, J.W., H.W. Browsers, and R.J. Gates, *Seasonal movements of sage grouse in southeastern Idaho*. Journal of Wildlife Management, 1988. 52:116–122. **33.** Wakkinen, W.L., *Nest Site Characteristics and Spring–Summer Movements of Migratory Sage Grouse in Southeastern Idaho*. 1990: University of Idaho. **34.** Fischer, R.A., K.P. Reese, and J.W. Connelly, *Influence of vegetative moisture content and nest fate on timing of female Sage Grouse migration*. The Condor, 1996. 98:868–872. **35.** Fischer, R.A., et al., *Nesting-area fidelity of Sage Grouse in southeastern Idaho*. The Condor, 1993. 95:1038–1041. **36.** Herzog, P.W., and D.M. Keppie, *Migration in a local population of Spruce Grouse*. The Condor, 1980. 82:366–372. **37.** Schroeder, M.A., *Behavioral differences of female Spruce Grouse undertaking short and long migrations*. The Condor, 1985. 87:281–286. **38.** Schroeder, M.A., and D.A. Boag, *Dispersal in Spruce Grouse: Is inheritance involved? Animal Behaviour*, 1988. 36:305–307. **39.** Weeden, R.B., and L.N. Ellison, *Upland Game Birds of Forest and Tundra*. 1968: Alaska Division of Game. **40.** Mossop, D.H., *Winter survival and spring breeding strategies of Willow Ptarmigan, in Adaptive Strategies and Population Ecology of Northern Grouse: Theories and Synthesis*, A.T. Bergerud and M.W. Graton, Editors. 1988, University of Minnesota Press. pp. 330–378. **41.** Kessel, B., *Birds of the Seward Peninsula, Alaska: Their Biogeography, Seasonality, and Natural History*. 1989: University of Alaska Press. **42.** Gruys, R.C., *Autumn and winter movements and sexual segregation of Willow Ptarmigan*. Arctic, 1993. 46:228–239. **43.** Braun, C.E., and R.K. Schmidt, *Effects of snow and wind on wintering populations of White-tailed Ptarmigan in Colorado*. Proc. Snow and Ice Symposium, 1971. pp. 238–250. **44.** Hoffman, R.W., and C.E. Braun, *Migration of a wintering population of White-tailed Ptarmigan in Colorado*. The Journal of Wildlife Management, 1975. 39:485–490. **45.** Braun, C.E., R.W. Hoffman, and G.E. Rogers, *Wintering areas and winter ecology of White-tailed Ptarmigan in Colorado, in Colorado Division of Wildlife Special Report*. 1976, Colorado Division of Wildlife. **46.** Herzog, P.W., *Summer habitat use by White-tailed Ptarmigan in southwestern Alberta*. Canadian Field-Naturalist, 1977. 91:367–371. **47.** Hoffman, R.W., and C.E. Braun, *Characteristics of a wintering population of White-tailed Ptarmigan in Colorado*. The Wilson Bulletin, 1977. 89:107–115. **48.** Anthony, A.W., *Migration of Richardson's Grouse*. The Auk, 1903. 20:24–27. **49.** Wing, L., *Seasonal movements of the Blue Grouse*. North American Wildlife Conference Transcript, 1947. 12:504–509. **50.** Hamerstrom, F.N., and F. Hamerstrom, *Mobility of the Sharp-tailed Grouse in relation to its ecology and distribution*. The American Midland Naturalist, 1951. 46:174–226. **51.** Ulliman, M.J., *Winter habitat ecology of Columbian Sharp-tailed Grouse in southeastern Idaho*. 1995, University of Idaho: Moscow, ID. **52.** Snyder, L.L., *A study of the Sharp-tailed Grouse*. University Toronto Studies Biol., 1935(40):1–65. **53.** Jackson, J.A., *American Black Vulture, in Handbook of North American Birds*, R. Palmer, Editor. 1988. **54.** Mossman, M.J. *Black and Turkey Vultures, in Midwest Raptor Management Symposium and Workshop*. 1991. **55.** Goodrich, L.J., and J.P. Smith, *Raptor Migration in North America, in The State of North American Birds of Prey*, K.L. Bildstein, J.P. Smith, and E. Ruelas Inzunza, Editors. 2008, Nuttall Ornithological Club and American Ornithologist's Union: Cambridge, MA. **56.** Friedmann, H., et al., *Distributional Check-list of the Birds of Mexico, Part 1*. 1950. **57.** Steenhof, K., M.N. Kochert, and M.Q. Moritsch, *Dispersal and migration of southwestern Idaho raptors*. Journal of Field Ornithology, 1984. 55:357–368. **58.** Greij, E.D., *Common Moorhen, in Migratory Shore and Upland Game Bird Management in North America*, T.C. Tacha and C.E. Braun, Editors. 1994, International Association of Fish and Wildlife Agencies, US Department of the Interior: Washington, D.C. pp. 144–157. **59.** Gorenzel, W.P., R.A. Ryder, and C.E. Braun, *American Coot distribution and migration in Colorado*. The Wilson Bulletin, 1981. 93:115–118. **60.** Gorenzel, W.P., R.A. Ryder, and C.E. Braun, *Reproduction and nest site characteristics of American Coots at different altitudes in Colorado*. The Condor, 1982. 84:59–65. **61.** Wright, N.J.R., and D.W. Matthews, *New nesting colonies of the Ivory Gull Pagophila eburnea in southern East Greenland*. Dansk Ornithologisk Forenings Tidsskrift, 1980. 74:59–64. **62.** Frisch, T., and W.C. Morgan, *Ivory Gull colonies in southern Ellesmere Island, Arctic Canada*. Canadian Field-Naturalist, 1979. 93:173–174. **63.** Jehl, J.R., Jr., and C. Chase III, *Foraging patterns and prey selection by avian predators: A comparative study in two colonies of California Gulls*. Studies in Avian Biology, 1987. 10:91–101. **64.** Wiley, J.W., *The White-Crowned Pigeon in Puerto Rico: Status, distribution, and movements*. The Journal of Wildlife Management, 1979. 43:402–413. **65.** Holt, D.W., and R. Kline, *Glaring grouse*. Montana Outdoors, 1989. 20:13–15. **66.** Cameron, E.S., *The birds of Custer and Dawson counties, Montana*. The Auk, 1907. 24:389–406. **67.** Johnson, H.C., *Pigmy Owl in town*. The Condor, 1903. 5:81. **68.** Bent, A.C., *Life Histories of North American Birds of Prey, Part Two*. 1938, New York, NY: Dover Publications, Ltd. **69.** Farley, F.L., *The Pygmy Owl: An Alberta Bird*. Canadian Field-Naturalist, 1987. 51:86–87. **70.** Campbell, R.W., N.K. McTaggart-Cowan, and I. Dawe, *Birds of British Columbia, Volume 2: Nonpasserines: Diurnal Birds of Prey through Woodpeckers*. 1990, Vancouver, BC: UBC Press. **71.** Laymon, S., *Ecology of the Spotted Owl in the central Sierra Nevada, California*. 1988, University of California. Berkeley: Ann Arbor. p. 303. **72.** Verner, J., R.J. Gutierrez, and G.I. Gould, Jr., *The California Spotted Owl: General biology and ecological relations, in The California Spotted Owl: A Technical Assessment of its Current Status*, J. Verner, et al., Editors. 1992, US Forest Service: Pacific Southwest Research Station. pp. 55–78. **73.** Ganey, J.L., R.B. Duncan, and W.M. Block, *Use of oak and associated woodlands by Mexican Spotted Owls in Arizona, in Ecology and Management of Oak and Associated Woodlands*. 1992, USDA Rocky Mountain Forest and Range Experiment Station: Sierra Vista, AZ. pp. 125–128. **74.** Allen, H., and L. Brewer, *Progress report no. 2 for the cooperative administrative study to monitor Spotted Owl management areas in national forests in Washington*. 1986, Washington Department of Game: Olympia, WA. **75.** Willey, D.W., *Home-range characteristics and juvenile dispersal ecology of Mexican Spotted Owls in*

TABLE 1. Continued.

- southern Utah, in *Recovery Plan for the Mexican Spotted Owl*. 1993, Utah Division of Wildlife Resource; US Dept. Interior Fish and Wildlife Service: Salt Lake City, UT. **76.** Zwank, P.J., and G.M. Southward, *Habitat characteristics of Mexican Spotted Owls in southern New Mexico*. Journal of Field Ornithology, 1994. 65:324–334. **77.** Franklin, A.B., *Breeding biology of the Great Gray owl in southeastern Idaho and northwestern Wyoming*. The Condor, 1988. 90:689–696. **78.** Bull, E.L., M.G. Henjum, and R.S. Rohweder, *Home range and dispersal of Great Gray Owls in northeastern Oregon*. Journal of Raptor Research, 1988. 22:101–106. **79.** Nero, R.W., *The Great Gray Owl: Phantom of the northern forest*. 1980, Washington, DC: Smithsonian Institution Press. **80.** Cannings, R., *Northern Saw-whet Owl*. Birds of North America, 1993. 42:1–20. **81.** Barb, M.A., *Natural History of the Northern Saw-whet Owl (Aegolius acadicus) in Southern Appalachian Mountains*. 1995, Johnson City, TN: East Tennessee State University. p. 136. **82.** Garrett, K., and J. Dunn, *Birds of Southern California: Status and Distribution*. 1981, Los Angeles, CA: Los Angeles Audubon Society. **83.** Woods, C., *Ecological aspects of torpor use and inactivity during winter by Common Poorwills*. 2003, University of Regina, SK, Canada. **84.** Howell, S.N.G., and S. Webb, *A Guide to the Birds of Mexico and Northern Central America*. 1995, Oxford, England: Oxford University Press. **85.** Andrews, R., and R. Righter, *Colorado Birds: A Reference to their Distribution and Habitat*. 1992, Denver, CO: Denver Museum of Natural History. **86.** Monson, G., and A.R. Phillips, *Annotated Checklist of the Birds of Arizona*. 1981, Tucson, AZ: University of Arizona Press. **87.** Lyon, D.L., *A montane hummingbird territorial system in Oaxaca, Mexico*. The Wilson Bulletin, 1976. 88:280–299. **88.** Des Granges, J.L., *Organization of a tropical nectar feeding bird guild in a variable environment*. Living Bird, 1978. 17:199–236. **89.** Land, H.C., *Birds of Guatemala*. 1970, International Committee for Bird Preservation, Pan-American Section. **90.** Monroe, B.L., *A Distributional Survey of the Birds of Honduras*. 1968, Ornithological Monographs 7:1–458. **91.** Stiles, F.G., and A.F. Skutch, *A Guide to the Birds of Costa Rica*. 1989, Ithaca, NY: Cornell University Press. **92.** Arizmendi, M.D., *Interacciones ecológicas múltiples: el caso del sistema mutualista colibríes-plantas y el ladrón de néctar Diglossa baritula (Passeriformes: Aves)*. 1994, Universidad Nacional Autónoma de México: México, D.F. **93.** Fox, R.P., *Plumages and territorial behavior of the Lucifer Hummingbird in the Chisos Mountains, Texas*. The Auk, 1954. 71:465–466. **94.** Gaines, D., *Birds of Yosemite and the East Slope*. 1988, Lee Vining, CA: Artemisia Press. **95.** Grinnell, J., *The Biota of the San Bernardino Mountains*. Vol. 5. 1908, Berkeley, CA: University of California, Berkeley, Museum of Vertebrate Zoology. **96.** Ortiz-Crespo, F.I., *Agonistic and Foraging Behavior of Hummingbirds Co-occurring in Central Coastal California*. 1980, University of California: Berkeley, CA. **97.** Stiles, F.G., *Food supply and the annual cycle of the Anna Hummingbird*. 1973, Berkeley, CA: University of California Press. **98.** Weathers, W.W., and D. Weathers, *Birds of Southern California's Deep Canyon*. 1983, Berkeley, CA. **99.** Willet, G., *Birds of the Pacific Slope of Southern California*. Pacific Coast Avifauna, J. Grinnell and H.S. Swarth, eds. 1912, Hollywood, CA: Cooper Ornithological Club. **100.** Bendire, C., *Life Histories of North American Birds, from the Parrots to the Grackles: With Special Reference to their Breeding Habits and Eggs*. 1895, Washington DC: Govt. Printing Office. **101.** Brewster, W., *Birds of the Cape Region of Lower California*. 1902, Cambridge, MA: Museum of Comparative Zoology at Harvard College. **102.** Wilbur, S.R., *Birds of Baja California*. 1987, Berkeley, CA: University of California Press. **103.** Bock, C.E., *The Ecology and Behavior of the Lewis Woodpecker (Asyndesmus lewis)*. University of California Publications in Zoology, G.A. Bartholomew, et al., eds. Vol. 92. 1970, Berkeley, CA: University of California Press. **104.** Snow, R.B., *A natural history of the Lewis Woodpecker Asyndesmus lewis (Gray)*. 1941, University of Utah: Salt Lake City, UT. **105.** Hadow, H.H., *Winter ecology of migrant and resident Lewis' Woodpeckers in southeastern Colorado*. The Condor, 1973. 75:210–224. **106.** Cooper, J.M., and C. Gillies, *Breeding distribution of the Lewis's Woodpecker in the East Kootenay Trench in relation to fire history*. In *Biology and Management of Species and Habitats at Risk*. 1999, Kamloops, BC: B.C. Ministry of Environment, Lands and Parks and University College of the Cariboo. **107.** Kaufman, K., and R. Troy, *Lives of North American Birds*. 1996, Boston, MA: Houghton Mifflin Co. **108.** Phillips, A., J. Matshall, and G. Monson, *The Birds of Arizona*. 1964, Tucson, AZ: University of Arizona Press. **109.** Hadow, H.H., *Audible communication and its role in the species recognition of Red-naped and Williamson's sapsuckers (Aves: Piciformes)*. 1977, University of Colorado: Boulder, CO. p. 77. **110.** Devillers, P., *Identification and distribution in California of the Syphrapicus various group of sapsuckers*. California Birds, 1970. 1(2):47–76. **111.** Howell, T.R., *Natural history and differentiation in the Yellow-bellied Sapsucker*. The Condor, 1952. 54:237–282. **112.** Richards, E.B., *A list of the land birds of the Grass Valley District, California*. The Condor, 1924. 26:98–104. **113.** Voous, K.H., *On the history of the distribution of the genus Dendrocopos*. Limosa, 1947. 20:1–142. **114.** Bent, A.C., *Life Histories of North American Woodpeckers: Order Piciformes*. 1939, Washington, DC: US Govt. Printing Office. **115.** Short, L.L., *Hybridization in the flickers (Colaptes) of North America*. Bulletin of the American Museum of Natural History, 1965. 129:307–428. **116.** Burleigh, T.D., *Birds of Idaho*. 1972, Caldwell, ID: Caxton Printers. **117.** Simpson, M.B., Jr., *Birds of the Blue Ridge Mountains*. 1992, Chapel Hill, NC: University of North Carolina Press. **118.** Gram, W.K., *Winter participation by neotropical migrant and resident birds in mixed-species flocks in northeastern Mexico*. The Condor, 1998. 100:44–53. **119.** Hutto, R.L., *A description of mixed-species insectivorous bird flocks in western Mexico*. The Condor, 1987. 89:282–292. **120.** Miller, A.H., et al., *Distributional Checklist of the Birds of Mexico: Part 2*. Pacific Coast Avifauna. Vol. No. 33. 1957: Cooper Ornithological Society. **121.** Whetstone, J., *Yecora, Sonora, Mexico, in Ninety-eighth Christmas Bird Count*, G.S. Le Baron, Ed. 1998, New York, NY: National Audubon Society. **122.** Russell, S.M., and G. Monson, *The Birds of Sonora*. 1998, Tucson, AZ: University of Arizona Press. **123.** Lehman, P.E., *The Birds of Santa Barbara County, California*. 1994, Santa Barbara, CA: Vertebrate Museum, University of California. **124.** Rosenberg, K.V., et al., *Birds of the Lower Colorado River Valley, in Birds of the Lower Colorado River Valley*. 1991. **125.** Beedy, E.C., and S.L. Granholm, *Discovering Sierran Birds: Western Slope*. 1985, Three Rivers, CA: Yosemite Natural History Association and Sequoia National History Association. **126.** Gilligan, J., et al., *Birds of Oregon: Status and Distribution*. 1994, McMinnville, OR: Cinclus Publications. **127.** Raffaele, H.A., T. Pederson, and K. Williams, *A Guide to the Birds of the West Indies*. 1998, Princeton, NJ: Princeton University Press. **128.** Bailey, A.M., and R.J. Niedrach, *Birds of Colorado*. 1965, Denver, CO: Museum of Natural History. **129.** Simpson, G.B., *Oregon Jays*. The Canadian Field-Naturalist, 1925. 39:29–30. **130.** Skinner, M.P., *Notes on the Rocky Mountain Jay in the Yellowstone National Park*. The Condor, 1921. 23:147–151. **131.** Bent, A.C., *Life Histories of North American Jays, Crows, and Titmice, Part 1*. U.S. National Museum Bulletin, 1946. 191:118–128. **132.** Westcott, P.W., *Relationships among three species of jays wintering in southeastern Arizona*. The Condor, 1969. 71:353–359. **133.** Dunn, J.L., and K.L. Garrett, *Parapatry in Woodhouse's and California Scrub-Jays revisited*. Western Birds, 2001. 32:186–187. **134.** Hubbard, J.P., *Revised check-list of the birds of New Mexico*. 1978. **135.** Linsdale, J.M., *A list of the birds of Nevada*. The Condor, 1951. 53:228–249. **136.** Monson, G., *Recent notes from the lower Colorado River*

TABLE 1. Continued.

- Valley of Arizona and California. The Condor, 1949. 51:262–265. **137.** Skinner, M.P., *The nutcrackers of Yellowstone Park*. The Condor, 1916. 18:62–64. **138.** Guintoli, J., and L.R. Mewaldt, *Stomach contents of Clark's Nutcracker collected in western Montana*. The Auk, 1978. 95:595–598. **139.** Hutchins, H.E., and R.M. Lanner, *The central role of Clark's Nutcracker in the dispersal and establishment of whitebark pine*. Oecologia, 1982. 55:192–201. **140.** Tomback, D.F., *Foraging strategies of Clark's Nutcracker*. Living Bird, 1978. 16:123–161. **141.** Tomback, D.F., and K.A. Kramer, *Limber pine seed harvest by Clark's Nutcracker in the Sierra Nevada: Timing and behavior*. The Condor, 1980. 82:467–468. **142.** Torick, L.L., *The interaction between Clark's Nutcracker and Ponderosa Pine, a "wind-dispersed" pine: Energy efficiency and multi-genet growth forms*. 1995, University of Colorado at Denver. **143.** Vander Wall, S.B., and H.E. Hutchins, *Dependence of Clark's Nutcracker, Nucifraga columbiana, on conifer seeds during the postfledging period*. Canadian Field-Naturalist, 1983. 97:208–214. **144.** Campbell, R.W., et al., *Birds of British Columbia, Volume 3: Passerines: Flycatchers through Vireos*. 1997, Vancouver, BC: UBC Press. **145.** Alcorn, J.R., *The Birds of Nevada*. 1988, Fallon, NV: Fairview West Publishers. **146.** Giffin, J.G., J.M. Scott, and S. Mountainspring, *Habitat selection and management of the Hawaiian Crow*. Journal of Wildlife Management, 1987. 51:485–494. **147.** Bent, A.C., *Life Histories of North American Flycatchers, Larks, Swallows, and their Allies*. US National Museum Bulletin, 1942. 179. **148.** Jewett, S.G., *A new Horned Lark from the state of Washington*. The Auk, 1943. 60:262–263. **149.** McCallum, D.A., *Alternative emigration strategies and the adaptive significance of natal dispersal in a population of Mountain Chickadees (Parus gambeli)*. 1988, University of New Mexico, Albuquerque. **150.** McCallum, D.A., *Variable cone crops, migration, and dynamics of a population of Mountain Chickadees (Parus gambeli), in Population Biology of Passerine Birds: An Integrated Approach*, J. Blondel, et al., Editors. 1990. **151.** Nosedal, J., *Local migrations of insectivorous birds in western Mexico: Implications for the protection and conservation of their habitats*. Bird Conservation International, 1994. 4(2/3):129–142. **152.** Nosedal, J., *Seasonal dynamics of foliage-gleaning insectivorous birds in southern Durango, Mexico, in Conservation of Neotropical Migratory Birds in Mexico*, M.H. Wilson and S.A. Sader, Editors. 1995, Maine Agric. For. Exp. Stn., Misc. Publ. 727. pp. 81–97. **153.** Swarth, H.S., *The California forms of the genus Psaltriparus*. The Auk, 1914. 31:499–526. **154.** Phillips, A.R., *The Known Birds of North and Middle America: Hirundinidae to Mimidae; Certhiidae*. In *The Known Birds of North and Middle America: Distributions and Variation, Migrations, Changes, Hybrids, Etc.* 1986, Denver, CO: A.R. Phillips. **155.** Cannings, R.A., R.J. Cannings, and S.G. Cannings, *Birds of the Okanagan Valley, British Columbia*. Nature Canada, 1988. 17(2):50. **156.** Grinnell, J., and A.H. Miller, *The Distribution of the Birds of California*. Pacific Coast Avifauna 1944, Berkeley, CA: Cooper Ornithological Club. **157.** Davis, W.A., and S.M. Russell, *Birds in Southeastern Arizona*. 1979: Tucson Audubon Society. **158.** Norris, R.A., *Comparative biosystematics and life history of the nuthatches Sitta pygmaea and Sitta pusilla*. University of California Publications in Zoology. 1958, Berkeley, CA: University of California Press. **159.** Roberson, D., *Pygmy Nuthatch, in Atlas of the Breeding Birds of Monterey County, California*, C. Tenney and S.F. Bailey, Editors. 1993, Monterey Peninsula Audubon Society: Monterey, CA. **160.** Unitt, P., *The birds of San Diego County*. 1984, San Diego, CA: San Diego Society of Natural History. **161.** Campbell, K.F., R.A. Erickson, and S.F. Bailey, *The autumn migration: Middle Pacific coast region*. American Birds, 1988. 42:127–134. **162.** Campbell, K.F., R.A. Erickson, and S.F. Bailey, *The winter season: Middle Pacific coast region*. American Birds, 1988. 42:314–320. **163.** McCaskie, G., *The autumn migration: Southern Pacific coast region*. American Birds, 1998. 42:314–320. **164.** McCaskie, G., *The winter season: Southern Pacific coast region*. American Birds, 1988. 42:113–116. **165.** Witzeman, J., and J. Witzeman, *The autumn migration: Southwest region, Arizona*. American Birds, 1988. 42:303–305. **166.** Stupka, A., *Notes on the Birds of Great Smoky Mountains National Park*. 1963, Knoxville, TN: University of Tennessee Press. **167.** Baumgartner, F.M., and A.M. Baumgartner, *Oklahoma Bird Life*. 1992, Norman, OK: University of Oklahoma Press. **168.** Baumgartner, F.M., and C.P. Nicholson, *Winter Wren*. Atlas of the Breeding Birds of Tennessee. 1997, Knoxville, TN: University of Tennessee Press. **169.** McCaskie, G., et al., *Birds of northern California, an annotated field list*. 2nd ed. 1988, Berkeley, CA: Golden Gate Audubon Society. **170.** Stedman, B.H., and C.P. Nicholson, *Winter Wren*. Atlas of the Breeding Birds of Tennessee. 2009. 40:29–34. **171.** Price, F.E., and C.E. Bock, *Population Ecology of the Dipper (Cinclus mexicanus) in the Front Range of Colorado*. Studies in Avian Biology No. 7. 1983, Las Cruces, NM: University of Colorado and Cooper Ornithological Society. **172.** Morrissey, C.A., *Effect of altitudinal migration within a watershed on the reproductive success of American Dippers*. Canadian Journal of Zoology, 2004. 82:800–807. **173.** Morrissey, C.A., and R.J. Olenick, *American Dipper, Cinclus mexicanus, preys upon larval tailed frogs, Ascaphus truei*. Canadian Field-Naturalist, 2004. 118:446–448. **174.** Faulkner, D., *Birds of Wyoming*. 2010, Greenwood Village, CO: Roberts and Company. **175.** Hayward, C.L., et al., *Birds of Utah*. Great Basin Naturalist Memoirs no. 1. 1976: Brigham Young University Press. **176.** Marshall, D.B., M.G. Hunter, and A. Conteras, *Birds of Oregon: A General Reference*. 2003, Corvallis, OR: Oregon State University Press. **177.** Wahl, T.R., B. Tweit, and S.G. Mlodinow, eds. *Birds of Washington: Status and Distribution*. 2005, Oregon State University Press. **178.** Wahl, T.R., B. Tweit, and S.G. Mlodinow, eds. *Birds of Washington: Status and Distribution*. 1949, Washington, DC: U.S. Govt. Printing Office. **179.** Bent, A.C., *Life Histories of North American Thrushes, Kinglets, and their Allies: Order Passeriformes*. 1949, Washington, DC: U.S. Govt. Printing Office. **180.** Popper, D.M., *Notes on the birds of Mount Locke, Texas*. The Condor, 1951. 53:51–54. **181.** Wauer, R.H., *Birds of Zion National Park and vicinity*. 1997, Logan, UT: Utah State University. **182.** Johnson, T.B., *Resident and North American migrant bird interactions in the Santa Marta highlands, northern Colombia, in Migrant Birds in the Neotropics: Ecology, Behavior, Distribution, and Conservation*, A. Keast and E.S. Morton, Editors. 1980, Smithsonian Institution Press: Washington, DC. pp. 239–247. **183.** Engels, W.L., *Structural Adaptations in Thrashers (Mimidae: Genus Toxostoma): With Comments on Interspecific Relationships*. 1940, Berkeley, CA: University of California Press. **184.** Grinnell, J., *Midwinter birds at Palm Springs, California*. The Auk, 1904. 6:40–45. **185.** Meams, E.A., *Some birds of Arizona*. The Auk, 1886. 3:289–307. **186.** Stephens, F., *Notes on birds observed in the Colorado Desert in winter*. The Auk, 1890. 7:296–297. **187.** Van Rossem, A.J., *Winter birds of the Salton Sea region*. The Condor, 1911. 13:129–137. **188.** Wauer, R.H., *Birds of Big Bend National Park and Vicinity*. 1973, Austin, TX: University of Texas Press. **189.** Verbeek, N.A.M., *Breeding ecology of the Water Pipit*. The Auk, 1970. 87:425–451. **190.** Wilson, S., and K. Martin, *Songbird use of high-elevation habitat during the fall post-breeding and migratory periods*. *Ecoscience*, 2005. 12:561. **191.** Root, T.L., *Atlas of Wintering North American Birds: An Analysis of Christmas Bird Count Data*. 1988, Chicago, IL: University of Chicago Press. **192.** Phillips, A.R., *The Known Birds of North and Middle America: Bombycillidae; Sylviidae to Sturnidae; Vireonidae*. In *The Known Birds of North and Middle America: Distributions and Variation, Migrations, Changes, Hybrids, Etc.* 1991, Denver, CO: A.R. Phillips. **193.** Anderson, B.W., and R.D. Ohmart, *Phainopepla utilization of honey mesquite forests in the Colorado River Valley*. The Condor,

TABLE 1. Continued.

1978. 80:334–338. **194.** Crouch, J.E., *Distribution and habitat relationships of the Phainopepla*. The Auk, 1943. 60:319–333. **195.** Swarth, H.S., *Birds of the Huachuca Mountains, Arizona*. 1904, Los Angeles, CA: Cooper Ornithological Club of California. **196.** Walsberg, G.E., *Ecology and energetics of contrasting social systems in Phainopepla (Aves: Ptilonotidae)*. University of California Publications in Zoology, 1977. 108:1–63. **197.** Marshall, J.T., Jr., *Birds of pine-oak woodland in southern Arizona and adjacent Mexico*. Pacific Coast Avifauna, 1957(32):1–125. **198.** Banks, K.W., et al., *Origins, population structure and movements of Snow Buntings *Plectrophenax nivalis* wintering in Highland Region, Scotland*. Bird Study, 1991. 38:10–19. **199.** Smith, R.D., et al., *Age and sex variation in choice of wintering site by Snow Buntings: The effect of altitude*. Ardea, 1993. 81:47–52. **200.** Lefebvre, G., and B. Poulin, *Seasonal abundance of migrant birds and food resources of Panamanian mangrove forests*. The Wilson Bulletin, 1996. 108:748–759. **201.** Rimmer, C.C., and K.P. McFarland, *Migrant stopover and postfledging dispersal at a montane forest site in Vermont*. Wilson Bulletin, 2000. 112:124–136. **202.** Curson, J., D. Quinn, and D. Beadle, *Warblers of the Americas: An Identification Guide*. 1994, New York, NY: Houghton Mifflin. p. 252. **203.** Hubbard, J.P., *The extent and sequence of the molts of the Yellow-rumped Warbler*. Nemouria, 1980. 25:9. **204.** Nosedal, J., *Foraging ecology of foliage-gleaning insectivorous birds of an oak-pine woodland of southern Durango, Mexico*. 1994, New Mexico State University: Las Cruces, NM. **205.** Chapman, F.M., *The Warblers of North America*. 1907, New York, NY: Appleton & Company. **206.** Bailey, F.M., and W.W. Cooke, *Birds of New Mexico*. 1928, Santa Fe, NM: New Mexico Department of Game and Fish in cooperation with the State Game Protective Association and the Bureau of Biological Survey. **207.** Morton, M.L., *Postfledging dispersal of Green-tailed Towhees to a subalpine meadow*. The Condor, 1991. 93:466–468. **208.** Brown, B.T., S.W. Carothers, and R.R. Johnson, *Grand Canyon Birds: Historical Notes, Natural History, and Ecology*. 1987, Tucson, AZ: University of Arizona Press. **209.** Gaines, D., *Birds of the Yosemite Sierra: A Distributional survey*. 1977, Oakland, CA: CAL-SYL Press. **210.** Hubbard, J.P., *Geographic Variation in Non-California Populations of the Rufous-crowned Sparrow*. 1975: Delaware Museum of Natural History. **211.** Phillips, A.R., *Aimophila ruficeps scottii* (Sennett) *Scott's Rufous-crowned Sparrow*, in *Life Histories of North American Cardinals, Grosbeaks, Buntings, Towhees, Finches, Sparrows, and Allies*, A.C. Bent and O.L. Austin, Editors. 1968, Dover Publications: New York, NY. pp. 923–930. **212.** Cord, B., and J.R. Jehl, *Distribution, biology, and status of a relict population of Brown Towhee (*Pipilo fuscus eremophilus*)*. Western Birds, 1979. 10:131–156. **213.** Davis, J., *Distribution and Variation of the Brown Towhees*. 1951, Berkeley, CA: University of California Publications in Zoology no 52, pp. 1–120. **214.** Oberholser, H.C., *The Bird Life of Texas*. Vol. 2. 1974, Austin, TX: University of Texas Press. **215.** Zimmerman, D.A., M.A. Zimmerman, and J.N. Durrie, *New Mexico Bird Finding Guide*. 1992, Albuquerque, NM: New Mexico Ornithological Society. **216.** Johnson, N.K., and J.A. Marten, *Macrogeographic patterns of morphometric and genetic variation in the Sage Sparrow complex*. The Condor, 1992. 94:1–19. **217.** Packard, F.M., *The birds of Rocky Mountain National Park, Colorado*. The Auk, 1945. 62:371–394. **218.** Streeter, R.G., *Field Notes*. Colorado Field Ornithologist, 1969. 6:27. **219.** Garrett, K., et al., *Call notes and winter distribution in the Fox Sparrow complex*. Birding, 2000. 32:412–417. **220.** Swarth, H.S., *Revision of the avian genus *Passerella* with special reference to the distribution and migration of the races in California*. University of California Publications in Zoology, 1920. 21(4). **221.** Behle, W.H., *The Birds of Northeastern Utah*. 1981, Salt Lake City, UT: Utah Museum of Natural History, University of Utah. **222.** Davis, A., and P. Arceze, *An examination of migration in Song Sparrows using banding recovering data*. North American Bird Bander, 1999. 24:124–128. **223.** Patten, M., G. McCaskie, and P. Unitt, *Birds of the Salton Sea: Status, Biogeography, and Ecology*. 2003, Berkeley, CA: University of California Press. **224.** Cortopassi, A.J., and L.R. Mewaldt, *The circumannual distribution of White-Crowned Sparrows*. Bird-Banding, 1965. 36(3):141–169. **225.** Grinnell, J., and T.I. Storer, *Animal Life in Yosemite: An Account of the Mammals, Birds, Reptiles, and Amphibians in a Cross-section of the Sierra Nevada*. 1924: University Press. **226.** Jewett, S.G., *Birds of Washington State*. 1953, Seattle, WA: University of Washington Press. **227.** Rabenold, K.N., and P.P. Rabenold, *Variation in altitudinal migration, winter segregation, and site-tenacity in two subspecies of Dark-eyed Juncos in the southern Appalachians*. The Auk, 1985. 102:805–819. **228.** Horvath, E.G., and K.A. Sullivan, *Facultative migration in Yellow-eyed Juncos*. The Condor, 1988. 90:482–484. **229.** Moore, N.J., *Ethology of the Mexican Junco (*Junco phaeonotus palliatus*)*. 1972, Washington State University: Ann Arbor. p. 192. **230.** Isler, M.L., and P.R. Isler, *The Tanagers: Natural History, Distribution, and Identification*. 1999, Washington, DC: Smithsonian Institution Press. **231.** Zimmer, J.T., *A Study of the Tooth-billed Red Tanager, *Piranga flava**. 1929, Zoological Series of Field Museum of Natural History 7(5):169–219. **232.** DeGraaf, R.M., and J.H. Rappole, *Neotropical Migratory Birds: Natural History, Distribution, and Population Change*. 1995, Ithaca, NY: Cornell University Press. **233.** Bent, A.C., *Life Histories of North American Blackbirds, Orioles, Tanagers, and Allies*. 1958, Washington, DC: U.S. Govt. Printing Office. **234.** Tweit, J.C., and R.C. Tweit, *Messquite-hackberry dominated canyon. The Birds of North America*. A. Poole and F. Gill, eds. Vol. 235. 1991. **235.** DeHaven, R.W., and J.A. Neff, *Recoveries and returns of Tricolored Blackbirds*, 1941–1964. Western Bird Bander, 1973. 48:10–11. **236.** Neff, J.A., *Migration of the Tricolored Red-wing in Central California*. The Condor, 1942. 44:45–53. **237.** Payne, R.B., *Breeding seasons and reproductive physiology of Tricolored Blackbirds and Redwinged Blackbirds*. University California Publications in Zoology, 1969. 90:1–137. **238.** Unitt, P., A. E. Klovestad, W. E. Haas, P. J. Mock, K. J. Winter, A. Mercieca, San Diego County Bird Atlas. 2004, San Diego, CA: San Diego Natural History Museum. **239.** Gullion, G.W., *Birds of the southern Willamette Valley, Oregon*. The Condor, 1951. 53:129–149. **240.** Verbeek, N.A.M., *An ecological study of the Brewer's Blackbird during the winter*. 1962, University of British Columbia. p. 55. **241.** Linsdale, J.M., *Coloration of downy young birds and of nest linings*. The Condor, 1936. 38:111–117. **242.** Pérez-Rivera, R.A., *Parasitism by the Shiny Cowbird in the interior parts of Puerto Rico*. Journal of Field Ornithology, 1986. 57:99–104. **243.** King, J.R., and E.E. Wales, *Observations on migration, ecology, and population flux of wintering Rosy Finches*. The Condor, 1964. 66:24–31. **244.** Hendricks, P., and J.E. Swenson, *Dynamics of the winter distribution of Rosy Finches, *Leucosticte arctica*, in Montana*. Canadian Field-Naturalist, 1983. 97:307–310. **245.** Cooke, W.W., *The migration of North American sparrows*. Bird-Lore, 1913. 15:364–365. **246.** Lane, J.A., and H.R. Holt, *A Birder's Guide to Eastern Colorado*. 1979, Denver, CO: L & P Press. **247.** Packard, F.M., *Brown-capped Rosy Finch*, in *Life Histories of North American Cardinals, Grosbeaks, Buntings, Towhees, Finches, Sparrows, and Allies*, A.C. Bent and O.L. Austin, Jr., Editors. 1968, US National Museum Bulletin: Washington, DC. pp. 373–383. **248.** Grinnell, J., *Birds of the Kotzebue Sound region, Alaska*. 1900, Santa Clara, CA: Cooper Ornithological Club of California. **249.** Price, W.W., *Description of a new Pine Grosbeak from California*. The Auk, 1897. 14:182–186. **250.** Bradley, R.A., and D.W. Bradley, *Co-occurring groups of wintering birds in the lowlands of southern California*. The Auk, 1983. 100:491–493. **251.** Salt, G.W., *The relation of metabolism to climate and distribution in three finches of the*



TABLE 1. Continued.

genus *Carpodacus*. Ecological Monographs, 1952. 22:121–152. **252**. Wheelock, I.G., *Birds of California*, 2nd ed. 1910, Chicago, IL: A.C. McClury and Co. **253**. Widirlechner, M.P., and S.K. Dragula, *Relation of cone-crop size to irruptions of four seed-eating birds in California*. American Birds, 1984. 38:840–846. **254**. Henshaw, H.W., *Report on the Ornithology of the Portions of California Visited During the Field-Season of 1875*, in *United States Geographical Surveys West of the One Hundredth Meridian Annual Reports*, 1876, US Geological Survey: Washington, DC. pp. 224–278. **255**. Mewaldt, L.R., and J.R. King, *Breeding site faithfulness, reproductive biology, and adult survivorship in an isolated population of Cassin's Finches*. The Condor, 1985. 87:494–510. **256**. Orr, R.T., *Cassin's Finch, in Life Histories of North American Cardinals, Grosbeaks, Buntings, Towhees, Finches, Sparrows, and Allies*, A.C. Bent and O.L. Austin, Jr., Editors. 1968. Dover Publications: New York, NY. pp. 280–289. **257**. Samson, F.B., *Territory, breeding density, and fall departure in Cassin's Finch*. The Auk, 1976. 93:477–497. **258**. Van Rossem, A.J., *Birds of the Charleston Mountains, Nevada: Pacific Coast avifauna no. 24*, 1936, Berkeley, CA: Cooper Ornithological Club. **259**. Thompson, W.L., *Agonistic behavior in the House Finch. Part I: Annual cycle and display patterns*. The Condor, 1960. 62:245–271. **260**. Van Riper III, C., *Aspects of House Finch breeding biology in Hawaii*. The Condor, 1976. 78:224–229. **261**. Gaines, D., *Birds of Yosemite and the East Slope*. 2nd edition. 1992. Lee Vining, CA: Artemisia Press. **262**. Van Rossem, A.J., *A Distributional Survey of the Birds of Sonora, Mexico*. Occasional Papers of the Museum of Zoology, Louisiana State University, 1945(21). **263**. Linsdale, J.M., *Goldfinches on the Hastings Natural History Reservation*. The American Midland Naturalist, 1957. 57:1–119. **264**. Butler, A.W., *Some notes concerning the Evening Grosbeak*. The Auk, 1892. 9:238–247. **265**. Scott, A.C., and M. Bekoff, *Breeding behavior of Evening Grosbeaks*. The Condor, 1991. 93:71–81. **266**. Verner, J., and A.S. Boss, *California wildlife and their habitats: Western Sierra Nevada*. 1980, Berkeley, CA: U.S. Dept. of Agriculture, Forest Service, Pacific Southwest Forest and Range Experiment Station. **267**. Banko, P.C., et al., *Availability of food resources, distribution of invasive species, and conservation of a Hawaiian bird along a gradient of elevation*. Journal of Biogeography, 2002. 29:789–808. **268**. Hess, S.C., et al., *Drepanidine movements in relation to food availability in subalpine woodland on Mauna Kea, Hawaii*. Studies in Avian Biology, 2001. 22:154–163. **269**. Baldwin, P.H., *Annual cycle, environment and evolution in the Hawaiian Honeycreepers: (Aves: Drepaniidae)*. Series: conservation. Studies in Avian Biology, 1986(9). **271**. Van Riper III, C., *Parasites of the Hawaii Nmakiki (*Loxops virens virens*)*, in *Technical Report*. 1975, US International Biological Program. **272**. Van Riper III, C., *The influence of nectar resources on nesting success and movement patterns of the Common Amakihi (*Hemignathus virens*)*. The Auk, 1984. 101:38–46. **273**. Ralph, C.J., and S.G. Fancy, *Demography and movements of 'Apapane and 'iwi in Hawaii*. The Condor, 1995. 97:729–742. **274**. MacMillen, R.E., and F.L. Carpenter, *Evening roosting flights of the honeycreepers *Himatione sanguinea* and *Vestiaria coccinea* on Hawai'i*. The Auk, 1980. 97:28–37. **275**. Conant, S., *Recent observations of endangered birds in Hawai'i's national parks*. 'Elepaio, 1981. 41:55–61.

TABLE 2. The number and percentage of species for which there is evidence of altitudinal migration in the *Birds of North America* life history accounts (Rodewald 2015). Included are all landbird orders and the 14 families within Passeriformes represented by at least 5 species in the region.

Taxon	altitudinal migrant species	
	N	%
Anseriformes	10	21.7
Galliformes	11	57.9
Ciconiiformes	0	0.0
Accipitriformes	4	14.8
Falconiformes	1	14.3
Gruiformes	2	15.4
Charadriiformes	4	4.6
Columbiformes	3	37.5
Psittaciformes	0	0.0
Cuculiformes	0	0.0
Strigiformes	5	26.3
Caprimulgiformes	3	33.3
Apodiformes	8	42.1
Trogoniformes	0	0.0
Coraciiformes	0	0.0
Piciformes	11	52.4
Passeriformes	102	31.6
Tyrannidae	8	23.5
Vireonidae	4	30.8
Corvidae	7	38.9
Hirundinidae	2	28.6
Paridae	3	25.0
Troglodytidae	4	40.0
Turdidae	7	43.8
Mimidae	1	10.0
Calcariidae	1	16.7
Parulidae	8	16.0
Emberizidae	17	39.5
Cardinalidae	4	28.6
Icteridae	5	23.8
Fringillidae	18	52.9

*nobilis*), which apparently made diurnal and seasonal altitudinal movements in search of nectar. The percent of migrants in the extant Hawaiian communities is likely similar to that of historical communities; the scant data on extinct taxa indicates that probably 11 of 38 endemic landbird species were altitudinal migrants. Some nonnative species recently established in North America engage in altitudinal migrations; Chukars (*Alectoris chukar*) breeding  $\geq 1800$  m descend below the snow line during winter (Christensen 1954), mirroring movements in their native range (McGowan and Kirwan 2016).

Fewer than half (46.7%) of the sources referenced in descriptions of altitudinal migration came from peer-reviewed journals. Of 275 references cited in Table 1, 44.2% were books or lengthy monographs (bird atlases, distributional monographs, annotated checklists, or other similar literature). Reports and theses comprised 9.1% of the references. The median publication date was 1980, 25% of

references dated from 1954 or earlier, and the oldest dated from 1876 (Henshaw 1876).

## DISCUSSION

What does this review tell us about altitudinal bird migration generally, and in North America specifically? On a global scale, altitudinal bird migration appears to be common (Figure 1). The summaries of information compiled by North America's leading ornithologists paint a surprising picture: altitudinal migration is similarly common in North America. At a minimum, 26.8% of our landbird species migrate altitudinally somewhere within their range, 20.2% do so within the USA and Canada, representing 30.3% of the species that spend the winter within the continent. Many of these species combine altitudinal and latitudinal movements in the same or different parts of their range. The complexities of migration behaviors exhibited by the North American avifauna is grossly simplified by such coarse tallies. Furthermore, estimates will certainly increase as more research is devoted to short-distance movements. Nevertheless, we can rule out the first hypothesis for why this behavior has been ignored—North America is *not* a global anomaly in having few altitudinal migrants.

The second hypothesis is also largely incorrect; understanding altitudinal migration does advance our fundamental understanding of animal movement generally. Most research questions relating to migration are equally as applicable to altitudinal migrants as latitudinal, long-distance migrants. Examples of core questions in migration research that are well suited to tackling in altitudinal migrant systems include documentation of movement patterns and migratory connectivity via new tracking methods, molt-migration (e.g., Fraser et al. 2010), the consequences of migration for community ecology, changes in migration timing and patterns under changing climates, and studies of how migration constrains and is affected by life-history trade-offs. For each of these topics, it matters little whether birds are migrating to sites 10,000s of kilometers away or only over a few 100s of meters in elevation.

There are certainly some topics in migration research unsuited to altitudinal migration systems. For instance, one cannot study the physiological adaptations that permit sustained, energetically costly flight for long, uninterrupted bouts in a short-distance migrant (Watanabe 2016). Likewise, understanding how trans-equatorial migrants correctly respond to and integrate both austral and northern day-length cues by necessity must involve migrants traversing large latitudinal gradients. However, for plenty of questions, selecting an altitudinal migrant would be preferable. The phenotypic flexibility we typically see in altitudinal migrants and other short-distance

migrants makes addressing fundamental questions more tractable. Evolutionary ecologists can exploit intraspecific behavioral variation to understand how endogenous and exogenous factors interact to influence costs and benefits of migration vs. residency—something that is impossible in obligate long-distance migrants.

The third hypothesis is highly plausible; we probably know so little about these birds' movements due to chance artifacts of ornithological history. While apparently the topic was interesting to naturalists a century ago, publications on this topic are dated. The reason may be related to the geographic distribution of altitudinal migration, which is necessarily constrained by the location of mountains. In the above tallies, half the altitudinal migrants were "western" species, and although 35% more were widespread, often only western populations of those species migrated altitudinally. The historic and current densities of researchers, cities, and centers of higher learning are roughly the inverse of topographical complexity on this continent. To this day, only 33% of ornithologists belonging to the American Ornithological Society live and work in states west of the 100th meridian where the most of the species making these movements live (data courtesy S. W. Gillihan; May 2016). Consequently, it is highly likely that such movements have flown under the radar of the ornithological community because they do not occur in most ornithologists' back yards.

### Why Study Altitudinal Migration?

There are at least 4 main reasons to fill this gap in knowledge.

**(1) Embrace the messy.** No avifauna, including that of North America, consists of species falling neatly into tidy migrant and resident categories (e.g., Seifert et al. 2016), or even the arbitrary "long-distance" and "short-distance" migrant groups—terms that mean different things to different authors (e.g., short-distance = 10s of km (Nilsson et al. 2008), or up to 1,000 km (Kondo and Omland 2007)). Furthermore, other movement categorizations including irruptions and nomadism belie the vast variation found in both latitudinal (e.g., Strong et al. 2015) and altitudinal movements (e.g., White-headed Woodpecker, *Leuconotopicus albolaryvatus*; Richards 1924). Embracing the sometimes messy, facultative, partial migration behaviors brings opportunities to understand the causes of many potential movement responses arising from environmental selective pressures and constraints. In a recent example, Himalayan altitudinal migrants provided the opportunity to contrast physiological adaptations to high elevations in year-round residents compared to more mobile species that must be phenotypically flexible to cope with changing atmospheric conditions over their annual cycle (Barve et al. 2016). Additionally, the study of "messy" movements can lead to results that challenge dogma arising from study of a biased

sample of movement behaviors. An example of this is the molt-migration of many Western songbirds, a pattern important to the understanding of interspecific differences in behavior, evolution, and conservation that has only been described and appreciated in the past few decades (Rohwer and Manning 1990, Leu and Thompson 2002).

A second argument is that studies of “messy” migrations are necessary to fully understand classic obligate migrations. For example, Ramenofsky et al. (2012) highlighted the paucity of studies describing the physiological regulation of facultative migrations, and explained how all migrations are likely governed by combinations of mechanisms. The hypothalamic–pituitary–gonad axis (HPG) is the primary pathway controlling the timing and physiological preparation for obligate migrations, while the hypothalamic–pituitary–adrenal axis (HPA) is probably involved in cueing facultative migrations. However, both types of migrations likely are controlled by both the HPG and HPA to different degrees and/or under different circumstances. As geolocator data is making increasingly obvious (e.g., Stutchbury et al. 2016), obligate migrants frequently engage in nonbreeding movements that do not fit neatly into the paradigm of tidy flights between breeding ranges and wintering ranges via discrete, short-term stopover sites (Newton 2012). A better understanding of the control of more flexible migration systems would inform how and why obligate migrants move in the way they do during these facultative phases of their migratory cycle.

**(2) Variation is the fodder of evolution.** Many hypotheses have been proposed for how long-distance, obligate migration arises in birds (Alerstam and Enckell 1979, Cox 1985, Lundberg 1988, Holmgren and Lundberg 1993). All share a common feature: individuals vary in their movement behavior, and selection acts on that variation. Altitudinal migration and other short-distance facultative migrations have been proposed to be precursors of long-distance migration in Neotropical lineages (Levey and Stiles 1992). Understanding the selective costs and benefits in obligate long-distance migrants requires that we adopt a comparative approach and make inferences regarding the migratory behavior of ancestral species (e.g., Outlaw and Voelker 2006). However, we know that migration is extremely labile; European Starlings (*Sturnus vulgaris*) in North America have altered their migratory propensity since being introduced to the continent in 1890 (Bitton and Graham 2015), and House Finches (*Haemorrhous mexicanus*) transported to eastern North America have “evolved” migratory behavior over the past 8 decades (Able and Belthoff 1998). Thus, inferences regarding the selective pressures that have tipped fitness cost–benefit trade-offs toward migrating may be obscured by saturation of gains and losses in phylogenetic contexts. By studying altitudinal migrants

and other partial, “messy” migrations, we can study ecological selective pressures in the here and now. In effect, such systems permit us to conduct empirical tests of hypotheses regarding the evolution of migration as it is happening.

**(3) Efficiency.** We live in an age of competitive job markets (Marshall et al. 2009) and scarce funding (<https://debblog.nsfbio.com/category/deb-numbers/>), factors that favor high research efficiency per unit time and dollar. Studying migration over large spatial scales frequently involves costly travel and technology. If such studies take place in wintering areas, they can also involve environmentally costly air travel and time-intensive acquisition of foreign research permits. In altitudinal migration systems, we can usually test the same hypotheses with fewer logistical challenges at the expense of fewer grant dollars relative to comparable studies with latitudinal migrants (e.g., Boyle 2008a vs. McKinnon et al. 2010). Such considerations are particularly relevant to graduate student research where careful choice of study system can permit full annual cycle field-based migration studies, even while completing academic requirements (e.g., Morrissey 1997, Lundblad 2014).

It is possible that the shorter-distance, facultative movements have been bypassed because overlapping summer and winter ranges make broad movement patterns harder to pin down. New tracking technologies are overcoming many of these logistical constraints as ever-smaller tracking devices come on the market, increasing the volume, precision, and accuracy of spatial data. Additionally, altitudinal gradients typically vary in isotopic signatures over short distances (Hobson 2005). By judicious choice of tissue sampling, detecting individual differences in altitudinal migration using isotopes is now possible in birds of any size at relatively low cost (Hobson et al. 2003, Boyle et al. 2011b).

**(4) Save the birds.** Just as we cannot understand the causes of population declines or implement effective conservation for long-distance migrants if we do not know where they go, we cannot ensure the future of the rest of our avifauna if we are ignorant of their movements. Declines in migratory birds are not limited to long-distance latitudinal migrants. In some cases, critically endangered birds undergo seasonal altitudinal migrations; understanding where, when, and how they move may prove crucial to protecting their future (Guevara et al. 2015). To date, no comparison exists of population trends of altitudinal migrants relative to nonmigrants or long-distance migrants. However, we do know that montane species are subject to unique challenges as climates change (La Sorte and Jetz 2010), and that the biology of montane birds is not as well understood as lowland species (Boyle and Martin 2015). Explicitly designing studies to allow detection and documentation of the patterns and prevalence of altitudinal

migration at community or population levels should be a priority in future ornithological studies.

## ACKNOWLEDGMENTS

I am indebted to the hundreds of authors who compiled and synthesized literature to create and update the *Birds of North America* series. Space precluded me from citing each of the life history accounts here. Many thanks to C. Conway for feedback on early stages of this project, and to C. Doty and S. Replogle Curnutt for assistance tabulating data and obtaining obscure references. WAB conceived of the idea, collected the data, and wrote the paper. This is contribution no. 17-139-J from the Kansas Agricultural Experiment Station.

## LITERATURE CITED

- Able, K. P., and J. R. Belthoff (1998). Rapid 'evolution' of migratory behaviour in the introduced House Finch of eastern North America. *Proceedings of the Royal Society B: Biological Sciences* 265:2063–2071.
- Alerstam, T., and P. H. Enckell (1979). Unpredictable habitats and evolution of bird migration. *Oikos* 33:228–232.
- Alves, M. A. S. (2007). Bird migration systems in Brazilian terrestrial habitats: Examples, gaps of knowledge, and directions for future studies. *Revista Brasileira De Ornitologia* 15:231–238.
- American Ornithologists' Union (1998). Checklist of North American Birds, 7th edition. American Ornithologists' Union, Washington, DC.
- Areta, J. I., and A. Bodrati (2010). A longitudinal migratory system within the Atlantic Forest: Seasonal movements and taxonomy of the Golden-rumped Euphonia (*Euphonia cyanocephala*) in Misiones (Argentina) and Paraguay. *Ornitología Neotropical* 21:71–86.
- Barve, S., A. A. Dhondt, V. B. Mathur, and Z. A. Cheviron (2016). Life-history characteristics influence physiological strategies to cope with hypoxia in Himalayan birds. *Proceedings of the Royal Society B: Biological Sciences* 283. doi:10.1098/rspb.2016.2201
- Beebe, W. (1947). Avian migration at Rancho Grande in north-central Venezuela. *Zoologica* 32:153–168.
- Bencke, G. A., and A. Kindel (1999). Bird counts along an altitudinal gradient of Atlantic forest in northeastern Rio Grande do Sul, Brazil. *Ararajuba* 7:91–107.
- Berthold, P. (1991). Genetic control of migratory behavior in birds. *Trends in Ecology & Evolution* 6:254–257.
- Berthold, P. (1998). Bird migration: Genetic programs with high adaptability. *Zoology-Analysis of Complex Systems* 101:235–245.
- Berthold, P. (2001). *Bird Migration: A General Survey*, 2nd edition. Oxford University Press, Oxford, UK.
- Bildstein, K. L. (2004). Raptor migration in the Neotropics: Patterns, processes, and consequences. *Ornitología Neotropical* 15:83–99.
- Bitton, P. P., and B. A. Graham (2015). Change in wing morphology of the European Starling during and after colonization of North America. *Journal of Zoology* 295:254–260.
- Borras, A., J. C. Senar, F. Alba-Sanchez, J. A. Lopez-Saez, J. Cabrera, X. Colome, and T. Cabrera (2010). Citril Finches during the winter: Patterns of distribution, the role of pines and implications for the conservation of the species. *Animal Biodiversity and Conservation* 33:89–115.
- Boyle, W. A. (2008a). Can variation in risk of nest predation explain altitudinal migration in tropical birds? *Oecologia* 155:397–403
- Boyle, W. A. (2008b). Partial migration in birds: Tests of three hypotheses in a tropical lekking frugivore. *Journal of Animal Ecology* 77:1122–1128.
- Boyle, W. A. (2011). Short-distance partial migration of Neotropical birds: A community-level test of the foraging limitation hypothesis. *Oikos* 120:1803–1816.
- Boyle, W. A., C. J. Conway, and J. L. Bronstein (2011a). Why do some, but not all, tropical birds migrate? A comparative study of diet breadth and preference *Evolutionary Ecology* 25:219–236.
- Boyle, W. A., C. G. Guglielmo, K. A. Hobson, and D. R. Norris (2011b). Lekking birds in a tropical forest forego sex for migration. *Biology Letters* 7:661–663.
- Boyle, W. A., and K. Martin (2015). The conservation value of high elevation habitats to migrant birds in southern British Columbia. *Biological Conservation* 192:461–476.
- Boyle, W. A., D. R. Norris, and C. G. Guglielmo (2010). Storms drive altitudinal migration in a tropical bird. *Proceedings of the Royal Society B: Biological Sciences* 277:2511–2519.
- Brambilla, M., and D. Rubolini (2009). Intra-seasonal changes in distribution and habitat associations of a multi-brooded bird species: Implications for conservation planning. *Animal Conservation* 12:71–77.
- Brennan, L. A., W. M. Block, and R. J. Gutiérrez (1987). Habitat use by Mountain Quail in northern California. *The Condor* 89:66–74.
- Brown, M. (2006). Annual and seasonal trends in avifaunal species richness in a coastal lowlands forest reserve in South Africa. *Ostrich* 77:58–66.
- Burgess, N. D., and C. O. F. Mlingwa (2000). Evidence for altitudinal migration of forest birds between montane Eastern Arc and lowland forests in East Africa. *Ostrich* 71:184–190.
- Burleigh, T. D. (1972). *Birds of Idaho*. Caxton Printers, Caldwell, ID, USA.
- Cade, B. S., and R. W. Hoffman (1993). Differential migration of Blue Grouse in Colorado. *The Auk* 110:70–77.
- Capllonch, P. (2007). Migraciones de especies de Tyrannidae de la Argentina: Parte 1. *Acta Zoologica Lilloana* 51:151–160.
- Chassot, O., and G. Monge-Arias (2012). Connectivity conservation of the Great Green Macaw's landscape in Costa Rica and Nicaragua (1994–2012). *Parks* 18:61–69.
- Chaves-Campos, J. (2004). Elevational movements of large frugivorous birds and temporal variation in abundance of fruits along an elevational gradient. *Ornitología Neotropical* 15:433–445.
- Chaves-Campos, J., J. E. Arévalo, and M. Araya (2003). Altitudinal movements and conservation of Bare-necked Umbrellabird *Cephalopterus glabricollis* of the Tilarán Mountains, Costa Rica. *Bird Conservation International* 13:45–58.
- Chesser, R. K., R. C. Banks, F. K. Barker, C. Cicero, J. L. Dunn, A. W. Kratter, I. J. Lovette, P. C. Rasmussen, J. V. Remsen, J. D. Rising, D. F. Stotz, and K. Winker (2013). Fifty-fourth

- supplement to the American Ornithologists' Union Check-list of North American Birds. *The Auk* 130:558–571.
- Christensen, G. C. (1954). The Chukar Partridge in Nevada. *Nevada Biological Bulletin* 1:77.
- Chu, M., and G. E. Walsberg. (1999). Phainopepla (*Phainopepla nitens*). In *The Birds of North America Online* (P. G. Rodewald, Ed.). Cornell Lab of Ornithology, Ithaca, NY, USA. doi:10.2173/bna.415
- Clark, C. J., and S. M. Russell (2012). Anna's Hummingbird (*Calypte anna*). In *The Birds of North America Online* (A. Poole, Editor). Cornell Lab of Ornithology, Ithaca, NY, USA. doi:10.2173/bna.226
- Connelly, J. W. (1988). Movements of Sage Grouse in South-eastern Idaho. *The Journal of Wildlife Management* 52:116–122.
- Cox, G. W. (1985). The evolution of avian migration systems between temperate and tropical regions of the New World. *The American Naturalist* 126:451–474.
- Crouch, J. E. (1943). Distribution and habitat relationships of the Phainopepla. *The Auk* 60:319–333.
- De La Hera, I., J. Gomez, T. Andres, P. Gonzalez-Ocio, P. Salmon, M. Salvador, A. Unanue, F. Zufiaur, and A. Onrubia (2014). Inferring the migratory status of woodland birds using ringing data: The case of a constant-effort site located in the Iberian highlands. *Ardeola* 61:77–95.
- Des Granges, J. L. (1979). Organization of a tropical nectar feeding bird guild in a variable environment. *Living Bird* 17: 199–236.
- Dingle, H. (2004). The Australo-Papuan bird migration system: Another consequence of Wallace's Line. *Emu* 104:95–108.
- Dingle, H. (2008). Bird migration in the southern hemisphere: A review comparing continents. *Emu* 108:341–359.
- Dixit, S., V. Joshi, and S. Barve (2016). Bird diversity of the Amrutganga Valley, Kedarnath, Uttarakhand, India with an emphasis on the elevational distribution of species. *Checklist* 12:1874.
- Dixon, K. L., and J. D. Gilbert (1964). Altitudinal migration in the Mountain Chickadee. *The Condor* 66:61–64.
- Dobzhansky, T. (1973). Nothing in biology makes sense except in the light of evolution. *The American Biology Teacher* 35:125–129.
- Dunning, J. B., and R. K. Bowers, Jr. (1984). Local movements of some Arizona montane birds. *North American Bird Bander* 9: 7.
- Ferrari, S., C. Albrueli, S. Imberti, and C. Lishman (2008). The up-to-date knowledge of the Magellanic Plover (*Pluvianellus socialis*), an endemic shorebird of southern Patagonia: Putting together the pieces of the puzzle. *Ornitología Neotropical* 19:433–443.
- Ford, H., and J. Pursey (1982). Status and feeding of the Eastern Spinebill *Acanthorhynchus tenuirostris* at the New England National Park, north-eastern NSW. *Emu* 82:203–211.
- Fraser, K. C., A. W. Diamond, and L. Chavarria (2010). Evidence of altitudinal moult-migration in a Central American hummingbird, *Amazilia cyanura*. *Journal of Tropical Ecology* 26:645–648.
- Fraser, K. C., T. K. Kyser, and L. M. Ratcliffe (2008). Detecting altitudinal migration events in Neotropical birds using stable isotopes. *Biotropica* 40:269–394.
- Fretwell, S. D. (1980). Evolution of migration in relation to factors regulating bird numbers. In *Migrant Birds in the Neotropics: Ecology, Behavior, Distribution and Conservation* (A. Keast and E. S. Morton, Editors). Smithsonian Institution Press, Washington, DC, USA. pp. 517–527.
- Galetti, M. (2001). Seasonal movements and diet of the Plumbeous Pigeon (*Columba plumbea*) in a Brazilian Atlantic Forest. *Melopsittacus* 4:39–43.
- Garwood, J. M., K. L. Pope, R. M. Bourque, and M. D. Larson (2009). High mountain lakes provide a seasonal niche for migrant American Dippers. *The Wilson Journal of Ornithology* 121:600–609.
- Gilbert, W. M., M. K. Sogge, and C. Van Riper III (2010). Orange-crowned Warbler (*Oreothlypis celata*). In *The Birds of North America* (A. Poole, Editor). Cornell Lab of Ornithology, Ithaca, NY, USA. doi:10.2173/bna.101
- Gill, F. B. (2007). *Ornithology*, 3rd edition. W. H. Freeman and Company, New York, NY, USA.
- Gillis, E. A., D. J. Green, H. A. Middleton, and C. A. Morrissey (2008). Life history correlates of alternative migratory strategies in American Dippers. *Ecology* 89:1687–1695.
- Grant, K. A., and V. Grant (1967). Effects of hummingbird migration on plant speciation in the California flora. *Evolution* 21:457–465.
- Green, D. J., I. B. J. Whitehorne, H. A. Middleton, and C. A. Morrissey (2015). Do American Dippers obtain a survival benefit from altitudinal migration? *PLOS One* 10:e0125734. doi:10.1371/journal.pone.0125734
- Greenberg, R. (1980). Demographic aspects of long-distance migration. In *Migrant Birds in the Neotropics: Ecology, Behavior, Distribution and Conservation* (A. Keast and E. S. Morton, Editors). Smithsonian Institution Press, Washington, DC, USA. pp. 493–504.
- Grinnell, J., and A. H. Miller (1944). The distribution of the birds of California. *Pacific Coast Avifauna* 27:1–608.
- Guevara, E. A., E. Bonaccorso, and J. F. Duivenvoorden (2015). Multi-scale habitat use analysis and interspecific ecology of the critically endangered Black-breasted Puffleg *Eriocnemis nigrivestis*. *Bird Conservation International* 25:479–488.
- Gutiérrez, R. J., and D. J. Delehanty (1999). Mountain Quail (*Oreortyx pictus*). In *The Birds of North America* (A. Poole and F. Gill, Editors). The Birds of North America, Inc., Philadelphia, PA, USA. doi:10.2173/bna.457
- Hahn, T. P., K. W. Sockman, C. W. Breuner, and M. L. Morton (2004). Facultative altitudinal movements by Mountain White-crowned Sparrows (*Zonotrichia leucophrys oriantha*) in the Sierra Nevada. *The Auk* 121:1269–1281.
- Hegemann, A., P. P. Marra, and B. I. Tieleman (2015). Causes and consequences of partial migration in a passerine bird. *The American Naturalist* 186:531–546.
- Hendricks, P., and J. E. Swenson (1983). Dynamics of the winter distribution of Rosy Finches, *Leucosticte arctoa*, in Montana. *Canadian Field-Naturalist* 97:307–310.
- Henshaw, H. W. (1876). Report on the Ornithology of the Portions of California Visited During the Field-Season of 1875. In *United States Geographical Surveys West of the One Hundredth Meridian Annual Reports*. United States Geological Survey, Washington, DC. pp. 224–278.
- Hess, S. C., C. R. Leopold, K. Misajon, D. Hu, and J. J. Jeffrey (2012). Restoration of movement patterns of the Hawaiian Goose. *The Wilson Journal of Ornithology* 124:478–486.
- Hilty, S. L. (1997). Seasonal distribution of birds at a cloud-forest locality, the Anchicayá Valley, in western Colombia. In *Studies*

- in Neotropical Ornithology Honoring Ted Parker (J. V. Remsen, Jr., Editor). Ornithological Monographs 48:321–343.
- Hobson, K. A. (2005). Stable isotopes and the determination of avian migratory connectivity and seasonal interactions. *The Auk* 122:1037–1048.
- Hobson, K. A., L. I. Wassenaar, B. Milá, I. Lovette, C. Dingle, and T. B. Smith (2003). Stable isotopes as indicators of altitudinal distributions and movements in an Ecuadorean hummingbird community. *Oecologia* 136:302–308.
- Holmgren, N., and S. Lundberg (1993). Despotic behavior and the evolution of migration patterns in birds. *Ornis Scandinavica* 24:103–109.
- Horvath, E. G., and K. A. Sullivan (1988). Facultative migration in Yellow-eyed Juncos. *The Condor* 90:482–484.
- Hughes, R. A. (1980). Additional puna zone bird species on the coast of Peru. *The Condor* 82:475.
- Hughes, R. A. (1984). Further notes on puna bird species on the coast of Peru. *The Condor* 86:93.
- Jahn, A. E., S. P. Bravo, V. R. Cueto, D. J. Levey, and M. V. Morales (2012). Patterns of partial avian migration in northern and southern temperate latitudes of the New World. *Emu* 112:17–22.
- Johnson, D. N., and G. L. Maclean (1994). Altitudinal migration in Natal. *Ostrich* 65:86–94.
- Johnson, R. E. (2002). Black Rosy-Finch (*Leucosticte atrata*). In *The Birds of North America* (A. Poole, Editor). Cornell Lab of Ornithology, Ithaca, NY, USA. doi:10.2173/bna.678
- Ketterson, E. D., and V. Nolan, Jr. (1983). The evolution of differential bird migration. *Current Ornithology* 1:357–402.
- Kimura, K., T. Yumoto, and K. Kikuzawa (2001). Fruiting phenology of fleshy-fruited plants and seasonal dynamics of frugivorous birds in four vegetation zones on Mt. Kinabalu, Borneo. *Journal of Tropical Ecology* 17:833–858.
- King, J. R., and E. E. Wales (1964). Observations on migration, ecology, and population flux of wintering Rosy Finches. *The Condor* 66:24–31.
- Klemp, S. (2003). Altitudinal dispersal within the breeding season in the Grey Wagtail *Motacilla cinerea*. *Ibis* 145:509–511.
- Kondo, B., and K. E. Omland (2007). Ancestral state reconstruction of migration: Multistate analysis reveals rapid changes in New World orioles (*Icterus* spp.). *The Auk* 124:410–419.
- Körner, C. (2007). The use of 'altitude' in ecological research. *Trends in Ecology & Evolution* 22:569–574.
- La Sorte, F. A., and W. Jetz (2010). Projected range contractions of montane biodiversity under global warming. *Proceedings of the Royal Society B: Biological Sciences* 277:3401–3410.
- Laymon, S. A. (1989). Altitudinal migration movements of Spotted Owls in the Sierra Nevada, California. *The Condor* 91:837–841.
- Leopold, C. R., and S. C. Hess (2014). Corridor- and stopover-use of the Hawaiian Goose (*Branta sandvicensis*), an intratropical altitudinal migrant. *Journal of Tropical Ecology* 30:67–78.
- Leu, M., and C. W. Thompson (2002). The potential importance of migratory stopover sites as flight feather molt staging areas: A review for Neotropical migrants. *Biological Conservation* 106:45–56.
- Levey, D. J., and F. G. Stiles (1992). Evolutionary precursors of long-distance migration: Resource availability and movement patterns in Neotropical landbirds. *The American Naturalist* 140:447–476.
- Loiselle, B. A., and J. G. Blake (1991). Temporal variation in birds and fruits along an elevational gradient in Costa Rica. *Ecology* 72:180–193.
- Lorenz, T. J., and K. A. Sullivan (2009). Seasonal differences in space use by Clark's Nutcrackers in the Cascade Range. *The Condor* 111:326–340.
- Lu, X., C. Wang, and T. Yu (2010). Nesting ecology of the Grey-backed Shrike (*Lanius tephronotus*) in South Tibet. *The Wilson Journal of Ornithology* 122:395–398.
- Lundberg, P. (1988). The evolution of partial migration in birds. *Trends in Ecology & Evolution* 3:172–175.
- Lundblad, C. G. 2014. Partial altitudinal migration: Tests of four mechanistic hypotheses in Yellow-eyed Juncos (*Junco phaeonotus*). M.S. thesis, University of Arizona, Tucson, AZ, USA.
- Mackas, R. H., D. J. Green, I. B. J. Whitehorne, E. N. Fairhurst, H. A. Middleton, and C. A. Morrissey (2010). Altitudinal migration in American Dippers (*Cinclus mexicanus*): Do migrants produce higher quality offspring? *Canadian Journal of Zoology* 88: 369–377.
- Marra, P. P., K. A. Hobson, and R. T. Holmes (1998). Linking winter and summer events in a migratory bird by using stable-carbon isotopes. *Science* 282:1884–1886.
- Marshall, J. C., P. Buttars, T. Callahan, J. J. Dennehy, D. J. Harris, B. Lunt, M. Mika, and R. Shupe (2009). In the academic job market, will you be competitive? A case study in ecology and evolutionary biology. *Israel Journal of Ecology & Evolution* 55: 381–389.
- Martínez del Río, C., P. Sabat, R. Anderson-Sprecher, and S. P. Gonzalez (2009). Dietary and isotopic specialization: The isotopic niche of three *Cinclodes* ovenbirds. *Oecologia* 161: 149–159.
- McGowan, P. J. K., and G. M. Kirwan (2016). Chukar (*Alectoris chukar*). In *Handbook of the Birds of the World Alive* (J. del Hoyo, A. Elliott, J. Sargatal, D. A. Christie, and E. de Juana, Editors). Lynx Editions, Barcelona, Spain.
- McGuire, L. P., and W. A. Boyle (2013). Altitudinal migration in bats: Evidence, patterns, and processes. *Biological Reviews* 88:767–786.
- McKinnon, L., P. A. Smith, E. Nol, J. L. Martin, F. I. Doyle, K. F. Abraham, H. G. Gilchrist, R. I. G. Morrison, and J. Bêty (2010). Lower predation risk for migratory birds at high latitudes. *Science* 327:327–328.
- Middleton, H. A., C. A. Morrissey, and D. J. Green (2006). Breeding territory fidelity in a partial migrant, the American Dipper *Cinclus mexicanus*. *Journal of Avian Biology* 37:169–178.
- Morrissey, C. A. 1997. The ecological and toxicological significance of altitudinal migration by the American Dipper (*Cinclus mexicanus*). Ph.D. dissertation, Simon Fraser University, Vancouver, BC, Canada.
- Newsome, S. D., P. Sabat, N. Wolf, J. A. Rader, and C. M. del Rio (2015). Multi-tissue  $\delta^2\text{H}$  analysis reveals altitudinal migration and tissue-specific discrimination patterns in *Cinclodes*. *Ecosphere* 6:art213.
- Newton, I. (2012). Obligate and facultative migration in birds: Ecological aspects. *Journal of Ornithology* 153:5171–5180.
- Nilsson, A. L. K., T. Alerstam, and J. A. Nilsson (2008). Diffuse, short and slow migration among Blue Tits. *Journal of Ornithology* 149:365–373.
- Nocedal, J. (1994). Local migrations of insectivorous birds in western Mexico: Implications for the protection and conser-

- vation of their habitats. *Bird Conservation International* 4: 129–142.
- Norbu, N., M. Wikelski, D. S. Wilcove, J. Partecke, Ugyen, U. Tenzin, Sherub, and T. Tempa (2013). Partial altitudinal migration of a Himalayan forest Pheasant. *PLOS One* 8: e60979. doi:10.1371/journal.pone.0060979
- Norment, C. J., and K. Green (2004). Breeding ecology of Richard's Pipit (*Anthus novaeseelandiae*) in the Snowy Mountains. *Emu* 104:327–336.
- Ogonowski, M. S., and C. J. Conway (2009). Migratory decisions in birds: Extent of genetic versus environmental control. *Oecologia* 161:199–207.
- O'Neill, J. P., and T. A. Parker III (1978). Responses of birds to a snow-storm in the Andes of southern Peru. *The Wilson Bulletin* 90:446–449.
- Ornelas, J. F., and M. d. C. Arizmendi (1995). Altitudinal migration: Implications for the conservation of the Neotropical migrant avifauna of western Mexico. In *Conservation of Neotropical Migratory Birds in Mexico* (M. H. Wilson and A. Sader, Editors). Maine Agricultural and Forest Experiment Station, Miscellaneous Publications 727, Orono, ME. pp. 98–109.
- Outlaw, D. C., and G. Voelker (2006). Phylogenetic tests of hypotheses for the evolution of avian migration: A case study using the Motacillidae. *The Auk* 123:455–466.
- Powell, G. V. N., and R. D. Bjork (2004). Habitat linkages and the conservation of tropical biodiversity as indicated by seasonal migrations of Three-wattled Bellbirds. *Conservation Biology* 18:500–509.
- Presnall, C. C. (1935). Altitudinal migration in southern Utah. *The Condor* 37:37–38.
- Rabenold, K. N., and P. P. Rabenold (1985). Variation in altitudinal migration, winter segregation, and site-tenacity in two subspecies of Dark-eyed Juncos in the southern Appalachians. *The Auk* 102:805–819.
- Ramenofsky, M., J. Cornelius, and B. Helm (2012). Physiological and behavioral responses of migrants to environmental cues. *Journal of Ornithology* 153:181–191.
- Ramos-Olmos, M. A. 1983. Seasonal movements of bird populations at a Neotropical study site in southern Veracruz, Mexico. Ph.D. dissertation, University of Minnesota, Minneapolis, MN, USA.
- Richards, E. B. (1924). A list of the land birds of the Grass Valley District, California. *The Condor* 26:98–104.
- Rodewald, P. E. (2015). *The Birds of North America*. Cornell Laboratory of Ornithology, Ithaca, NY, USA. <https://birdsna.org>
- Rohwer, S., and J. Manning (1990). Differences in timing and number of molts for Baltimore and Bullock's Orioles: Implications to hybrid fitness and theories of delayed plumage maturation. *The Condor* 92:125–140.
- Rosene, W. (1969). *The Bobwhite Quail, Its Life and Management*. Rutgers University Press, New Brunswick, NJ, USA.
- Rosselli, L. (1994). The annual cycle of the White-ruffed Manakin, *Corapipo leucorrhoa*, a tropical frugivorous altitudinal migrant, and its food plants. *Bird Conservation International* 4: 143–160.
- Ryan, G. C. 2012. Cross-seasonal habitat selection of altitudinal migratory birds in Taiwan. M.Sc. thesis, National Dong Hwa University, Hualien, Taiwan, ROC.
- Schroeder, M. A., J. R. Young, and C. E. Braun (1999). Greater Sage-Grouse (*Centrocercus urophasianus*). In *The Birds of North America* (A. Poole, Editor). Cornell Lab of Ornithology, Ithaca, NY, USA. doi:10.2173/bna.425
- Seifert, N., M. Haase, S. L. Van Wilgenburg, C. C. Voigt, and A. S. Ornés (2016). Complex migration and breeding strategies in an elusive bird species illuminated by genetic and isotopic markers. *Journal of Avian Biology* 47:275–287.
- Sick, H. (1993). *Ornitología Brasileira*. Editora Nova Fronteira, Rio de Janeiro, Brazil.
- Simpson, M. B., and H. D. Pratt (1992). *Birds of the Blue Ridge Mountains: A Guide for the Blue Ridge Parkway, Great Smoky Mountains, Shenandoah National Park, and Neighboring Areas*. University of North Carolina Press, Chapel Hill, NC, USA.
- Skutch, A. F. (1969). *Life Histories of Central American Birds III: Families Cotingidae, Pipridae, Formicariidae, Furnariidae, Dendrocolaptidae, and Picidae*. *Pacific Coast Avifauna* 35:1–580.
- Slud, P. (1964). The birds of Costa Rica: Distribution and ecology. *Bulletin of the American Museum of Natural History* 128:1–430.
- Solórzano, S., S. Castillo, T. Valverde, and L. Ávila (2000). Quetzal abundance in relation to fruit availability in a cloud forest of southeastern Mexico. *Biotropica* 32:523–532.
- Sommeille, M., A. Manica, S. H. M. Butchart, and A. S. L. Rodrigues (2013). Mapping global diversity patterns for migratory birds. *PLOS One* 8:e70907. doi:10.1371/journal.pone.0070907
- Stanley, C. Q., M. MacPherson, K. C. Fraser, E. A. McKinnon, and B. J. M. Stutchbury (2012). Repeat tracking of individual songbirds reveals consistent migration timing but flexibility in route. *PLOS One* 7:e40688. doi:10.1371/journal.pone.0040688
- Steenhof, K. (2013). Prairie Falcon (*Falco mexicanus*). In *The Birds of North America* (A. Poole, Editor). Cornell Lab of Ornithology, Ithaca, NY, USA.
- Stiles, F. G. (1980). The annual cycle in a tropical wet forest hummingbird community. *Ibis* 122:322–343.
- Stiles, F. G. (1988). Altitudinal movements of birds on the Caribbean slope of Costa Rica: Implications for conservation. In *Tropical Rainforests: Diversity and Conservation* (F. Alameda and C. M. Pringle, Editors). California Academy of Sciences, San Francisco, CA, USA. pp. 243–258.
- Stiles, F. G., and A. F. Skutch (1989). *A Guide to the Birds of Costa Rica*. Cornell University Press, Ithaca, NY, USA.
- Strewe, R. (2006). The importance of altitudinal migration in frugivorous birds on the Pacific slope of the Colombian Andes. *Journal of Ornithology* 147:27.
- Strewe, R., and C. Navarro (2003). New distributional records and conservation importance of the San Salvador Valley, Sierra Nevada de Santa Marta, northern Colombia. *Ornitología Colombiana* 1:28–40.
- Strong, C., B. Zuckerberg, J. L. Betancourt, and W. D. Koenig (2015). Climatic dipoles drive two principal modes of North American boreal bird irruption. *Proceedings of the National Academy of Sciences USA* 112:E2795–E2802.
- Stutchbury, B. J. M., R. Siddiqui, K. Applegate, G. T. Hvenegaard, P. Mammenga, N. Mickle, M. Pearman, J. D. Ray, A. Savage, T. Shaheen, and K. C. Fraser (2016). Ecological causes and consequences of intratropical migration in temperate-breeding migratory birds. *The American Naturalist* 188:528–540.

- Swanson, D. L., J. L. Ingold, and R. Galati (2012). Golden-crowned Kinglet (*Regulus satrapa*). In *The Birds of North America* (P. E. Rodewald, Editor). Cornell Lab of Ornithology, Ithaca, NY, USA.
- Swarth, H. S. (1904). Birds of the Huachuca Mountains, Arizona. *Pacific Coast Avifauna* 4:1–73.
- Terrill, S. B., and K. P. Able (1988). Bird migration terminology. *The Auk* 105:205–206.
- Tidemann, S. C., S. J. Wilson, and T. G. Marples (1988). Some results from a long-term banding project in the Brindabella Range, A.C.T. *Corella* 12:1–6.
- Tinoco, B. A., P. X. Astudillo, S. C. Latta, and C. H. Graham (2009). Distribution, ecology and conservation of an endangered Andean hummingbird: The Violet-throated Metaltail (*Metalura baroni*). *Bird Conservation International* 19:63–76.
- Tomback, D. F. (1998). Clark's Nutcracker (*Nucifraga columbiana*). In *The Birds of North America* (A. Poole, Editor). Cornell Lab of Ornithology, Ithaca, NY, USA.
- Walsberg, G. E. (1977). Ecology and energetics of contrasting social systems in Phainopepla (Aves: Ptilonotidae). *University of California Publications in Zoology* 108:1–63.
- Watanabe, Y. Y. (2016). Flight mode affects allometry of migration range in birds. *Ecology Letters* 19:907–914.
- Wheelwright, N. T. (1983). Fruits and the ecology of Resplendent Quetzals. *The Auk* 100:286–301.
- Ydenberg, R. C., R. W. Butler, D. B. Lank, C. G. Guglielmo, M. Lemon, and N. Wolf (2002). Trade-offs, condition dependence and stopover site selection by migrating sandpipers. *Journal of Avian Biology* 33:47–55.