

Responses of foraging Eurasian beavers *Castor fiber* to predator odours

Authors: Rosell, Frank, and Czech, Andrzej

Source: *Wildlife Biology*, 6(1) : 13-21

Published By: Nordic Board for Wildlife Research

URL: <https://doi.org/10.2981/wlb.2000.033>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Responses of foraging Eurasian beavers *Castor fiber* to predator odours

Frank Rosell & Andrzej Czech

Rosell, F. & Czech, A. 2000: Responses of foraging Eurasian beavers *Castor fiber* to predator odours. - Wildl. Biol. 6: 13-21.

The ability of Eurasian beavers *Castor fiber* to recognise different predator odours has received little research, nor has the use of predator odours to deter Eurasian beavers from damaging agricultural crops, fruit and forest trees. Recognition of and response to predator odours by prey is of adaptive significance because it reduces predation risk. We tested the hypothesis that predator odours decrease foraging and predicted that: human and wolf *Canis lupus* odour would decrease foraging more effectively than other predator odours. Our results showed that all tested predator odours (red fox *Vulpes vulpes*, river otter *Lutra lutra*, lynx *Lynx lynx*, wolf and brown bear *Ursus arctos*), except those from human and dog *Canis familiaris*, significantly decreased foraging during summer. River otter, red fox, lynx, wolf and brown bear odours had the strongest effects during summer. During autumn, river otter odour was significantly more effective than the other predator odours, except those from lynx, human and red fox, in decreasing foraging. Only odour from river otter, human, lynx and red fox had a significantly stronger effect than the three controls during autumn. Overall, the river otter odour was most effective in decreasing foraging. Odours from predators sympatric with the Eurasian beaver did not have a larger effect than those of originally sympatric, but now absent species. Beavers ate more sticks with predator odour in autumn than in summer. Our results have clear practical implications, and several are suggested.

Key words: *Castor fiber*, Eurasian beaver, feeding repellents, foraging, odours, Poland, predators

Frank Rosell, Faculty of Arts and Sciences, Department of Environmental and Health Studies, Telemark College, N-3800 Bø in Telemark, Norway & Department of Zoology, Norwegian University of Science and Technology, N-7491 Trondheim, Norway - e-mail: Frank.Rosell@hit.no
Andrzej Czech, Institute of Environmental Sciences, Jagiellonian University, ul. Oleandry 2a, PL-30-063 Krakow, Poland

Received 16 April 1999, accepted 12 November 1999

Associate Editor: Tommy Asferg

The Eurasian beaver *Castor fiber* is slowly becoming re-established over much of its former range in Europe, and presently numbers are estimated at roughly half a million in Europe and Asia (Rosell & Pedersen 1999). The spread of beavers into cultural landscapes leads to conflict with man. In several countries the populations

of beavers are large enough to be harvested, and hunting is again allowed. In other countries where hunting is not permitted, other measures to control the population and limit damage by beavers need to be considered (Nolet & Rosell 1998).

In Eurasia nearly all beaver-man conflicts are caused

by beavers feeding on cultivated plants such as crops and trees, or dam building (Richard 1986, Heidecke & Klenner-Fringes 1992, Rosell & Parker 1995). Dam building is by far the most important beaver damage problem although feeding damage can locally be problematic (Rosell & Parker 1995). Most of the feeding damage (>75%) is reported from within a distance of 20 m from the water's edge (Heidecke & Klenner-Fringes 1992). Nolet & Rosell (1998) suggested that restoration of at least 20 m wide zones of natural vegetation along the banks of waterways (not accessible to cattle and horses) is the best durable solution to the problem of beaver feeding damage. Alternatively, feeding damage can be reduced by chemical repellents (slaked lime, quick lime and linseed oil), fencing or by using wire around individual trees (e.g. Richard 1986).

The Eurasian beaver is territorial and usually lives in family units (Wilsson 1971, Djoshkin & Safonow 1972, Nolet & Rosell 1994). A typical family consists of a monogamous adult pair, young of the year, yearlings and sometimes two-year olds or older (e.g. Wilsson 1971, Kudrjasov 1973). The territories are scent marked with castoreum and/or anal gland secretion, which play an important role in territory defence (Rosell & Nolet 1997, Rosell & Bergan 1998, Rosell, Bergan & Parker 1998). Beavers are strict herbivores. In summer, they mainly feed on aquatic plants and various herbs, and to a lesser extent woody food (e.g. Heidecke 1988, Histøl 1989, Nolet, van der Veer, Evers & Ottenheim 1995). In autumn and winter, they mainly eat woody food, although in some areas aquatics may also be used (Simonsen 1973). Although beavers can cut down mature trees of up to 1 m in diameter, they prefer small trees with a diameter of less than 10 cm (Wilsson 1971, Simonsen 1973, Basey & Jenkins 1995, Donkor & Fryxell 1999). Most trees are cut in autumn, when the beavers are preparing their food cache. Such a cache is mainly built in climates with harsh winters (Rosell & Parker 1995).

The beaver's aquatic lifestyle and habit of constructing partially submerged dens under masses of earth and sticks (Wilsson 1971, Zurowski 1992) has seemingly limited the number of its effective predators (Rosell & Parker 1996). The main cause for the near disappearance of the beaver in the 19th century was over-hunting (Nolet & Rosell 1998). Other species known to prey on beavers in Europe are wolves *Canis lupus*, bears *Ursus arctos*, lynxes *Lynx lynx*, wolverines *Gulo gulo*, red foxes *Vulpes vulpes* and dogs *Canis familiaris* (Dezhkin & Safonov 1966, Tyurnin 1984, Zurowski 1989, Kile, Nakken, Rosell & Espe-

land 1996, Rosell, Parker & Kile 1996). Humans and wolves are regarded as the beaver's main predators in Europe. Humans have preyed upon Eurasian beavers both in historic and presumably prehistoric time, and human predation was the major cause for the near extirpation of the beaver in Europe (Nolet 1996, Nolet & Rosell 1998, Rosell et al. 1996). In Isle Royale National Park wolf predation is so common that it is considered to regulate the local North American beaver *C. canadensis* population (Shelton & Peterson 1983). Recently, Andersone (1998) reported that beavers appeared to be the most important food item for wolves in Latvia during summer, and concluded that wolves switched to beavers when ungulates were few. Substantial predation by the black bear *U. americanus* on beaver has only been reported from an island in Lake Superior (Smith, Trauba & Anderson 1994). The lynx, wolverine, red fox and dog are rarely reported as predators on beavers and probably have minor effects on beaver populations. The river otter *Lutra lutra* and pine marten *Martes martes* are also suspected to prey on the Eurasian beaver in some areas (see references in Tyurnin 1984, Rosell & Hovde 1998). However, in many areas where the Eurasian beaver and the river otter are sympatric beaver remains have not been found in otter excrements (see references in Tyurnin 1984). Also, North American beavers are rare in the diet of the river otter *Lontra canadensis*. In a northeast Alberta area with high beaver and otter densities, Reid (1984) found that only 5 of 1,140 (0.4%) otter scats contained beaver remains. However, the otter may have fed on beaver carcasses. No beaver remains were found in seven other studies of otter droppings, cited by Reid (1984). Studies on the otter diet, undertaken in beaver lodges in Canada, also showed that otters did not prey on young beavers (Tumilson & Karnes 1987). The mink *Mustela vison* is probably not a predator of Eurasian beaver. Brzezinski & Zurowski (1992) found that young beavers were not eaten by mink during spring in northern Poland. Beaver kits were probably sufficiently guarded by their parents and mink were not able to enter the beaver lodges without exposing themselves to attacks by adults.

In North America, predator odours, especially of the coyote *C. latrans*, lynx *L. canadensis* and river otter, are promising as feeding repellents for the North American beaver (Engelhart & Müller-Schwarze 1995). Although mammals respond to scent from allopatric predators and often are repelled by them, chemical cues from sympatric species are generally more effec-

tive (Mason, Epple & Nolte 1993). Predator odours also have the potential to reduce feeding in many other genera of rodents (e.g. Vernet-Maury, Polak & Demael 1984, Robinson 1990, Sullivan, Crump, Wieser & Dixon 1990, Merckens, Harestad & Sullivan 1991, Swihart 1991, Nolte, Mason, Epple, Aronov & Campbell 1994, Borowski 1998). Recognition of and response to predator odours by prey is of adaptive significance because it reduces predation risk (Merckens et al. 1991).

The Eurasian beaver's ability to recognise different predator odours has apparently received no attention, nor has the use of predator odours to deter Eurasian beaver from damaging agricultural crops, fruit and forest trees. Therefore, we designed a study to test the hypothesis that predator odours decrease beaver foraging by investigating the effects of such scents on food consumption in the field. The following prediction was tested: human and wolf odour would be more effective in reducing beaver foraging than the other predator odours.

Material and methods

Our study was conducted in Suwalki Lakeland in northeastern Poland (53°50'N, 23°15'E). Gently rolling hills up to 220 m a.s.l., intersected by lakes and rivers form the landscape. Approximately 3.4% of the area is covered by water (Zurowski & Kasperczyk 1986). The natural vegetation on the banks of the watersheds consists mostly of shrubs of willow *Salix* spp. and aspen *Populus tremula*. Suwalki Lakeland is densely forested and human population is relatively low (about 10 inhabitants/km²; Rocznik Statystyczny 1997). World War II, and a resulting shift of national boundaries, caused the complete extirpation of Eurasian beavers from Poland by 1945, and the species is now strictly protected (Dzieciolowski & Gozdziowski 1999). The mean density of sites on the entire beaver range in the Suwalki Lakeland in 1984/85 was 15.3/100 km² (Zurowski & Kasperczyk 1986). The population in Suwalki Lakeland was about 4,500 in 1994 (Dzieciolowski 1996). In our study area, river otter (Brzezinski, Romanowski, Cygan & Pabin 1996), wolves (Okarma 1993), red foxes (Buchalczyk 1981, 1983, Pielowski 1982) and lynxes (Kamieniarz & Panek 1996) are present, while the brown bear occur only in the Carpathian Mountains in southern Poland (Jakubiec & Buchalczyk 1987). Information on month-

ly average rainfall was supplied by the closest meteorological station located about 40 km from the centre of the research area (Rocznik Statystyczny 1998). There was only a small difference in monthly average rainfall between the summer (51.5 mm) and autumn (63.5 mm) trials.

To compare our results from the Eurasian beaver with those from the North American beaver, we used the methods of Engelhart & Müller-Schwarze (1995). Predator odour samples were prepared from excrements (see Table 1). All faeces, except those from dogs, were collected from animals in Zoo Lodz, Zoo Poznan and Zoo Warsaw during April-June 1997. The dog samples were collected from pets. We assume that the faeces from the zoo animals were similar enough to those of wild animals because the zoo animals were fed with the same type of food as the animals eat in the wild. The river otters were fed with fish and frogs, the red foxes and the lynxes with rabbit meat, and the wolves and the brown bears with beef and pork. For all samples, one part faeces (by weight) was suspended in two parts pure methanol for two hours and then filtered with a vacuum filter. The filtrate was used for the experiment. The odour samples were stored in glass jars with alcohol-proof plastic lids in a refrigerator until use.

Aspen sticks cut from aspen saplings in the study area were used as carriers for the odours. Stems with a diameter of 1-2 cm and with as few lateral twigs as possible, were selected. The sticks were cut fresh on the first day of each trial, pruned, and cut into 30 cm long sections. The bark of all sticks, except the 'intact' control, was perforated by rolling them on a nail-board, to facilitate uptake of the methanol solutions. The sticks were dried at room temperature for two hours before being scented by dipping two thirds of their length into the extracts for two seconds (except those left intact). One stick was perforated and untreated (blank), and one unperforated and untreated (intact). In addition, sticks were soaked in a deer repellent based on human sweat named HUKINOL and produced by KIEFERLE GmbH, Germany. This repellent is used to protect forest plantations (P. Janiszewski, pers. comm.). We also soaked sticks in pure methanol as a solvent control. The sticks were dried for two hours at room temperature before further handling.

For each trial we used 10 randomly selected beaver colonies. We used different colonies for each of five consecutive trials, i.e. 50 colonies per season. The colonies used in the summer trials (16 July - 9 August 1997) were used again in the autumn trials (11 Octo-

ber - 4 November 1997). A trial consisted of 10 sticks placed in a row on the ground, 30 cm apart, at the bank of each pond parallel to the water's edge, near feeding places that were frequently visited by beavers. Each stick carried one of the seven predator odours, and the remaining three sticks were controls. The sticks were placed in a random order, determined by lot, and the same pattern was used for all colonies during one trial. A trial lasted for five days and nights, i.e. five beaver activity periods, without replacing any missing or altered sticks. New sticks were always used on a new trial. We used a total number of 1,000 sticks during the summer and autumn trials.

We recorded the results of nocturnal beaver activity each day between 7 and 9 a.m., and classified the sticks into: 1) eaten, i.e. completely or peeled $\geq 80\%$, or missing sticks, and 2) not eaten, i.e. peeled $< 80\%$, left intact and in place or left intact and moved. Beavers sample sticks and often discard partially peeled sticks (Engelhart & Müller-Schwarze 1995). Sticks that were not eaten after five activity periods were regarded as rejected.

We used χ^2 -tests to test for overall significance in the differences between odours in number of sticks eaten after five activity periods during both summer and autumn trials (Sokal & Rohlf 1995). Thereafter we tested for which odours were significantly different using Tukeys T-method for multiple comparisons (approximate) (Berenson & Levine 1989). Sign tests were used to test if the median number of sticks eaten during the summer were less than the median number of sticks eaten during the autumn. All probabilities presented are one-tailed unless otherwise indicated (Sokal & Rohlf 1995).

Results

Summer

The number of experimental sticks eaten by the beaver (Fig. 1) differed significantly between odours ($\chi^2 = 257.6$, $P < 0.0001$). After five activity periods, it ranged from 0% for river otter and red fox odours to 90% for blank and intact aspen sticks. After river otter and red fox, the most repellent odours were

lynx (6% eaten), wolf (8% eaten), and brown bear (10% eaten). Among the responses of beavers to the different samples, significant differences were found in the following comparisons: all odours except blank vs intact, and all odours, except human and dog vs methanol control (Table 1). Generally, there were no significant differences between the single predator odours. All predator odours, except human and dog odours, had a significantly stronger effect on beaver foraging than the three controls.

Autumn

In the autumn, the number of experimental sticks eaten by the beaver (see Fig. 1) also differed significantly between odours ($\chi^2 = 139.7$, $P < 0.0001$). As in summer, river otter odour was avoided most, but not quite as pronounced as in summer (0% vs 18% eaten). The most frequently eaten samples were blank and intact (98% each). Generally, river otter odour was significantly different from all other predator odours, except human odours (26% eaten), lynx (42% eaten) and red fox (46% eaten) (see Table 1).




Figure 1. Responses by beaver (% samples completely eaten or peeled $\geq 80\%$, and/or missing) to different odours and predator odours at Suwalki Lakeland in Poland during summer and autumn, 1997. The activity period lasted for five consecutive days and nights (1-5), without replacing any missing or altered sticks.

Table 1. Difference in beaver responses to odour samples after five activity periods for summer and autumn trials during 1997.

| Odour | RO ^a | Lynx | Wolf | BB ^a | Dog | HS ^a | M ^a | I ^a | B ^a |
|------------------|-----------------|------|------|-----------------|-----|-----------------|----------------|----------------|----------------|
| Summer trials | | | | | | | | | |
| Red fox | - | - | - | - | - | - | * | * | * |
| River otter | | - | - | - | - | - | * | * | * |
| Lynx | | | - | - | - | - | * | * | * |
| Wolf | | | | - | - | - | * | * | * |
| Brown bear | | | | | - | - | * | * | * |
| Dog | | | | | | - | - | * | * |
| Human sweat | | | | | | - | - | * | * |
| Methanol control | | | | | | | | * | * |
| Intact | | | | | | | | | - |
| Autumn trials | | | | | | | | | |
| Red fox | - | - | - | - | - | - | * | * | * |
| River otter | | - | * | * | * | - | * | * | * |
| Lynx | | | - | - | - | - | * | * | * |
| Wolf | | | | - | - | * | - | * | * |
| Brown bear | | | | | - | - | - | * | * |
| Dog | | | | | | * | - | - | - |
| Human sweat | | | | | | | * | * | * |
| Methanol control | | | | | | | | - | - |
| Intact | | | | | | | | | - |

^a Abbreviations for odours: RO: river otter; BB: brown bear; HS: human sweat; M: methanol control; I: intact; B: blank.

* and - state if the comparison is significant or not by Tukey's T-method for multiple comparisons (approximate) at total significance level 5%.

Seasonal variation

Only 10.3% of sticks with predator odours were eaten in summer, compared with 44.9% in autumn. This difference was significant (two-tailed, $S = 7$ positive differences, $P = 0.0156$, $N = 7$). In summer trials, 23.3% of all sticks were consumed, compared with 49.0% in autumn. This difference is also significant (two-tailed, $S = 10$ positive differences, $P = 0.002$, $N = 10$). Only sticks with human odour were eaten in similar amount during the summer and autumn trials (22 and 26%, respectively).

Discussion

Significance of predator odours in influencing foraging

Our results supported the hypothesis that predator odours on aspen sticks decrease foraging by the Eurasian beaver. The results also showed that Eurasian beavers are able to distinguish between different odour treatments of food. We predicted that human and wolf odours would be most effective. However, the results showed that during summer all predator odours, except those of human and dog, significantly decreased beaver foraging. River otter, red fox, lynx, wolf and brown bear odours had the strongest effects during summer. During autumn, river otter odour was significantly more effective than the other predator

odours, except those of human, lynx and red fox, in decreasing beaver foraging. Only river otter, lynx, human and red fox odours had a significantly stronger effect than the three controls during autumn. Similar results were found by Engelhart & Müller-Schwarze (1995) for the North American beaver. They showed that coyote, lynx and river otter odours had the strongest effects and concluded that predator odours were promising as feeding repellents.

The effectiveness of predator odours as natural repellents may depend on factors such as geographic distribution of predator and prey, duration of their geographic association and cultural transmission of predator responses among prey (Swihart 1991). An innate response by prey to a predator cue such as odour is likely to result if prey and predator have coexisted over evolutionary time (e.g. Gorman 1984, Berdoy & Macdonald 1991, Ward, Macdonald & Doncaster 1997). However, research done with house mice *Mus musculus* conducted on an island without any mammalian predators showed that reaction to predator odour (cat *Felis catus* and red fox) may disappear after a certain number of generations (Dickman 1992), and therefore is not passed on genetically. Avoidance of a predator odour can be either species-specific (Swihart 1991, Jedrzejewski, Rychlik & Jedrzejewska 1993, Nolte, Farley, Campbell, Mason & Epple 1993) or general (Stoddart 1982a, Weldon, Graham & Mears 1993, Nolte et al. 1994). Furthermore, some prey learn to respond only to predators that are actively

dangerous (Dickman 1992). The dog had the weakest effect during both summer and autumn trials. This finding may be due to the novelty of the pet dogs' odour. Indeed, no wild dogs are present in our study area. Interactions between beaver and humans have occurred in Poland for a very long time before the protection. Fossils of Eurasian beavers appear in early Pliocene deposits (12 million years ago) in Europe (Novak 1987). The duration of the geographic association between the different predators and the Eurasian beaver in our area, and/or the past or present predator pressure probably explains the results we found. Engelhart & Müller-Schwarze (1995) found that the slightly more avoided odours were those of predator species that were sympatric with the respective North American beaver populations. Beavers avoided the odours of sympatric predators such as river otter and coyote that actually occur within the range of the tested beaver slightly more than those of originally sympatric, but now absent, species such as the wolf. However, the North American beaver did not avoid the odours of sympatric species significantly more than the odour of the allopatric species, i.e. African lion *Panthera leo* (Engelhart & Müller-Schwarze 1995). Our results did not show that odours from predators sympatric with the Eurasian beaver, like wolf and lynx, had a significantly larger effect than those of originally sympatric, but now absent, species such as the brown bear. Responses to chemical cues from allopatric predators and failure to habituate to predator odours have been interpreted as evidence that responses to these stimuli are innate (Müller-Schwarze 1972, Stoddart 1980, Dickman & Doncaster 1984, Robinson 1990). Dickman & Doncaster (1984) suggested that similar chemicals eliciting avoidance in rodents may commonly occur in the faeces and urine of carnivores; this is supported by observations that rodents often avoid the odours of carnivores with which there has been no evolutionary contact (Stoddart 1982a, b).

Beavers showed the strongest response to the scent of river otters, a predator species not only sympatric but actually often living together with the beaver in the same habitats, indeed in the same lodges built by beavers apparently without preying upon them (Reid 1984, Brzezinski et al. 1996, V.E. Sidorovich, pers. comm.). From an ecological point of view, it is an extremely interesting and novel finding. It does not match the classical interpretation of 'repellents' as the river otter does not appear to be a predator on beaver. A plausible interpretation may be that river otter ex-

crement inhibited feeding by beavers more than the other predator excrements because it simply tastes or smells foul, and not because of predator recognition. Indeed, the beavers ate other sticks close by the otter sticks.

Seasonal variation

Beavers in our study ate more sticks with predator odours in autumn than in summer trials. We also found that beavers ate significantly more of all sticks during autumn than in summer. Engelhart & Müller-Schwarze (1995) also found a similar seasonal difference, and concluded that this was most likely correlated with reduced abundance of palatable food. In summer, beavers feed very little on tree bark and prefer to eat leaves and chew the bark of a twig only after all leaves are consumed. They also eat grass and herbs. This might explain less interest in the aspen sticks during summer. In late autumn, the growing season is over and beavers depend almost exclusively on bark for food. The beaver become less selective and the aspen samples might become more attractive (Engelhart & Müller-Schwarze 1995).

Predator odours may also become less effective after heavy rains (Sullivan, Nordstrom & Sullivan 1985, Engelhart & Müller-Schwarze 1995), and this might vary for each predator odour. In our study area there was only a small difference in monthly average of rainfall between the two field trials, so we assume rainfall did not affect our results. However, rain may have had an effect on the total result. The biggest problem in the work with odours as repellents has not been to find substances that animals will avoid, but to find substances that will have a long-term effect, i.e. substances that beavers do not habituate to, that does not wash away with rainfall or evaporate too fast (Sullivan et al. 1985, Engelhart & Müller-Schwarze 1995).

Management implications

Our results may have several practical implications. One implication is that predator odours could reduce damage by free-ranging Eurasian beaver to agricultural crops, fruit and forest trees, and be a humane, environmentally acceptable chemical that can be used to manage wild Eurasian beaver populations. In a few species predator odour has been a successful means to reduce feeding damage in field trials by 60-100% for time periods ranging from 1 to 5 months (Sullivan & Crump 1984, Sullivan 1986, Sullivan, Crump & Sullivan 1988a, Swihart 1991). Woodchucks *Marmota*

monax, for example, avoid marking fruit trees sprayed with bobcats' *L. rufus* urine for as long as 93 days, and continue to avoid bobcats' urine even when exposed to the odour in consecutive years (Swihart 1991). Trees are long-lived, and rodent damage is difficult to prevent completely. No repellent is likely to provide total protection. Nevertheless, predator odours, and in particular river otter odour, could reduce damage by beavers during periods when trees are most vulnerable, e.g. during early spring and late autumn. A component of the weasel's *M. nivalis* scent has successfully suppressed feeding damage to seedlings by hares *Lepus americanus* for a six-week period in a field bioassay trial (Sullivan & Crump 1984). This period covered the early spring when conifer seedlings were susceptible to hare damage, just after snow-melt and prior to hares switching to preferred summer herbaceous foods.

Because other rodents avoid burrows treated with predator odours (e.g. Sullivan, Crump & Sullivan 1988b), it is also conceivable that Eurasian beavers might avoid treated burrows, lodges and/or dens. Indeed, adult pocket gophers *Thomomys talpoides* were deterred from colonising an area treated with sulphur compounds from the anal glands of the stoat *M. erminea* (Sullivan & Crump 1986, Sullivan et al. 1990).

Attempts to prevent Eurasian beavers from repairing destroyed dams and plugging culverts may also be possible at least for a short period when it is most necessary. Guenther (1956) reported that the North American beavers temporarily avoided dams with bear *Ursus* sp. and cougar *F. concolor* scent, but resumed repair when the scent dissipated. However, Buech (1985) tried wolf faeces as deterrents to plugged culverts with no success.

Further clarification of the role and utility of predator odours as repellents of the Eurasian beaver is warranted. Also, testing of the organic components, especially of the river otter, and particularly the volatile components, may enable identification of the compounds responsible for the response we observed. This may explain several of the results reported here.

Acknowledgements - we thank Per Christian Hagen for statistical advice, Göran Hartman, Thrine M. Heggberget, Jan Heggnes, Arild Landa, Howard Parker, Bruce Schulte, Geir A. Sonerud and Lixing Sun, and two anonymous reviewers for their constructive comments on earlier drafts of this paper. We also thank the board of National Forests in Suwalki region and students of the Biology Department at Jagiellonian University for help in field works and Dr.

Andrzej Górecki, Director of Institute of Environmental Biology at Jagiellonian University for administrative support. This research was founded by grant no. 97/1/A/4 obtained from Castor Research Society, Poland.

References

- Andersone, Z. 1998: Beaver: a new wolf's prey in Latvia? Comparison of winter and summer diet of *Canis lupus* Linnaeus, 1758. - In: Reig, S. (Ed.); Abstracts. Euro-American Mammal Congress, Santiago de Compostela, Spain, July 19-24, 1998, pp. 166.
- Basey, J.M. & Jenkins, S.H. 1995: Influences of predation risk and energy maximization on food selection by beavers (*Castor canadensis*). - *Canadian Journal of Zoology* 73: 2197-2208.
- Berdoy, M. & Macdonald, D.W. 1991: Factors affecting feeding in wild rats. - *Acta Oecologica* 12: 261-279.
- Berenson, R. & Levine, M. 1989: Basic business statistics: Concepts and applications. - Prentice-Hall, Inc. A Division of Simon & Schuster Englewood Cliffs, New Jersey, 904 pp.
- Borowski, Z. 1998: Influence of predator odour on the feeding behaviour of the root vole (*Microtus oeconomus* Pallas, 1776). - *Canadian Journal of Zoology* 76: 1791-1794.
- Brzezinski, M., Romanowski, J., Cygan, J.P. & Pabin, B. 1996: Otter *Lutra lutra* distribution in Poland. - *Acta Theriologica* 41: 113-126.
- Brzezinski, M. & Zurowski, W. 1992: Spring diet of the American mink *Mustela vison* in the Mazurian and Brodnica Lakelands, northern Poland. - *Acta Theriologica* 37: 193-198.
- Buchalczyk, T. 1981: Red fox - *Vulpes vulpes* (Linnaeus, 1758). - In: Pucek, Z. (Ed.); Keys to vertebrates of Poland - Mammals, PWN, Warszawa, pp. 270-273.
- Buchalczyk, T. 1983: *Vulpes vulpes* (Linnaeus, 1758). - In: Pucek, Z. & Raczynski, J. (Eds.); Atlas of Polish mammals. PWN, Warszawa, pp. 141-145. (In Polish).
- Buech, R.R. 1985: Beaver in water impoundments: understanding a problem of water-level management. - In: Knighton, M.D. (Ed.); Water impoundments for wildlife: a habitat management workshop; 1982 August 31-September 2; Bemidji, MN. General Technical Report NC-100. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station, pp. 95-105.
- Dezhkin, V.V. & Safonov, G. 1966: Biology and economical usefulness of beaver. - Ekonomika Publication House, Moscow, U.S.S.R., 92 pp. (Translated from Russian).
- Dickman, C.R. 1992: Predation and habitat shift in the house mouse, *Mus domesticus*. - *Ecology* 73: 313-322.
- Dickman, C.R. & Doncaster, C.P. 1984: Responses of small mammals to red fox (*Vulpes vulpes*) odour. - *Journal of Zoology* 204: 521-531.

- Djoshkin, W.W. & Safonow, W.G. 1972: Die Biber der alten und neuen Welt. - A. Ziemsen Verlag, Wittenberg, 168 pp. (In German).
- Donkor, N.T. & Fryxell, J.M. 1999: Impact of beaver foraging on structure of lowland boreal forests of Algonquin Provincial Park, Ontario. - *Forest Ecology and Management* 118: 83-92.
- Dzieciolowski, R. 1996: Bobr. - Wydawnictwo SGGW, Warszawa, 124 pp. (In Polish).
- Dzieciolowski, R. & Gozdziwski, J. 1999: The reintroduction of European beaver, *Castor fiber*, in Poland. A success story. - In: Buser, P. & Dzieciolowski, R. (Eds.); *Beaver protection, management, and utilization in Europe and North America*. Kluwer Academic/Plenum Publishers, New York, pp. 31-35.
- Engelhart, A. & Müller-Schwarze, D. 1995: Responses of beaver (*Castor canadensis* Kuhl) to predator chemicals. - *Journal of Chemical Ecology* 21: 1349-1364.
- Gorman, M.L. 1984: The responses of prey to stoat (*Mustela erminea*) scent. - *Journal of Zoology* 202: 419-423.
- Guenther, S. 1956: A beaver baffler. - *Washington State Game Bulletin* 8: 3, 6.
- Heidecke, D. 1988: Methodik zur Auswahl von Wiederansiedlungsgebieten, Ansiedlung und Förderung des Elbesbibers (*Castor fiber albus*). - *Berichte Akademie der Landwirtschaftswissenschaften der DDR, Institut für Landschaftsforschung und Naturschutz, Halle*, 39 pp. (In German).
- Heidecke, D. & Klenner-Fringes, B. 1992: Studie über die Habitatnutzung des Bibers in der Kulturlandschaft. - In: Schröpfer, R., Stubbe, M. & Heidecke, D. (Eds.); *Materialien des 2. Internationalen Symposiums Semiaquatische Säugetiere*. Martin-Luther-Universität, Halle/Saale, pp. 215-265. (In German).
- Histøl, T. 1989: Sommerdiett hos bever (*Castor fiber*) i ett utvalg av skogsvann i Vennesla kommune, Vest-Agder. (In Norwegian with English summary: Summer diet of beavers *Castor fiber* L. in some small lakes in Vennesla, Vest-Agder). - *Fauna* 42: 96-103.
- Jakubiec, Z. & Buchalczyk, T. 1987: The brown bear in Poland: its history and present numbers. - *Acta Theriologica* 32: 289-306.
- Jedrzejewski, W., Rychlik, L. & Jedrzejewska, B. 1993: Responses of bank voles to odours of seven species of predators: experimental data and their relevance to natural predator-vole relationships. - *Oikos* 68: 251-257.
- Kamieniarz, R. & Panek, M. 1996: The occurrence and population trends of lynx in Poland, 1982-1994. - *Journal of Wildlife Research* 1: 171-173.
- Kile, N.B., Nakken, P.J., Rosell, F. & Espeland, S. 1996: Red fox, *Vulpes vulpes*, kills a European Beaver, *Castor fiber*, Kit. - *Canadian Field-Naturalist* 110: 338-339.
- Kudrjasov, V.S. 1973: Individernas sammensättning i bäverkolonier, samt förhållandet mellan arterna. - In: Den 5. allsovjjetiske konferansen om bever, pp. 82-85. (Translated to Swedish from Russian by fil. mag. Torsten Jansson).
- Mason, J.R., Epple, G. & Nolte, D.L. 1993: Semiochemicals and improvements in rodent control. - In: Galef, B.E., Valsecchi, P. & Mainardi, M. (Eds.); *Ontogeny and Social Transmission of Food Preferences in Mammals: Basic and Applied Research*. Horwood Academic Press, London, pp. 327-346.
- Merkens, M., Harestad, A.S. & Sullivan, T. 1991: Cover and efficacy of predator-based repellents for Townsend's vole, *Microtus townsendii*. - *Journal of Chemical Ecology* 17: 401-412.
- Müller-Schwarze, D. 1972: Responses of young black-tailed deer to predator odors. - *Journal of Mammalogy* 53: 393-394.
- Nolet, B.A. 1996: Management of the beaver (*Castor fiber*): towards restoration of its former distribution and ecological function in Europe? - *Report Council of Europe, Strasbourg*, 29 pp.
- Nolet, B.A. & Rosell, F. 1994: Territoriality and time budgets in beavers during sequential settlement. - *Canadian Journal of Zoology* 72: 1227-1237.
- Nolet, B.A. & Rosell, F. 1998: Comeback of the beaver *Castor fiber*: an overview of old and new conservation problems. - *Biological Conservation* 83: 165-173.
- Nolet, B.A., van der Veer, P.J., Evers, E.G.J. & Ottenheim, M.M. 1995: A linear programming model of diet choice of free-ranging beavers. - *Netherlands Journal of Zoology* 45: 317-335.
- Nolte, D.L., Farley, J.P., Campbell, D.L., Mason, J.R. & Epple, G. 1993: Potential repellents to prevent mountain beaver damage. - *Crop Protection* 12: 624-626.
- Nolte, D.L., Mason, J.R., Epple, G., Aronov, E. & Campbell, D.L. 1994: Why are predator urines aversive to prey? - *Journal of Chemical Ecology* 20: 1505-1516.
- Novak, M. 1987: Beaver. - In: Novak, M., Baker, J.A., Obbard, M.E. & Malloch, B. (Eds.); *Wild Furbearer Management and Conservation in North America*. Ontario, Ontario Ministry of Natural Resources, USA, pp. 283-312.
- Okarma, H. 1993: Status and management of the wolf in Poland. - *Biological Conservation* 66: 153-158.
- Pielowski, Z. 1982: Über die Bedeutung des Fuchses *Vulpes vulpes* (L. 1758) in der Jagdwirtschaft der Volksrepublik Polen. - *Beiträge zur Jagd- und Wildforschung* 12: 71-77. (In German).
- Reid, D.G. 1984: Ecological interactions of river otters and beavers in a boreal ecosystem. - M.Sc. thesis, University of Calgary, Alta, 199 pp.
- Richard, P.B. 1986: The Status of the Beaver in France. - *Zoologische Abhandlungen (Dresden)* 41: 121-130.
- Robinson, I. 1990: The effect of mink odor on rabbits and small mammals. - In: Macdonald, D.W., Müller-Schwarze, D. & Natynczuk S.E. (Eds.); *Chemical signals in vertebrates V*. Oxford University Press, Oxford, UK, pp. 567-572.
- Rocznik Statystyczny 1997: Główny Urząd Statystyczny. - Warszawa, 715 pp. (In Polish).
- Rocznik Statystyczny 1998: Główny Urząd Statystyczny. - Warszawa, 691 pp. (In Polish).

- Rosell, F. & Bergan, F. 1998: Free-ranging Eurasian beavers, Castor fiber, deposit anal gland secretion when scent marking. - *Canadian Field-Naturalist* 112: 532-535.
- Rosell, F., Bergan, F. & Parker, H. 1998: Scent-marking in the Eurasian beaver (*Castor fiber*) as a means of territory defense. - *Journal of Chemical Ecology* 24: 207-219.
- Rosell, F. & Hovde, B. 1998: Pine Marten, *Martes martes*, as a Eurasian Beaver, *Castor fiber*, lodge occupant and possible predator. - *Canadian Field-Naturalist* 112: 535-536.
- Rosell, F. & Nolet, B.A. 1997: Factors affecting scent-marking behavior in the Eurasian beaver (*Castor fiber*). - *Journal of Chemical Ecology* 23: 673-689.
- Rosell, F. & Parker, H. 1995: Forvaltning av bever: dagens tilstand og fremtidig behov. (In Norwegian with English summary: Beaver management: present practice and Norway's future needs.) - Report Telemark College, Bø in Telemark, Norway, 137 pp.
- Rosell, F. & Parker, H. 1996: Beverens innvirkning på økosystemet - en nøkkelart vender tilbake. (In Norwegian with English summary: The beaver's (*Castor spp.*) role in forest ecology: a key species returns.) - *Fauna* 49: 192-211.
- Rosell, F., Parker, H. & Kile, N.B. 1996: Dødsårsaker hos bever. (In Norwegian with English summary: Causes of mortality in beaver (*Castor fiber* & *canadensis*)). - *Fauna* 49: 34-46.
- Rosell, F. & Pedersen, K.V. 1999: Bever. - *Landbruksforlaget*, 272 pp. (In Norwegian).
- Shelton, P.C. & Peterson, R.O. 1983: Beaver, wolf and moose interactions in Isle Royale National Park, USA. - *Acta Zoologica Fennica* 174: 265-266.
- Simonsen, T.A. 1973: Beverens næringsøkologi i Vest-Agder. - *Meddelelser fra Statens viltundersøkelser* 2: 1-66. (In Norwegian).
- Smith, D.W., Trauba, D.R. & Anderson, R.K. 1994: Black bear predation on beavers on an island in Lake Superior. - *American Midland Naturalist* 132: 248-255.
- Sokal, R.R. & Rohlf, F.J. 1995: *Biometry: The principles and practice of statistics in biological research.* - W.H. Freeman and company, 3rd edition, New York, USA, 887 pp.
- Stoddart, D.M. 1980: Some responses of a free living community of rodents to the odors of predators. - In: Müller-Schwarze, D. & Silverstein, R.M. (Eds.); *Chemical signals: vertebrates and aquatic invertebrates.* Plenum Press, New York, USA, pp. 1-10.
- Stoddart, D.M. 1982a: Demonstrations of prey olfactory discrimination by the short-tailed vole, *Microtus agrestis*. - *Animal Behaviour* 20: 293-294.
- Stoddart, D.M. 1982b: Does trap odour influence estimation of population size of the short-tailed vole, *Microtus agrestis*? - *Journal of Animal Ecology* 30: 375-386.
- Sullivan, T.P. 1986: Influence of wolverine (*Gulo gulo*) odor on feeding behavior of snowshoe hares (*Lepus americanus*). - *Journal of Mammalogy* 67: 385-388.
- Sullivan, T.P. & Crump, D.R. 1984: Influence of Mustelid scent-gland compounds on suppression of feeding by snowshoe hares (*Lepus americanus*). - *Journal of Chemical Ecology* 10: 1809-1821.
- Sullivan, T.P. & Crump, D.R. 1986: Avoidance response of pocket gophers (*Thomomys talpoides*) to mustelid anal gland compounds. - In: Duvall, D., Müller-Schwarze, D. & Silverstein, R.M. (Eds.); *Chemical signals in vertebrates IV, ecology, evolution and comparative biology.* Plenum, New York, New York, USA, pp. 519-531.
- Sullivan, T.P., Crump, D.R. & Sullivan, D.S. 1988a: Use of predator odors as repellents to reduce feeding damage by herbivores. III. Montane and meadow voles (*Microtus montanus* and *Microtus pennsylvanicus*). - *Journal of Chemical Ecology* 14: 363-377.
- Sullivan, T.P., Crump, D.R. & Sullivan, D.S. 1988b: Use of predator odors as repellents to reduce feeding damage by herbivores. IV. Northern pocket gophers (*Thomomys talpoides*). - *Journal of Chemical Ecology* 14: 379-389.
- Sullivan, T.P., Crump, D.R., Wieser, H. & Dixon, A. 1990: Response of pocket gophers (*Thomomys talpoides*) to an operational application of synthetic semiochemicals of stoat (*Mustela erminea*). - *Journal of Chemical Ecology* 16: 941-949.
- Sullivan, T.P., Nordstrom, L.O. & Sullivan, D.S. 1985: Use of predator odors as repellents to reduce feeding damage by herbivores. II. Black-tailed deer (*Odocoileus hemionus columbianus*). - *Journal of Chemical Ecology* 11: 921-935.
- Swihart, R.K. 1991: Modifying scent marking behavior to reduce woodchuck damage to fruit trees. - *Ecological Applications* 1: 98-103.
- Tumilson, R. & Karnes, M. 1987: Seasonal changes in food habits of river otters in southwestern Arkansas beaver swamps. - *Mammalia* 51: 225-231.
- Tyurnin, B.N. 1984: Factors determining numbers of the river beavers (*Castor fiber*) in the European North. - *Soviet Journal of Ecology* 14: 337-344.
- Vernet-Maury, E., Polak, E.H. & Demael, A. 1984: Structure-activity relationship of stress inducing odorants in the rat. - *Journal of Chemical Ecology* 10: 1007-1018.
- Ward, J.F., Macdonald, D.W. & Doncaster, C.P. 1997: Responses of foraging hedgehogs to badger odour. - *Animal Behaviour* 53: 709-720.
- Weldon, P.J., Graham, D.P. & Mears, L.P. 1993: Carnivore faecal chemicals suppress feeding by alpine goats (*Capra hircus*). - *Journal of Chemical Ecology* 19: 2947-2952.
- Wilsson, L. 1971: Observations and experiments on the ethology of the European Beaver (*Castor fiber* L.). - *Viltrevy* 8: 115-306.
- Zurowski, W. 1989: Bobr europejski. - In: *Lowiectwo, PWRiL, Warszawa*, pp. 321-323. (In Polish).
- Zurowski, W. 1992: Building activity of beavers. - *Acta Theriologica* 37: 403-411.
- Zurowski, W. & Kasperczyk, B. 1986: Characteristics of a European beaver population in the Suwalki Lakeland. - *Acta Theriologica* 31: 311-325.