

Breeding systems and cytology in Cyprian populations of six Limonium Species (Plumbaginaceae)

Authors: Iro Kouzali, Rea Artelari, and Ourania Georgiou

Source: Willdenowia, 42(2) : 307-313

Published By: Botanic Garden and Botanical Museum Berlin (BGBM)

URL: <https://doi.org/10.3372/wi.42.42217>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

IRO KOUZALI¹, REA ARTELARI^{1*} & OURANIA GEORGIU¹

Breeding systems and cytology in Cyprian populations of six *Limonium* species (*Plumbaginaceae*)

Abstract

Kouzali I., Artelari R. & Georgiou O.: Breeding systems and cytology in Cyprian populations of six *Limonium* species (*Plumbaginaceae*). – Willdenowia 42: 307–313. December 2012. – Online ISSN 1868-6397; © 2012 BGBM Berlin-Dahlem.

Stable URL: <http://dx.doi.org/10.3372/wi.42.42217>

The reproductive mechanisms and chromosome numbers have been investigated of the Cyprian endemics *Limonium cypricum* and *L. mucronulatum*, as well as of *L. aucheri*, *L. avei*, *L. meyeri* and *L. virgatum*, which also occur on the island of Cyprus. Their taxonomy is considered and the correlation between their breeding systems and cytology is discussed. *L. aucheri* and *L. cypricum* are apomictic and their chromosome numbers, $2n=5x=43$ and $2n=3x=27$ respectively, are reported here for the first time. For *L. mucronulatum* no chromosome number could be established, but pollen and stigma features indicate that it is apomictic too. *L. meyeri* is sexual with $2n=2x=18$ (first report), while *L. avei* and *L. virgatum* are apomictic with $2n=3x=27$.

Additional key words: Cyprus, chromosome numbers, sexual reproduction, apomictic reproduction

Introduction

The genus *Limonium* Mill., with about 400 species (Palacios & Gonzales-Candelas 1997) the largest genus of *Plumbaginaceae*, is one of the taxonomically most difficult plant taxa. This difficulty is due to its reproductive behaviour, i.e. the presence of both sexual and apomictic reproduction, as well as the frequent occurrence of hybridisation and polyploidy. Most of the *Limonium* taxa are concentrated in the Mediterranean region, which constitutes the centre of diversity of this genus (Cowan & al. 1998; Lledó & al. 2003).

In the island of Cyprus, *Limonium* is represented by eight species, a rather small number compared to the species numbers on other islands of the E Mediterranean.

According to Buttler & Hand (in Hand 2003), the Cyprus flora is relatively poor in *Limonium* species, e.g. in comparison to Crete, the genus being absent from many parts of the coast, especially from areas dominated by igneous rocks. Indeed, in Crete 18 *Limonium* species occur (Mayer 1995; Brullo & Guarino 2000), while from the much smaller island of Kithira still nine species are reported (Artelari & Georgiou 2002).

According to Bokhari & Edmondson (1982), Meikle (1985) and Greuter & al. (1989), the species known from Cyprus so far include the Cyprian endemics *Limonium cypricum* (Meikle) Hand & Buttler and *L. mucronulatum* (H. Lindb.) Greuter & Burdet, as well as *L. aucheri* (Girard) Greuter & Burdet, *L. avei* (De Not.) Brullo & Erben, *L. echioides* (L.) Mill., *L. meyeri* (Boiss.) Kuntze, *L. sin-*

1 Division of Plant Biology, Department of Biology, University of Patras, Rion 26500, Patras, Greece; *e-mail: artelari@upatras.gr (author for correspondence).

uatum (L.) Mill. and *L. virgatum* (Willd.) Fourr.

Very few cytological information exists for *Limonium* on Cyprus and there is a complete lack of data concerning their breeding systems. In the present work material from several localities of the island belonging to six of the eight above-mentioned species has been studied, focusing mainly on their cytology and reproductive mechanism.

Material and methods

This study is based on plants collected mainly by the first author from several localities of Cyprus in the framework of her undergraduate diploma thesis. This material includes thirty-four populations from six of the eight species present on the island. No material of *Limonium echioides* and *L. sinuatum* has been found. Voucher specimens are kept in the Herbarium of the University of Patras (UPA).

All the collected populations were studied to assess their reproductive mechanism. Pollen and stigma type combinations of the flowers were determined according to Erben (1978, 1979). The pollen stainability was estimated by using cotton blue as described in Artelari & Kamari (1986). Fifteen populations were used for the cytological studies. For this purpose, seeds were germinated in Petri dishes on moistened filter paper and the root tips were pretreated according to the method described in previous papers (Artelari 1984; Artelari & Kamari 1986).

Results and Discussion

1. *Limonium cypricum* (Meikle) Hand & Buttler in Willdenowia 33: 315. 2003 \equiv *Limonium albidum* subsp. *cypricum* Meikle in Ann. Mus. Goulandris 6: 88. 1983.

Distribution — Endemic to Cyprus.

Material examined — Kato Pafos, “Tafi ton Vasileon”, on calcareous maritime rocks, *Christodoulou* 27 (UPA).

Taxonomic notes — *Limonium cypricum* was described by Meikle (1983) as a subspecies of *L. albidum* (Guss.) Pignatti, which is endemic to the island of Lampion, S Italy. According to Meikle (1985), it differs from the typical *L. albidum* in the smaller flowers with straighter calyx tubes and bracts. Greuter & al. (1989) placed this taxon as a subspecies into the *L. intermedium* group, which else comprises the endemics of S Italy (around Sicily) *L. albidum*, *L. hyblaicum* Brullo, *L. intermedium* (Guss.) Brullo, *L. lopadusanum* Brullo, *L. mazararum* Pignatti and *L. panormitanum* (Tod.) Pignatti, as well as *L. zembrae* Pignatti, endemic to Tunisia. Buttler & Hand (in Hand 2003), comparing the plants of Cyprus to the typical *L. albidum*, reported that they are very similar but all taxo-

nomically important parts of the spikelets and flowers of the Cyprian plants are smaller than those of typical *L. albidum*. They therefore came to the conclusion that the Cyprian taxon should be considered as a separate species, which probably constitutes a relict of the *L. intermedium* group.

Cytology and breeding system — Our study revealed the triploid chromosome number $2n=3x=27$. The karyotype (Fig. 1A) lacks the long metacentric “marker” chromosomes characteristic for karyotypes with the basic number $x=8$. The only previous report, by Yıldız & Gücel (2006), gives the diploid chromosome number $2n=18$.

Concerning the pollen and stigma combination, the population of this taxon is monomorphic with the self-incompatible combination B, indicating self-sterility. Pollen stainability is 0 % and the pollen grains are misshaped, irregular in size (conspicuously large or conspicuously small), with variable morphology (tri- and tetrapolypate). Such pollen and stigma features characterise apomictic species (Baker 1953c). Our data combined with the good seed production of the population and the triploid chromosome number indicate that *Limonium cypricum* reproduces apomictically.

2. *Limonium mucronulatum* (H. Lindb.) Greuter & Burdet in Willdenowia 19: 40. 1989 \equiv *Statice mucronulata* H. Lindb. in Acta Soc. Sci. Fenn., Ser. B, Opera Biol. 2(7): 26. 1946.

— *Limonium narbonense* sensu Meikle (1985).

Distribution — Endemic to Cyprus.

Material examined — Larnaka, Alikes area, on sand, *Kouzali* 31 (UPA), *Christodoulou* 26 (UPA).

Taxonomic notes — *Limonium mucronulatum* is only known from a single locality (area of Alike), where it was rediscovered by Christodoulou & Tsintides (Tsintides & al. 2007) about 50 years after its first discovery by Lindberg in 1939 (Lindberg 1946).

Meikle (1985) erroneously used the name *Limonium narbonense* Mill. Greuter & Raus (1989), referring to it as *L. mucronulatum*, place it into the *L. sibthorpium* group, which comprises also *L. raddianum* (Boiss.) Pignatti from Egypt, *L. sibthorpium* (Guss.) Kuntze from Sicily as well as *L. teuchirae* Brullo and *L. vaccarii* Brullo from Libya.

Limonium mucronulatum is included in the “Red Book of the flora of Cyprus” (Tsintides & al. 2007) as “Critically Endangered” (CR), comprising only 600–700 plants distributed in an area of 3000 m² and its habitat being seriously threatened.

Cytology and breeding system — No cytological data are known so far for *Limonium mucronulatum*. In spite of



Fig. 1. Mitotic metaphase plates of *Limonium* from Cyprus – A: *L. cypricum*, $2n=3x=27$, material from Kato Pafos (*Christodoulou 27*); B: *L. aucheri*, $2n=5x=43$, material from Cavo Greco (*Kouzali 16*); C: *L. meyeri*, $2n=2x=18$, material from Agia Napa (*Kouzali 10*); D: *L. avei*, $2n=3x=27$, material from Cavo Greco (*Kouzali 36*). – Scale bar = 10 μm .

our strenuous efforts, we did not succeed to obtain an accurate chromosome count.

Concerning pollen and stigma combination, the material collected belongs to two populations which were found to be monomorphic with the self-incompatible combination B, low pollen stainability (3–13 %) and good seed production. Pollen grains are tri- to tetra-lobate, irregular in size, the stainable ones being well-shaped, the unstainable ones being much smaller and misshaped. As this situation is characteristic for apomictic taxa, the pollen and stigma features allow us to conclude that *Limonium mucronulatum* reproduces apomictically, although no data for its chromosome number are available so far.

3. *Limonium aucheri* (Girard) Greuter & Burdet in Willdenowia 19: 39. 1989 \equiv *Statice aucheri* Girard in Ann. Sci. Nat. Bot., ser. 3, 2: 328. 1844. = *Limonium ocymifolium* subsp. *bellidifolium* (Sm.) Meikle, Fl. Cyprus: 1070. 1985 \equiv *Statice bellidifolia* Sm., Fl. Graec. Prodr. 1: 211. 1806 [non (Gouan) DC. in Lamarck & Candolle, Fl. Franc. ed. 3, 3: 421. 1805].

Distribution — Reported so far from the Greek islands of Kithnos, Rodos and Kriti as well as from Cyprus.

Material examined — Protaras area, “Fig Tree” bay, near the hotel, on calcareous maritime rocks, *Kouzali 17* (UPA); Protaras area, 300 m east of the “Fig Tree” bay, on calcareous maritime rocks, *Kouzali 18*, *Christodou-*

lou 25 (UPA); Cape Cavo Greco, on calcareous maritime rocks, *Kouzali 16* (UPA).

Taxonomic notes — Greuter & al. (1989) placed this taxon into the *Limonium ocymifolium* group, which comprises allied taxa mainly distributed in the C and S Aegean and the S Peloponnisos. According to Greuter & al. (1989) and Artelari & Georgiou (1999), six species are included: *L. ocymifolium* (Poir.) Kunze, *L. hierapetrae* Rech. f., *L. creticum* Artelari, *L. cythereum* Artelari & Georgiou, *L. corinthiacum* (Boiss. & Heldr.) Kunze and *L. aucheri*. In addition, on the basis of morphological, cytological and breeding system features, we suggest that *L. runemarkii* Rech. f. and *L. doerfleri* (Halácsy) Rech. f. also belong to that group. The placement of *L. runemarkii* in the *L. palmare* group instead (Greuter & al. 1989) is not justified because of the morphological differences.

According to Greuter & Raus (1989), *Limonium aucheri* is a doubtful species corresponding to *Statice bellidifolia* Sm. This last taxon has been treated by Boissier (1879), Halácsy (1904) and Hayek (1928) as a variety of *S. ocymifolia* Poir., by Rechinger (1943a, b) as a variety of *L. ocymifolium* (Poir.) Kuntze and by Meikle (1985) as a subspecies of *L. ocymifolium*.

Mayer (1995) comments that the taxonomic status of *Limonium aucheri* concerns a case of great confusion, as the names *Statice aucheri* and *S. bellidifolia* are based on different types, the first one on a specimen collected by Aucher-Eloy from Kithnos, and the second on a specimen of Smith from Rodos, but that *L. aucheri* is possibly conspecific with *L. ocymifolium*. We also consider

L. aucheri and *L. ocymifolium* as very closely related. Smith's *Statice bellidifolia*, figured in Sibthorp & Smith (1821: t. 295), is very similar to *L. ocymifolium* and a detailed study of additional material of *L. aucheri* will probably lead to the fusion of these two species. This is also supported by the fact that they have the same chromosome number, $2n=43$ (see below).

Limonium aucheri is classified by Tsintides & al. (2007) as "Endangered" (EN), being known from two localities only (Protaras area and Cape Cavo Greco).

Cytology and breeding system — Our cytological study of *Limonium aucheri* revealed the pentaploid chromosome number $2n=5x=43$, derived from the combination of $2 \times 8 + 3 \times 9$ genomes. This is indicated by the presence of two long metacentric "marker" chromosomes (Fig. 1B), characteristic for karyotypes with the basic number $x=8$ (Erben 1978, 1979) and shows that the taxon is of hybrid origin. The chromosome number of *L. aucheri* is given here for the first time.

The study of the pollen and stigma combination showed that all three populations are monomorphic, two with the self-incompatible combination B and pollen stainability 0–3 %, and one with the self-incompatible combination A and pollen stainability 0–24 %. Pollen grains of all populations are irregular in size. Stainable pollen grains are well-shaped and much larger than the unstainable ones which are small and misshaped. The above data combined with the good seed production of the plants and the pentaploid chromosome number indicate that *Limonium aucheri* is an apomictic species. The closely related *L. ocymifolium* has the same pentaploid chromosome number $2n=43$ and is also apomictic (Artelari 1989; Artelari & Georgiou 2002).

4. *Limonium meyeri* (Boiss.) Kuntze, Revis. Gen. Pl. 1: 395. 1981 \equiv *Statice meyeri* Boiss. in Candolle, Prodr. 12: 645. 1848.

– *Limonium gmelinii* sensu Osorio-Tafall & Seraphim (1973)

– *Limonium vulgare* sensu Osorio-Tafall & Seraphim (1973)

Distribution — Bulgaria, Crimea, Anatolia and Cyprus.

Material examined — Agia Napa, islet of the bay "Nissi", on calcareous maritime rocks, *Kouzali 10* (UPA); Cape Cavo Greco, Agii Anargiri, on calcareous maritime rocks, *Kouzali 13* (UPA); Larnaka, Alikes area, Aliki Spiros, on sand, *Kouzali 33* (UPA).

Taxonomic notes — *Limonium meyeri* belongs to the Mediterranean *L. vulgare* group, which also comprises *L. brevipetiolatum* Artelari & Erben, *L. effusum* (Boiss.) Kuntze, *L. gmelinii* (Willd.) Kuntze, *L. humile* Mill., *L. narbonense* Mill., *L. vanense* Kit Tan & Sorger and *L.*

vulgare Mill. (Greuter & al. 1989). Its closest relative is *L. gmelinii* from which it differs in the taller growth, the more open, loosely paniculate inflorescence and the remotely spaced spikelets (Bokhari & Edmondson 1982).

Cytology and breeding system — All three populations studied were found to be diploid with $2n=18$ (Fig. 1C). The chromosome number of *Limonium meyeri* is reported here for the first time.

Concerning pollen and stigma combination, the populations are dimorphic and have the self-incompatible combinations A and B. This is in accordance with Baker's report (1953a) that *Limonium meyeri* is dimorphic. Pollen and stigma features are typical for sexual taxa, i.e. all pollen grains are regular in size, well-shaped and their stainability varies from 17 % to 100 %. These data, combined with the diploid chromosome number, indicate that *L. meyeri* is a sexually reproducing species.

According to available information, most species of the *Limonium vulgare* group are dimorphic and sexual with chromosome numbers derived from the basic number $x=9$ and karyotypes not possessing "marker" chromosomes. More specifically, *L. brevipetiolatum* is dimorphic sexual and hexaploid with $2n=54$ (Artelari & Erben 1986), *L. gmelinii* is dimorphic (Baker 1953a), *L. narbonense* is dimorphic sexual, tetraploid, hexaploid and octaploid with $2n=36, 54, 72$, respectively (Erben 1978, 1993; Brullo & Pavone 1981; Artelari 1992; Palacios & al. 2000; Georgakopoulou & al. 2006) and *L. vulgare* is dimorphic sexual and tetraploid with $2n=36$ (Baker 1953a; Erben 1979, 1993; Palacios & al. 2000). The only exception so far is *L. humile*, which according to Baker (1953a, b) is secondarily monomorphic with the self-compatible type D, and according to Erben (1979) has the pentaploid chromosome number $2n=54$.

5. *Limonium virgatum* (Willd.) Fourr. in Ann. Soc. Linn. Lyon, ser. 2, 17: 141. 1869 \equiv *Statice virgata* Willd., Enum. Pl. Hort. Berol.: 336. 1809.

= *Statice smithii* Ten., Fl. Napol. 3: 350. 1829.

– *Limonium oleifolium* Mill., Gard. Dict. Ed. 8, no 3. 1768, nom. amb.

– *Statice oleifolia* auct., non Scop., Del. Fl. Faun. Insubr. 1: 24, t. 10. 1786.

Distribution — Widely distributed in the Mediterranean region as well as in C and S Portugal.

Material examined — Agia Thekla area, on calcareous maritime rocks, *Kouzali 2* (UPA); in the small bay near the church of Agia Thekla, on calcareous maritime rocks, *Kouzali 3* (UPA); Agia Thekla area, on sand, *Kouzali 1* (UPA); Agia Napa, bay "Dome Hotel", on calcareous maritime rocks, *Kouzali 5* (UPA); Agia Napa, Makronisos, bay "Adamos", on calcareous maritime rocks, *Kouzali 6* (UPA); Agia Napa, bay "Nissi", on calcareous

maritime rocks, *Kouzali 8* (UPA); Agia Napa, bay “Yi-annoula Hotel”, on calcareous maritime rocks, *Kouzali 9* (UPA); Cape Cavo Greco, bay “Kamara tou Koraka”, on calcareous maritime rocks, *Kouzali 12* (UPA); Cape Cavo Greco, near the church of Agii Anargiri, on calcareous maritime rocks, *Kouzali 14* (UPA); Agios Thirsos, in the area of the church, on calcareous maritime rocks, *Kouzali 19* (UPA); in the area of the monastery Apostolos Andreas, on calcareous maritime rocks, *Kouzali 21* (UPA); Karpasia area, “Acheon Akti”, on sand dunes, *Kouzali 22* (UPA); Kato Pafos, “Tafi ton Vasileon”, on calcareous maritime rocks, *Christodoulou 28* (UPA); Protaras area, “Fig Tree” bay, on calcareous maritime rocks, *Kouzali 17* (UPA).

Taxonomic notes — *Limonium virgatum* is characterised by strongly dichotomously branched stems with numerous short sterile branches, compact, arcuate spikes and strongly banana-like curved spikelets. It often grows together with other *Limonium* species, forming intermediates (Pignatti 1972; Artelari 1984; Erben 1993). In Cyprus, in the area of Cavo Greco where *L. virgatum* coexists with *L. aucheri*, some plants present intermediate features such as sterile branches, somewhat curved spikelets (features of *L. virgatum*) and spatulate-obovate, ± emarginate leaves (features of *L. aucheri*), being the first record of intermediate forms of *Limonium* from Cyprus.

Cytology and breeding system — The triploid chromosome number $2n=27$ was counted in two populations and confirms previous counts from Greek material (Artelari 1984, 1989, 1992; Artelari & Georgiou 2002; Georgakopoulou & al. 2006) as well as from elsewhere (D’Amato 1949; Baker 1952, 1953b, c; Erben 1978, 1979; Chichiricco & Tammara 1980; Brullo & Pavone 1891; Ingrouille 1984; Arrigoni & Diana 1993; Palacios & al. 2000). Dolcher & Pignatti (1971) reported $2n=24-27$ and the tetraploid number $2n=4x=32$.

Concerning pollen and stigma combination, all studied populations are monomorphic with combination B. This confirms previous findings (Dulberger 1975; Ingrouille 1984; Artelari 1989a; Artelari & Georgiou 2002) with the exception of Baker (1953a), who states that *Limonium virgatum* is dimorphic and sexual but “some varieties of it may not be dimorphic”. Pollen stainability is low (0–8 %) and seed production is good. Pollen grains are irregular in size, the unstainable ones being misshaped and much smaller than the well-shaped stainable ones. These data and the triploid chromosome number confirm that *L. virgatum* reproduces apomictically, which was reported also by Erben (1979), Artelari (1984), Artelari & Georgiou (2002) and Georgakopoulou & al. (2006).

6. *Limonium avei* (De Not.) Brullo & Erben in Willdenowia 17: 17. 1988 ≡ *Statice avei* De Not., Prosp. Fl. Ligust.: 54. 1846.

= *Statice echioides* subsp. *exaristata* Murb. in Acta Univ. Lund. 35(3): 1. 1899 ≡ *Limonium echioides* subsp. *exaristatum* (Murb.) Maire in Jalandiez & Maire, Cat. Pl. Maroc: 571. 1934 ≡ *Limonium exaristatum* (Murb.) P. Fourn., Quatre Fl. France: 720. 1937.
= *Limonium longispicatum* Erben in Mitt. Bot. Staats-samml. München 14: 555. 1978.

Distribution — France, S Sardinia, Italy, Sicily, Greece, Crete, Cyprus, Egypt, Tunisia and Libya.

Material examined — Cape Cavo Greco, near the church of Agii Anargiri, on calcareous maritime rocks, *Kouzali 36* (UPA); Protaras area, under “Cavo Maris” hotel, on calcareous maritime rocks, *Christodoulou 33* (UPA); Larnaka, Alikes area, Aliki Spiros, on sand, *Kouzali 40* (UPA).

Taxonomic notes — *Limonium avei* is characterised by its annual to biennial life form. Meikle (1985) refers to this taxon as *L. echioides* subsp. *exaristatum* (Murb.) Maire, indicating its close relationship to the annual Mediterranean *L. echioides* (L.) Mill., also occurring in Cyprus (Meikle 1985; Brullo 1988; Hand 2009). *L. avei* mainly differs from *L. echioides* in having mucronate to acute leaves, calyx ribs not or just shortly excurrent into an awn and a generally more robust habit. Moreover, *L. avei* is triploid with $2n=27$ (see below), while *L. echioides* is diploid with $2n=18$ (Dolcher & Pignatti 1971; Erben 1979; Artelari & Georgiou 2002).

Cytology and breeding system — The triploid chromosome number $2n=27$ was found in the three populations of *Limonium avei* studied. This number has been already reported by Erben (1978, under *L. longispicatum*), Brullo (1988) and Brullo & Erben (1989). Its karyotype lacks the long metacentric “marker” chromosomes characteristic for karyotypes with the basic number $x=8$ (Fig. 1D).

The examination of pollen and stigma showed that all individuals of the three populations of *Limonium avei* are male sterile and have a cob-like stigma. Nevertheless, they have good seed production and this, combined with the triploid chromosome number, indicates that *L. avei* reproduces apomictically. Existence of male sterile plants in *L. avei* has not been reported so far, but Baker (1953a) and Ingrouille (1984) stated the occurrence of male sterile colonies with cob-like stigmata in the closely related diploid *L. echioides*, which is a secondarily monomorphic species with the self-compatible combination C.

Acknowledgements

We would like to thank Mr C. Christodoulou, PhD Student of the Centre for Agri-Environmental Research (CAER), Department of Agriculture, of the University of Reading, for the specimens of *Limonium* kindly of-

ferred to us, as well as for the valuable help and support to the first author during her field work. We also thank Dr S. Spanou for her linguistic comments to this manuscript.

References

- Arrigoni P. V. & Diana S. 1993: Contribution à la connaissance du genre *Limonium* en Corse. – *Candollea* **48**: 631–677.
- Artelari R. 1984: Biosistimatiki meléti tou génous *Limonium* (*Plumbaginaceae*) stin periochi tou Ioniou pelágous. – Ph. D. Thesis Patras University.
- Artelari R. 1989: Biosystematic study of the genus *Limonium* (*Plumbaginaceae*) in the Aegean area (Greece). I. Some *Limonium* species from the Kiklades islands. – *Willdenowia* **18**: 399–408.
- Artelari R. 1992: Reports 51–55. [In Kamari G., Felber F. & Garbari F. (ed.), *Mediterranean chromosome number reports* 2]. – *Fl. Medit.* **2**: 229–232.
- Artelari R. & Erben M. 1986: *Limonium brevipetiolatum* – eine neue hexaploide Sippe aus Süd-Griechenland. – *Mitt. Bot. Staatssamml. München* **22**: 507–511.
- Artelari R. & Georgiou O. 1999: Two new species of *Limonium* (*Plumbaginaceae*) from the island of Kithira (Greece). – *Bot. J. Linn. Soc.* **131**: 399–415.
- Artelari R. & Georgiou O. 2002: Biosystematic study of the genus *Limonium* (*Plumbaginaceae*) in the Aegean area, Greece. III. *Limonium* on the islands Kithira and Antikithira and the surrounding islets. – *Nordic J. Bot.* **22**: 483–501.
- Artelari R. & Kamari G. 1986: A karyological study of ten *Limonium* species (*Plumbaginaceae*) endemic in the Ionian area. – *Willdenowia* **15**: 497–513.
- Baker H. G. 1952: Cytotaxonomic studies in the *Plumbaginaceae*. – *Heredity* **6**: 279–280.
- Baker H. G. 1953a: Dimorphism and monomorphism in the *Plumbaginaceae*. II. Pollen and stigmata in the genus *Limonium*. – *Ann. Bot.* **17**: 433–445.
- Baker H. G. 1953b: Dimorphism and monomorphism in the *Plumbaginaceae*. III. Correlation of geographical distribution patterns with dimorphism and monomorphism in *Limonium*. – *Ann. Bot.* **17**: 615–627.
- Baker H. G. 1953c: The agamic complex in *Limonium*, subsections *Densiflorae* and *Dissitiflorae*. – Pp. 329–330 in: [Anon. ed.], *Proceedings of the 7th International Botanical Congress, Stockholm*. – Stockholm, Almquist & Wiksell.
- Boissier E. 1879: *Flora orientalis* **4**. – Basel: Georg.
- Bokhari M. H. & Edmondson J. R. 1982: *Limonium* Miller. – Pp. 465–476 in: Davis P. H. (ed.), *Flora of Turkey and the East Aegean Islands* **7**. – Edinburgh: University Press.
- Brullo S. 1988: Miscellaneous notes on the genus *Limonium* (*Plumbaginaceae*). – *Willdenowia* **17**: 11–18.
- Brullo S. & Erben M. 1989: The genus *Limonium* (*Plumbaginaceae*) in Tunisia. – *Mitt. Bot. Staatssamml. München* **28**: 419–500.
- Brullo S. & Guarino R. 2000: Contribution to the knowledge of flora and vegetation of Khrisi islet (Crete, SE Mediterranean sea). – *Fl. Medit.* **10**: 265–282.
- Brullo S. & Pavone P. 1981: Chromosome numbers in the Sicilian species of *Limonium* Miller (*Plumbaginaceae*). – *Anales Jard. Bot. Madrid* **37**: 535–555.
- Chichiricco G. & Tammario F. 1980: Numeri cromosomici per la flora italiana. – *Inform. Bot. Ital.* **12**: 324–325.
- Cowan R., Ingrouille M. J. & Lledó M. D. 1998: The taxonomic treatment of agamosperms in the genus *Limonium* (*Plumbaginaceae*). – *Folia Geobot.* **33**: 353–366.
- D'Amato F. 1949: Triploidia e apomissia in *Statice oleifolia* Scop. var. *confusa* Godr. – *Caryologia* **2**: 71–84.
- Dolcher T. & Pignatti S. 1971: Un'ipotesi sull'evoluzione dei *Limonium* del bacino del Mediterraneo. – *Giorn. Bot. Ital.* **105**: 95–107.
- Dulberger R. 1975: Intermorph structural differences between stigmatic papillae and pollen grains in relation to incompatibility in *Plumbaginaceae*. – *Proc. Roy. Soc. London, B*, **188**: 257–274.
- Erben M. 1978: Die Gattung *Limonium* im südwest-mediterranen Raum. – *Mitt. Bot. Staatssamml. München* **14**: 361–631.
- Erben M. 1979: Karyotype differentiation and its consequences in Mediterranean *Limonium*. – *Webbia* **34**: 409–417.
- Erben M. 1993: *Limonium* Mill. – Pp. 2–143 in: Castroviejo S., Aedo C., Cirujano S., Lainz M., Montserrat P., Morales R., Muñoz Garmendia F., Navarro C., Paiva J. & Soriano C. (ed.), *Flora iberica* **3**. – Madrid: Real Jardín Botánico.
- Georgakopoulou A., Manousou S., Artelari R. & Georgiou O. 2006: Breeding systems and cytology in Greek populations of five *Limonium* species (*Plumbaginaceae*). – *Willdenowia* **36**: 741–750.
- Greuter W. & Raus T. (ed.) 1989: *Med-Checklist Notulae*, 15. – *Willdenowia* **19**: 27–48.
- Greuter W., Burdet H. M. & Long G. (ed.) 1989: *Med-Checklist* **4**. – Genève: Conservatoire et Jardin botaniques; Berlin: Botanischer Garten and Botanisches Museum Berlin-Dahlem.
- Halácsy E. von 1904: *Conspectus florae Graecae* **3**. – Leipzig: Engelmann.
- Hand R. 2003: Supplementary notes to the Flora of Cyprus III. – *Willdenowia* **33**: 305–325.
- Hand R. 2009: Supplementary notes to the Flora of Cyprus VI. – *Willdenowia* **39**: 301–325.
- Hayek A. de 1928: *Prodromus florae peninsulae balcanicae* 2. – *Repert. Spec. Nov. Regni Veg. Beih.* **30(2)**: 1–96.
- Ingrouille M. J. 1984: A taxonomic analysis of *Limonium* (*Plumbaginaceae*) in western Europe. – *Pl. Syst. Evol.* **147**: 103–118.

- Lledó M. D., Erben M. & Crespo M. B. 2003: *Myriolepis*, a new genus segregated from *Limonium* (*Plumbaginaceae*). – *Taxon* **52**: 67–73.
- Lindberg H. 1946: Iter Cyprium. – *Acta Soc. Sci. Fenn., Ser. B, Opera Biol.* **2**(7).
- Mayer A. 1995: Comparative study of the coastal vegetation of Sardinia (Italy) and Crete (Greece) with respect to the effects of human influence. – *Libri Bot.* **15**.
- Meikle R. D. 1983: Addition to the flora of Cyprus. – *Ann. Mus. Goulandris* **6**: 87–94.
- Meikle R. D. 1985: *Flora of Cyprus* **2**. – Kew: Royal Botanic Gardens.
- Osorio-Tafall B. F. & Seraphim G. M. 1973: List of the vascular plants of Cyprus. – Nicosia: Ministry of Agriculture and Natural Resources.
- Palacios C. & Gonzalez-Candelas F. 1997: Lack of genetic variability in the rare and endangered *Limonium cavanillesii* (*Plumbaginaceae*) using RAPD markers. – *Molec. Ecol.* **6**: 671–675.
- Palacios C., Rosello, J. A. & Gonzalez-Candelas, F. 2000: Study of the evolutionary relationships among *Limonium* species (*Plumbaginaceae*) using nuclear and cytoplasmic molecular markers. – *Molec. Phylogen. Evol.* **14**: 232–249.
- Pignatti S. 1972: *Limonium*. – Pp. 38–50 in: Tutin T. G., Heywood V. H., Burges N. A., Moore D. M., Valentine D. H., Walters S. M. & Webb D. A. (ed.), *Flora europaea* **3**. – Cambridge: University Press.
- Rechinger K. H. 1943a: *Flora Aegaea. Flora der Inseln und Halbinseln des Ägäischen Meeres*. – *Akad. Wiss. Wien, Math.-Naturwiss. Kl., Denkschr.* **105**(1).
- Rechinger K. H. 1943b: *Neue Beiträge zur Flora von Kreta*. – *Akad. Wiss. Wien, Math.-Naturwiss. Kl., Denkschr.* **105**(2, 1).
- Sibthorp J. & Smith J. E. 1821: *Flora Graeca* **3**(2). – London: Richard Taylor et socii.
- Tsintides T., Christodoulou C. S., Delipetrou P. & Georgiou K. (ed.) 2007: *The Red Data Book of the flora of Cyprus*. – Lefkosia: Cyprus Forestry Association.
- Yıldız K. & Gücel S. 2006: Chromosome numbers of 16 endemic plant taxa from Northern Cyprus. – *Turk. J. Bot.* **30**: 181–182.